Original Article



In vitro and *in vivo* Production of Gingerols and Zingiberene in Ginger Plant (*Zingiber officinale* Roscoe)

Imaneh Dehghania, Akbar Mostajerana, Gholamreza Asgharib,*

^aPlant Science Division, Biology Department, University of Isfahan, Isfahan, I.R.Iran. ^bIsfahan Pharmaceutical Sciences Research Center, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, I.R.Iran.

Abstract

Ginger plant, Zingiber officinale Roscoe, is an important tropical plant used as spices and well known for its medicinal properties. It has a pungent and aromatic rhizome rich of zingiberen and gingerols. Many secondary metabolites are known to accumulate in the plant cell culture systems. So, in this research, gingerols and zingiberene production of callus cultures were compared with regenerated plant in vitro and glasshouse cultivated ginger in vivo. The experiment was conducted in a randomized treatment design with 4 replicates. Ginger rhizomes were imported from china and were cultured in dark at 28±2 °C for 30 days for sprouting and then were cut to small pieces (2-3 cm) and sterilized, and then the rhizomes were cut into 5 mm sized pieces. Bud explants were inoculated on different sterile medium at 25±2 °C under 16 L/8D photoperiod for 2 months. For in vivo culture, the ginger rhizome's surface was sterilized by using 1 g of l-1 of benomyle fungicide solution then was cut into small unique pieces and kept in perlite in control condition for 2 weeks for germination. Then unique sprouting seedlings were transferred to hydroponic system including Hoagland's solution for 3 months. Fresh calli, in vitro and in vivo gingers rhizomes were used for extraction with dichloromethane and analyzed by TLC. No gingerols and zingiberene spots on calli plates were detected, but in vitro and in vivo ginger rhizomes produced gingerols and zingiberene. Therefore, callus culture of ginger is proposed for rapid proliferation of plant cells.

Keywords: Gingerols; TLC; *Zingiber officinale*; Zingiberene. *Received:* December 23, 2010; *Accepted:* March 5, 2011

1. Introduction

Ginger plant, *Zingiber officinale* Roscoe, is an important medicinal horticultural crop in South East Asia and has a pungent, aromatic rhizome that is rich in secondary metabolites such as oleoresin, phenolic derivates, zingiberene and gingerols [1]. Ginger has been traditionally used for treatment of headaches, nausea, colds [2], arthritis, rheumatism, muscular discomfort [3], and inflammation [4]. Ginger therapeutic effects such as anti-inflammatory activities and

^{*}Corresponding author: Dr. Gholamreza Asghari, Department of Pharmacognosy, School of Pharmacy (http://pharm.mui.ac.ir/), Isfahan University of Medical Sciences, (http://www.mui.ac.ir), Hezar Jerib Ave. Isfahan, I.R. IRAN. Tel: +98-311-7922644; Fax: +98-311-6680011

I Dehghania et al / IJPS Spring 2011; 7(2): 129-133

moderate pungency have been attributed to the mixture of gingerol derivatives in oleoresins [5]. Recently, the production of secondary metabolites using plant cell and organ culture has been considered as a major research area. Micropropagation [6], organ culture [7], somatic embryogenesis [8], organogenesis [9], protoplast culture [10], and germplasm preservation [11] of ginger have been reported. Many compounds are known to accumulate in plant cell culture systems (in vitro), and their concentrations were equal to or higher than that of the plant (in vivo) [12]. Dornenburg and Knorr (1995) reported that cell culture have a higher rate of metabolism than intact differentiated plants because the initiation of cell growth in culture leads to a fast proliferation of the cell mass and to a condensed biosynthetic cycle [13].

It is reported that production of gingerol and zingiberene in callus culture of Z. *officinale* is correlated with some sort of differentiation [14]. Lack of information on in vitro and in vivo differentiation related to the secondary metabolite production in Zingiber officinale Rosc. was the reason for conducting this experiment. In this research, secondary metabolite (gingerols and zingiberene) production of callus culture (*in vitro*) and regenerated ginger has been compared with the ginger which was cultivated in greenhouse (*in vivo*).

2. Materials and methods

2.1. In vitro culture

Ginger rhizomes were obtained from China and then kept in dark at 28 ± 2 °C for 30 days for sprouting. The ginger rhizomes were cut into small pieces (2-3 cm) and sterilized in 2% NaClO solution with 4 drops of Tween 40 for 5 min, after tap washing and treatment with 70% ethanol for 30 sec. After a thorough wash in sterile distilled water, the rhizomes were surface sterilized with 0.2% (w/v) mercuric chloride solution enriched with few drops of 1 N HCl for 10-15 min. Following repeated washes with sterile water (4 washes, 3 min) the rhizomes were cut into 5 mm sized pieces having at least one bud. The outer sheaths covering buds were further removed [15]. The buds as explants were inoculated on a different sterile medium with 4 replicates [16].

2.2. Callus induction and plant regeneration

The basal Murashige and Skoog (MS) medium, containing 3% sucrose and 0.7% agar, pH 5.8, supplemented with 2,4-dichloro phenoxy acetic acid (2,4-D) and Kinetin (Kn) as plant growth regulators (PGRs) were used as the medium for different treatments (Table1). The buds of rhizomes in the medium were incubated for 2 months at 25 ± 2 °C under 16 L/8D photoperiod for callus establishment. The medium with 1 and 0.2 mg.l-1 of 2,4-D and Kn, respectively (M4), was used to obtain the highest percentage of callus induction. Each treatment resulted in callus formation accounted as positive results on callus induction. Then, the callus mass was also sub-cultured to hormone free medium M1 for 3 months in order to obtain shoot development and leaf emergence that lead to ginger regeneration [17].

2.3. In vivo culture

The ginger rhizome's surface was sterilized using 1 g/l of benomyle fungicide solution then cut into small pieces in such a way that each piece contained one vegetative bud. The rhizomes were planted in control condition in glasshouse (30 ± 5 °C, $55\pm5\%$ RH and 16 L/8D photoperiod) in small pots filled with perlite for germination. After 2 weeks, the unique sprouting seedlings were transferred to hydroponic system including Hoagland's solution for 3 months [18].

2.4. TLC analysis

Fresh calli, *in vitro* and *in vivo* ginger's rhizomes were cut into smaller pieces and macerated with dichloromethane (5 ml), at

M4

(MS) medium and supplements for callus induction.					
Medium	PGRs (mg/l)				
	2,4-D	Kn			
M1	-	-			
M2	1	-			
M3	-	0.2			

 Table 1. The treatments consisted of Murashige and Skoog

 (MS) medium and supplements for callus induction.

PGRs: plant growth regulators; 2,4-D: 2,4-dichloro phenoxy acetic acid; Kn: Kinetin

0.2

1

room temperature, then shaked for 20 min., filtered, and concentrated under nitrogen gas to give at least 1 ml of the extract, dehydrated by anhydrous sodium sulfate and chromatographed on a pre-coated TLC silica gel plate (silica gel G 060 F₂₅₄ plates). The TLC plates were put inside the development tank for the solvent to run up. N-hexane:diethyl ether (40:60 v/v) was used as developing solvent system. Then the plates were taken out before the solvent front reach to the top. The dried plates were then visualized by heating at 110 °C after spraying the reagent. The reagent used was sulfuric acid and vanillin. The individual zones were detected and identified in comparison with standards and the relevant literatures [19].

3. Results

3.1. Effect of PGRs on callus induction

The results indicated that different PGRs treatment has different effects on callus establishment. Using 0.2 mg/l of Kn had the lowest effect with 30%, but 1 mg of 2,4-D caused production of 50% of callus. The best results were obtained when the combination of 2,4-D and Kn were used. In this case 70% callus was produced (Table 2).

3.2. Effect of in vitro and in vivo on gingerols and zingiberene production

TLC analysis showed several spots with different R_f values in which gingerols and zingiberene spots were identified. No gingerols and zingiberene spots on calli was detected (Table 3), but two very apparent spots on plates related to in vitro regenerated and *in vivo* greenhouse cultivated ginger

Table 2. Percentage of callus induced in different mediums.				
Medium	PGRs (mgL ⁻¹) Callus induction			
	2,4-D	Kn		
M1		-		

 M2
 1
 50

 M3
 0.2
 30

 M4
 1
 0.2
 70

 PGRs: plant growth regulators; 2,4-D: 2,4-dichloro phenoxy acetic acid; Kn:
 Kinetin

rhizomes were observed. Gingerols, was revealed as a light violet spots which was equal to standard spot at Rf 0.3 and zingiberene as a dark purple spot at Rf 0.9. Indeed, results showed no more spots in regenerated in vitro ginger compare to *in vivo*.

4. Discusion

positive effect of high Despite concentration of 2,4-D on cell proliferation rate and consequently callus induction (70%) at M4 medium (Table1), no gingerols and zingiberene was detected in callus samples. Petiard et al. (1985) had reported that a number of physical and chemical factors could influence secondary metabolism in plant cell culture [20]. They concluded that the concentration of plant growth regulating hormones (PGRs) such as auxin (e.i 2,4-D) increases the cell growth, and higher concentrations of these hormones are deleterious to secondary metabolite production [20]. Similar results were reported by Ma and Gang (2006) on secondary metabolites formation in callus of ginger [21].

Indeed, the sites of secondary compounds synthesis and storage in plant cells often are located in individual or separated compartments such as highly specialized structures containing secretory and accumulatory elements and oil glands. In undifferentiated callus cultures, these accumulation sites are missing and probably due to the sites missing in calli, very low or no detectable yields of the secondary components were obtained [13]. According to Zarate and Yeoman (1994), the number of I Dehghania et al / IJPS Spring 2011; 7(2): 129-133

Table 3. Gingerols and zingiberene production on calli, in vitro regenerated, and in vivo mature ginger plant.

Physiological conditions	Morphological form	Gingerols	Zingiberene
In vitro	Callus	-	-
	Regenerated plant	+	+
In vivo	Mature plant	+	+

pigmented cells (cells containing the essential oils) is higher in mature rhizomes than in immature ones. Furthermore, they observed that there is a reasonable correlation between the number of pigmented cells and gingerols accumulation sites in the mature and immature rhizomes [22]. The finding of Zarate and Yeoman (1994) may support our results on no production of gingerols and zingibnerene in callus culture of ginger due to lack of morphological differentiation.

The amount and composition of secondary metabolites production in plant cells have been shown to be variable [23], and they often change due to physiological conditions [24]. Also, alterations in the environmental factors such as nutrient levels, light and temperature may affect the amount and composition of secondary metabolites [25]. Funk and Brodelius (1994) reported that higher amounts of N-sources in MS medium (*in vitro*) than in Hoagland's Solution (*in vivo*) play an important role in secondary products accumulation in ginger cells as well as in Vanilla planifolia [26]. Anasori and Asghari (2008) also reported that the presence of light is necessary for the gingerols and zingiberene production in differentiated Z. officinale [14]. So, in vitro and *in vivo* different physiological conditions such as nutrient solution components, light, temperature, pH, O₂ and CO2 uptake would caused differences between two environments and may influence gingerols and zingiberene production. However, no variation in composition of these two secondary metabolites was observed when different conditions were used in differentiation of Z. officinale in this experiment. It seems that in vitro conditions have not been effective on gingerols and zingiberene production resulting from variation of medium nutrients or morphological differentiation.

Acknowledgment

Authors would like to thank University of Isfahan and Isfahan University of Medical Sciences for supporting this work.

References

- [1] Bhagyalakshmi N, Singh NS. Meristem culture and micropropagation of a variety of ginger (*Zingiber officinale* Rosc.) with a high yield of oleoresin. *J Hortic Sci* 1988; 63: 321-7.
- [2] Grant KL, Lutz RB. Alternative therapies: ginger. *Am J Health Syst Pharm* 2000; 57: 945-7.
- [3] Blumenthal M, Werner BR. The complete German Commission E monographs: therapeutic guide to herbal medicines. Austin, Tex.: American Botanical Council; Boston, MA.: Integrative Medicine Communications. 1999.
- [4] Kiuchi F, Iwakami S, Shibuya M, Hanaoka F, Sankawa U. Inhibitors of prostaglandin biosynthesis from ginger. *Chem Pharm Bull* 1982; 30: 754-7.
- [5] Mustafa T, Srivastava KC, Jensen KB. Drug development report: 9. pharmacology of ginger, *Zingiber officinale. J Drug Dev* 1993; 6: 25-39.
- [6] Hosoki T, Sagawa Y. Clonal propagation of ginger (*Zingiber officinale* Rosc.) through tissue culture. *Hort Sci* 1977; 12: 451-2.
- [7] Sharma TR, Singh BM. High-frequency in vitro multiplication of disease-free *Zingiber officinale* Rosco. *Plant Cell Rep* 1997; 17: 68-72.
- [8] Kackar A, Bhat SR, Chandle KPS, Malik SK. Plant regeneration via somatic embrogenesis in ginger. *Plant Cell Tissue Organ Cult* 1993; 32: 289-92.
- [9] Babu KN, Samsudeen K, Ravindran PN. Direct regeneration of plantlets from immature inflorescence of ginger (*Zingiber officinale* Rosc.) by tissue culture. *J Spices Aromat Crops* 1992; 1: 43-8.
- [10] Geetha SP, Babu KN, Rema J. Isolation of protoplasts from cardamom (Elettaria cardamomum Maton.) and ginger (*Zingiber* officinale Rosc.). J Spices Aromat Crops 2000; 9: 23-30.
- [11] Dekker AJ, Rao AN, Gob CJ. In vitro storage of multiple shoot cultures of ginger at ambient temperatures of 24-29°C. *Sci Hort* 1991; 47: 157-67.

Gingerols and zingiberene production in cultured ginger plant

- [12] Werrmann U, Knorr D. Conversion of menthyl acetate or neomenthyl acetate to menthol or neomenthol by suspension cultures of Mentha canadensis and Mentha piperita. J Agric Food Chem1993; 41: 517-20.
- [13] Dornenburg H, Knorr D. Strategies for the improvement of secondary metabolite production in plant cell cultures. *Enzym Microb Technol* 1995; 17: 674-84.
- [14] Anasori P, Asghari G. Effects of light and differentiation on gingerol and zingiberene production in callus culture of *Zingiber officinale* Rosc. *Res Pharm Sci* 2008; 3: 59-63.
- [15] Chirangini P, Sharma GJ. In vitro propagation and microrhizome induction in Zingiber cassumunar (Roxb.) an antioxidant-rich medicinal plant. J Food Agricul Environ 2005; 3: 139-42.
- [16] Guo Y, Zhang Zh. Establishment and plant regeneration of somatic embryogenic cell suspension cultures of the *Zingiber officinale* Rosc. *Scientia Horticulturae* 2005; 107: 90-6.
- [17] Guo Y, Bai J, Zhang Zh. Plant regeneration from embryogenic suspension-derived protoplasts of ginger (*Zingiber officinale* Rosc.). *Plant Cell Tissue Organ Cult* 2007; 89: 151-7.
- [18] Dehghani I. Effect of salinity stress on growth rate (dry weight) and proline content in *Zingiber* officinale Roscoe plant. M.Sc. Thesis. Biology Department, University of Isfahan. 2008.
- [19] Wagner H, Bladt S, Zgainski EM. Plant drug analysis. New York, Springer-Verlag; 1984.

- [20] Petiard V, Baubault C, Bariaud A, Hutin M, and Courtois D. Studies variability of plant tissue cultures for alkaloid production in Catharanthus roseus and Papaver somniferum callus cultures. In: Primary and secondary metabolism of plant cell cultures (Neumann KH, Barz W, and Reinhard E, eds.).Springer-Verlage, New York, 1985; pp. 133-42.
- [21] Ma X, Gang DR. Metabolic profiling of in vitro micripropagated and conventionally greenhouse grown ginger (*Zingiber officinale*). *Phytochemistry* 2006; 67: 1673-85.
- [22]Zarate R, Yeoman MM. Studies of the cellular localization of the phenolic pungent principle of ginger, *Zingiber officinale* Roscoe. New Phytol 1994; 126: 295-300.
- [23] Parr AJ. Secondary products from plant cell cultures. Adv Biotechnol Proc 1988; 9: 1-34.
- [24] Pfeiffer N. FDA OKs Calgen's Flavr Savr TM tomato for marketing in supermarket. U.S. Genetic Engineering News 1994; 14: 1-31.
- [25] Stöckigt J. Biotransformations with cultivated plant cells. Agro Food-Industry Hi-Tech 1993; 4: 25-8.
- [26] Funk C, Brodelius PE. Vanilla planifolia: In vitro biosynthesis of vanillin and other phenylpropanoid derivatives. In: Biotechnology in agriculture and forestry, vol. 26. Medicinal and aromatic plants VI(Bajaj. Y. P.S. eds.). Springer-Verlag, Berlin, 1994; pp. 377-402.