Seasonal Influences on Different Stages of *In Vitro* Fertilization: Stimulation and Fertilization

Soraya Khafri, M.Sc.¹, Anoshirvan Kazemnejad, Ph.D.^{1*}, Mansoureh Movahedin, Ph.D.², Ebrahim Hajizadeh, Ph.D.¹

> 1. Biostatistics Department, Tarbiat Modares University, Tehran, Iran 2. Anatomy Department, Tarbiat Modares University, Tehran, Iran

Abstract.

Background: Effect of seasonal changes in human reproduction has been intensively researched. Some studies acknowledge influences of seasonal variation on natural conception, while others can not confirm them. The aim of this study was to investigate the effect of seasonal changes on different stages of assisted reproductive technology (ART) in Tehran.

Materials and Methods: This study was carried out on 268 cycles which were the first intracytoplasmic sperm injection (ICSI) cycle of the couples between July 2005 and March 2007. Multiple logistic regression was used to model the influence of different clinical factors on pregnancy rate and a bivariate Poisson regression model was used for modeling the number of mature oocytes and obtained embryos.

Results: Based on the logistic regression model, the duration of infertility, female age, number of mature oocytes, and transferred embryos were only the significant predictors. In this series, seasonal changes had no predictive value on the pregnancy rate.

Bivariate Poisson regression model showed that only the female age and duration of infertility were the factors simultaneously affecting the number of embryos and mature oocytes.

Conclusion: The suspected seasonal variability on the outcome of ICSI cycles has not been confirmed. Furthermore, among different seasons, there are no significant differences in the number of mature oocytes, obtained embryos, and transferred embryos.

Keywords: Oocyte, Embryos, ICSI, Pregnancy Rates, Seasonal Variation

Introduction

Every physician and biologist working in the field of reproductive medicine keeps wondering about variable pregnancy rates, because success rates are fluctuating without visible reason from excellent to very poor. Thus, the question of whether the different seasons of the year can have positive or negative influences on the characteristics of the oocytes, fertilization, or pregnancy rates in conjunction with assisted reproductive technologies (ART) in humans arises. Statisticians and epidemiologists have previously reported various in vitro fertilization (IVF) outcomes in correlation with or without seasonal and weather influences in different geographical areas. Some studies point to a variation in natural conception in humans (1-3); others cannot confirm the presence of the seasonal variations (4-7). Better sperm quality in the cooler seasons was observed by many studies (8), whereas other studies have been inconclusive (9, 10).

Different geographical areas with different climates may cause disparity in IVF outcomes. Thus, the aim

Received: 9 Aug 2008, Accepted: 6 Oct 2008

of this study was to investigate the effect of seasonal changes on various embryological parameters in IVF patients in Tehran.

Materials and Methods

Study sample

A database containing information on all ICSI cycles was used. In total, 306 cycles from 268 couples were performed between July 2005 and March 2007 in Shayamehr Limited Surgical Center (Tehran). The Research was approved by Medical Ethics Commitee of Tarbiat Modares University. This study was carried out on 268 cycles which were the first ICSI cycle of the couples. The season was expected to influence the oocyte during its development at the time of IVF treatment; it was also expected to influence the spermatozoon during their development before sperm collection on the day of oocyte aspiration. Therefore, to study seasonality, the date of aspiration was taken as an approximation of the moment of etiological impact.



Royan Institue International Journal of Fertility and Sterility Vol 2, No 2, Aug-Sep 2008, Pages: 90-95

^{*} Corresponding Address: P.O. Box 14115-111, Biostatistic Department, Tarbiat Modares University, Tehran, Iran Email: aklili@yahoo.com

The Iranian national regulations ensuring anonymity and privacy with respect to the patient data were followed. The ovarian stimulation protocol was as follows:

All of the patients in our study were underwent the Long protocol. All of them were down-regulated with the GnRH agonist and subsequently underwent controlled ovarian stimulation with gonadotrophins. When at least one follicle had reached a diameter of 18 mm and more than three had reached a diameter of greater than 15 mm, a single a single dose of 10,000 IU hCG via intramuscular injection was administered. Transvaginal ultrasound-guided follicular aspiration was performed 34-36 hours later. Oocytes were inseminated by the ICSI method 2-4 hours after follicular aspiration. According to embryo scoring, 1-3 embryos were transferred to each patient's uterus 48-72 hours post ICSI.

A clinical pregnancy was defined by the presence of at least one gestational sac in which a fetal heartbeat could be seen on transvaginal ultrasound examination 4 to 6 weeks after embryo transplantation.

Statistical Methods

One-way analysis of variance and chi-square were utilized to evaluate differences among the different seasons. A P value of less than 0.05 was considered to be statistically significant.

Multiple logistic regression was used to model the influence of the different clinical factors on pregnancy rate. All clinical, demographic, embryonic and temporal factors were entered into the analysis. Those factors were: the female age, duration of infertility, causes of infertility (male related, polycystic ovarian syndrome (PCOs), endometriosis, both male and female related, unexplained, etc.), season of treatment, abortion history, history of ectopic pregnancy, history of ovarian and/or uterine surgery, familial marriage, number of previous ART attempts, sperm characteristics (count, motility, and

morphology rates), and number of previous intrauterine inseminations (IUI). Weather parameters including: temperature, relative humidity, atmospheric pressure and cumulative daylight hours were acquired from the National Weather Bureau station of Tehran, Iran. Statistical analysis was performed using SPSS version 14.0 (SPSS Inc., Chicago, IL). Mature oocytes and obtained embryos are the outcomes of two important stages in IVF process: stimulation and fertilization. The number of mature oocytes and obtained embryos are likely to affect each other; hence their relationship must be analyzed with a simultaneous equation model. Because the number of mature oocytes and embryos are "counts" and they are correlated, the use of a standard method such as the standard linear regression or univariate Poisson regression was not thought to be suitable. Therefore, the bivariate Poisson regression model (11, 12) was alternatively chosen and adjusted to model the influence of seasonal variations on the number of mature oocytes, obtained embryos and other clinical factors. Because of the small sample size, the Bayesian approach was used for estimating the parameters and the Bayesian analysis was performed by using Winbugs v.1.4.3 (an open-source statistical computing environment described at http://www.mrc-bsu.cam.ac.uk/ bugs).

Results

The causes of infertility in the 268 cases which underwent ICSI cycles were male factor (38.1%), PCO (10%), endometriosis (3%), both male and female factor (12.3%), and Unexplained (8.2%). The mean duration of infertility and female age in our data were 8.53 4.83 and 31.59 5.32 years respectively.

Table 2 presents the numbers of mature oocytes, obtained embryos, transferred embryos and pregnancy rate in different seasons. However, the highest number of mature oocytes, obtained embryos, and

Tuble 1. Distribution of the cutical parameters of the patients in affected								
Clinical Parameters	Spring	Summer	Fall	Winter				
Number of Patients (%)	32 (11.9)	73 (28.0)	76 (28.4)	85 (31.7)				
Female Age	32.91±5.69	31.02±4.67	31.08±5.72	32.06±5.32				
Duration of Infertility	9.00 5.72	9.09±4.72	8.31±4.73	8.05±4.66				
Diagnosis of Infertility								
Male Factor	13.7	27.5	26.5	32.4				
PCO	15.4	8.5	30.8	42.3				
Endometriosis	7.7	15.4	23.1	53.8				
Both Male and Female Factor	9.1	9.1	33.3	48.5				
Unexplained	0	45.4	31.8	22.7				

Table 1: Distribution	of the clinical p	arameters of the	patients in	different seasons
-----------------------	-------------------	------------------	-------------	-------------------

Note: values are mean SD or percentages

Clinical Parameters (Mean±SD)	Spring	Summer	Fall	Winter	P-Value
Mature Oocytes	5.27±4.22	5.60±3.74	6.30±4.14	5.11±3.25	NS
Obtained Embryos	2.91 ± 2.82	3.40 ± 2.64	3.92 ± 2.92	3.18±2.36	NS
Transferred Embryos	2.09±1.33	2.32±1.16	2.47±1.22	2.52±1.32	NS
Pregnancy Rate (%)	25.0	22.7	10.5	15.3	NS

Table 2: The number of mature oocytes, obtained and transferred embryos, and pregnancy rate among different seasons

NS = Not Significant

rarely transferred embryos were in the fall but it was not statistically significant.



Fig 1: Pregnancy rate in different seasons in ICSI cycles

As shown in Fig 1, the highest pregnancy rate was in spring and the lowest in fall that was not statistically significant. Fig 2 shows the monthly weather parameter changes in Tehran. Maximum temperature was during the summer and was inversely related to the atmospheric pressure (2A and 2C). The relative humidity reached its peak from October to January (2B), and gradual declination of the cumulative daylight hours was observed from October to December (2D).

According to the logistic regression model on the outcome of pregnancy, among different clinical parameters only the duration of infertility, female age, number of mature oocytes, and transferred embryos were the significant predictors. Furthermore, seasonal changes had no predictive value on pregnancy rate in this series (Table 3).

The Bivariate Poisson regression model showed that only the female age and duration of infertility were



Fig 2: Monthly weather parameter changes

92

95% CI Parameters Estimation OR -0.040 Female age 0.954 0.714-00.982 **Duration of infertility** -0.055 0.946 0.785-00.971 The number of mature oocytes 0.131 1.140 1.006-01.291 The number of transferred embryos 0.462 1.590 1.022-64.300

Table 3: Variables selected as predictive of pregnancy by using multiple logistic regression

Table 4. Summar	v statistics a	of fitting t	he Rivariate	naissan regressian
<i>Iuvie</i> 4. Summur	ν διαπδιίζο υ	71 11111112 11	<i>ie Divariale</i>	DUISSUN TESTESSIUN

Variable definition	Mean	Std.Error	Lower	Upper			
Response variable: Mature Oocytes							
Constant	2.282	0.013	1.783	2.822 ª			
Female age	-0.017	0.004	-0.035	-0.004 ^a			
Indicator For season of treatment: 1=Fall, 0=otherwise	0.165	0.004	-0.021	0.348			
Familial marriage	0.091	0.002	-0.111	0.290			
Female history	0.146	0.003	-0.063	0.355			
Duration of infertility	-0.025	0.002	-0.044	-0.005 ^a			
Indicator for infertility diagnostic: 1=Tubal factor, 0=Male factor	-0.203	0.02	-0.381	0.027			
Indicator for previous ART attempts:1=2 twice, 0=1 twice	-0.105	0.003	-0.341	0.129			
Indicator for previous ART attempts:1=3 twice or more, 0=1 twice	0.258	0.003	-0.017	0.491			
History of ovarian problem	0.258	0.006	-0.132	0.636			
Response variable: Obtained embryos							
Constant	1.459	0.029	0.272	2.447 ^a			
Female age	-0.013	0.004	-0.035	-0.008 ^a			
Indicator for season of treatment: 1= Fall, 0= otherwise	0.163	0.004	-0.064	0.390			
Female history	0.121	0.004	-0.137	0.372			
Duration of infertility	-0.025	0.003	-0.049	-0.001ª			
Sperm motility	0.004	0.001	-0.004	0.012			
Sperm morphology	0.006	0.001	-0.007	0.009			

• "Lower" and "Upper" denote the 2.5th percentile and the 97.5th percentile of the simulated draws, respectively a: Significant at 0.05 level.

the factors simultaneously affecting the number of embryos and mature oocytes (Table 4).

Discussion

Carefully studying the literature on the seasonal variations in the outcome of ART cycles, the extent of discrepancies between these studies is remarkable. Some show the best pregnancy rates in the months of November thru February (13), others show the worst pregnancy rates in the months of January, February, and March (5) in IVF cycles. Consistent with the results of Rojansky et al (14) and Vahidi et al (15), we found an upper pregnancy rate in the spring and a lower rate in the fall (25.0% vs. 10.5%), but no statistical significances were observed. Our findings are in agreement with other studies showing no differences in the results of assisted reproductive techniques done in differ-

ent seasons of the year (4-7).

Better fertilization rates, embryo quality and pregnancy rates during the cold months have been observed (16, 13). In our study we showed no significant differences in the number of mature oocytes and embryos and transferred embryos among seasonal variations that this fact is disagreement with above-mentioned studies. However, maximum value of those variables is in the fall.

Melatonin, a hormone produced during darkness, may be one of the factors influencing gonadal function (17). The involvement of melatonin in the steroid genesis of human ovaries has been documented (18, 19). Webley and Luck (18) reported direct stimulation of melatonin on hCG. Melatonin is also thought to enhance hCG stimulation of progesterone production by human granulose cells in culture (19). Yie et al. (20, 21) reported higher concentrations of melatonin and progesterone during the dark season of the year and that of oestradiol during the light season in pre-ovulatory follicular fluids.

Rojansky et al. (14), showed the correlation between the absolute number of daylight hours and fertilization rate. Their result was highest in spring and lowest in autumn. Furthermore, Chang et al. (22) showed the cumulative daylight hours negatively correlated with the implantation and pregnancy rates of day 3 embryo transfers.

The higher pregnancy rate in spite of low oocyte and embryo numbers during spring may only be explained by increased endometrial receptivity during this season.

Fig 2 shows the maximum values for pressure and humidity, minimum as well as values for daylight hours and temperature in fall in Tehran, Iran. Furthermore, our results show the highest number of mature oocytes and obtained embryos in fall versus the other seasons. These results together confirm the conclusions of the previously mentioned studies.

Better sperm quality in the cooler seasons was observed by many authors (9). For those with severe poor sperm quality, normal fertilization could be easily achieved by ICSI. Thus, the impacts of poor sperm quality on ART outcome may be largely eliminated (23, 24). Therefore, the impact of the sperm quality was removed from the model in this study.

Multiple logistic regression showed that that the significant predictors influencing the outcome of ICSI cycle are the female age, duration of infertility, and the number of mature oocytes and transferred embryos.

Our results show that pregnancy rate increased with an increase in the number of transferred embryos and mature oocytes as well as with decreased patient age, and duration of infertility.

the significant predictors influencing the outcome of the ICSI cycle are the female age, duration of infertility, and the number of mature oocytes and transferred embryos.

When we used the bivariate Poisson regression model for studying the relationship between seasonal changes and two outcomes (mature oocytes and obtained embryos), we found no significant association. Our findings are in agreement with the studies showing no differences in the results of assisted reproductive techniques done in different seasons of the year (4-7). This may be due to the fact that seasonal changes should be taken into account together with other factors when evaluating infertility data (15). Furthermore, in addition to different ICSI variables, we found that the correlation factor between the two variables (the number of mature oocytes and embryos) should also be taken into account. More investigations using appropriate statistical modeling are needed in this regard.

This study has some limitations. We could not compare implantation rate among different seasons, therefore whether endometrial receptivity also played a role needs further investigations. The cross-sectional estimates of seasonal variation of hormone levels are not true measures of longitudinal changes, and individual variations cannot be identified. A longitudinal design following up the same participants throughout the year would have been preferable. However, true seasonal variations would be expected to be evident in this study due to a large sample size despite the cross-sectional design. Furthermore, the analysis included information from the first cycle which is likely to deviate from that in other cycles. In the next phase of our research, we will study the effect of the number of referring cycles on the outcomes in the presence of other factors. In addition, we will study the effect of different geographical areas with different climates on different embryological parameters with a nationwide multi-center study.

Conclusion

In conclusion, by the analysis of the first ICSI cycles, we confirmed that significant predictors for ICSI outcomes are the duration of infertility, female age, number of mature oocytes, and transferred embryos. However, the suspected seasonal variability of the outcome of ICSI cycles has not been confirmed. Furthermore, there are no significant differences in the numbers of mature oocytes, obtained embryos, and transferred embryos among different seasons.

A change in routine fertility treatments concerning different seasons should therefore not be considered.

Acknowledgements

The authors wish to thank the ART team of Shayamehr Limited Surgical Center, and Z. Alamalhoda for the Data collection.

The financial support by the Tarbiat Modares University for Scientific Research is greatly acknowledged.

References

1. Roenneberg T, Aschoff J. Annual rhythm of human reproduction. I. Biology, sociology or both? J Biol Rhythm. 1990; 5: 195-216.

2. Roenneberg T, Aschoff J. Annual rhythm of human reproduction. II. Environmental correlations. J Biol Rhythm. 1990; 5: 217-239.

3. Lam DA, Miron JA. Seasonality of birth in human popu-

Khafri et al.

lation. Soc Biol. 1991; 38: 51-78.

4. Daya I, Garcia JE, Smith RD, Padilla SL. Seasonal variation in in-vitro fertilization pregnancy rate: an analysis of 2,674 oocyte retrievals. Infertility. 1993; 15: 21-26.

5. Casper RF, Erskine HJ, Armstrong DT, Brown SE, Daniel SA, Graves GR, et al. In vitro fertilization: diurnal and seasonal variation in luteinizing hormone surge onset and pregnancy rate. Fertil Steril. 1988; 49: 644-648.

6. Fleming C, Nice L, Hughes AO, Hull MGR. Apparent lack of seasonal variation in implantation rates after in vitro fertilization. Hum Reprod 1994; 9: 2164-2166.

7. Wunder DM, Limoni C, Birkhäuser MH. the Swiss FIVNAT-Group. Lack of seasonal variations in fertilization, pregnancy and implantation rates in women undergoing IVF. Hum Reprod. 2005; 20(11): 3122-3129.

8. Centola GM, Eberly S. Seasonal variation and age related changes in human sperm count, motility, motion parameters, morphology, and white blood cell concentration. Fertil Steril. 1999; 72: 803-808.

9. Mortimer D, Tempelton AA, Lenton EA, Coleman RA. Annual pattern of human sperm production and semen quality. Arch Androl. 1983; 10: 1-5.

10. Ombelet W, Maes M, Vandeput H, Cox A, Janssen M, et al. Chronobiological uctuations in semen parameters with a constant abstinence period, Arch Androl. 1996; 37: 91-96.

11. Plassmann F, and Tideman TN. Confirming more guns, less crimes. Stanford Law Review 2003; 55(4): 1313-1370.

12. Thygesen HH, Zwinderman AH. Modeling Sage data with a truncated gamma-Poisson model. BMC Bioinformatics. 2006; 7: 157-165.

13. Stolwijk AM, Reuvers MJCM, Hamilton CJCM, Jongbloet PH, Hollanders JMG, Zeilhuis GA. Seasonality in the results of in-vitro fertilization. Hum Reprod. 1994; 2300-2305.

14. Rojansky N, Benshuhan A, Meirsdorf S, Lewin A, Laufer N, Safran A. Seasonal variability in fertilization and embryo quality rates in women undergoing IVF. Fertil Steril. 2000; 74: 476-481.

15. Vahidi A, Kalantar SM, Soleimani M, Amir Arjmand MH, Aflatoonian A, Karimzadeh MA, et al. The relationship between seasonal variability and pregnancy rates in women undergoing assisted reproductive technique, Iran. J Reprod Med. 2004; 2: 82-86.

16. Chang SY, Lan KC, Huang FJ, Kung FT, Tsai MY. Comparable clinical outcome of tubal embryo transfer for ligoasthenoteratozoospermia treated with intracytoplasmic sperm injection and for female infertility treated with in vitro fertilization. Chang Gung Med J. 2000; 23: 253-259.

17. Tamarkin L, Baird CJ, Almeida OFX. Melatonin: A coordinating signal for mammalian reproduction? Science. 1985; 227: 714-720.

18. Webley GE, Luck MR. Melatonin directly stimulates the secretion of progesterone by human and bovine granulosa cells in vitro. J Reprod Fertil. 1986; 78: 711-717.

19. Brzezinski A, Seibel MM, Lynch HJ, Deng MH, Wurtman RJ. Melatonin in human preovulatory fluid. J Clin Endocrinol Metab. 1987; 64: 865-867.

20. Yie SM, Niles LP, Younglai EV. Melatonin receptors on human granulosa cell membranes. J Clin Endocrinol Metab. 1995; 80: 1747-1749.

21. Yie SM, Brown GM, Liu GY, Collins JA, Daya S, Hughes EG. Melatonin and steroids in human preovulatory follicular fluid: Seasonal variations and granulosa cell steroid production. Hum Reprod. 1995; 10: 50-55.

22. Chang SY, Lan KC, Chen CY, Huang FJ, Tsai MY, Chang CY. The influences of weather on patients with different ovarian responses in the treatment of assisted reproductive technology. J Assist Reprod Genetics. 2005; 22(5): 191-198.

23. Lan KC, Chang SY, Huang FJ, Kung FT, Tsai MY. In vitro fertilization versus intracytoplasmic sperm injection in the embryo transfer model: a comparison of outcomes. Taiwanese J Obstet Gynecol. 2001; 40: 1-5.

24. Chamoun D, Udoff L, Scott L, Magder L, Adashi EY, McClamrock HD. Seasonal effect on pregnancy rates in an in vitro fertilization program. J Assist Reprod Genetics. 1995; 12: 585-589.