

GABA_A Receptor Subunits in Rat Testis and Sperm

Abolghasem Esmaeili, Ph.D.*

Cell, Molecular and Developmental Division, Biology Department, Faculty of Science,
The University of Isfahan, Isfahan, Iran

Abstract

Background: γ -Aminobutyric acid (GABA) is considered to be the predominant inhibitory neurotransmitter in mammalian central nervous systems (CNS). There are two major classes of GABA receptors: GABA_ARs and GABA_BRs. The GABA_A receptor is derived from various subunits such as α 1- α 6, β 1- β 3, γ 1- γ 4, δ , ϵ , π , and ρ 1-3. Intensive research has been performed to understand and establish the distribution and functions of these receptors in the CNS and peripheral tissues. The presence of some GABA_A receptors in sperm prompted us to explore the existence of GABA_A receptors in rat testis and sperm.

Materials and Methods: Total cellular RNA was isolated from Wistar rat sperm and testis and reverse transcribed to cDNA. PCR reactions were performed in a 20 μ l (microliter) volume containing specific GABA_AR subunits (forward and reverse primers) with other required materials. Reactions were carried out using a PCR machine to investigate the existence of GABA_A receptor subunits in rat testis and sperm. The amplification products were analyzed on 2% agarose gels stained with ethidium bromide.

Results: Our results showed that GABA_ARs composed of α 5, β 1, β 3, and γ 1 subunits were expressed in both testis and sperm. These results indicate that, in sperm, GABA_A receptors might have important functions.

Conclusion:

Sperm could be a novel site of GABA_A expression. Further studies should be taken to explore the role of these receptors on sperm acrosome reaction.

Keywords: GABA_AR, RT-PCR, Testis, Sperm

Introduction

The amino acid gamma-aminobutyric-acid (GABA) is a major inhibitory neurotransmitter in the CNS (1) that mediates most of its effects through GABA_A receptors. There are two major classes of GABA receptors: ionotropic GABA type-A receptors that form ion channels (GABA_ARs) and metabotropic GABA type-B (GABA_BRs) which are G protein-coupled receptors. The GABA_AR is a member of the cysteine-loop family of ligand-gated ion channels that also includes the nicotinic acetylcholine receptor cation channel (nAChR), the 5-HT₃ receptor cation channel (5-HT₃R) and the glycine receptor chloride channel (GABA_AR). Receptors of this class comprise 5 subunits arranged symmetrically around a central ion-conducting pore. Each subunit consists of 4 α -helical transmembrane (TM) domains and a large extracellular amino-terminal domain that harbors the ligand binding sites and the signature cysteine-loop.

Molecular biology revealed the complex subunit structural design of this receptor channel, in which a pentameric assembly resulting from five of at

least 21 subunits, grouped in the eight classes α 1-6, β 1-4, γ 1-4, δ , π , ϵ , θ , and ρ 1-3 (2, 3) permits an immense number of putative receptor isoforms. Their variety extended the existence of several splice forms, for example, the α 6, β 2 and γ 2 subunits (4). The subunit combination of a GABA_ARs determines the specific effects of allosterical modulators of benzodiazepines, barbiturates, steroids, general anaesthetics, some convulsants, polyvalent cations, and ethanol. These agents act in different binding sites, some of which are not yet known (5). Drugs and endogenous ligands bind either to the extracellular domain or channel domain of the GABA_ARS and act as positive or negative allosteric modulators (1). In heterologous expression systems, the presence of α and β classes are needed to have functional channels. These γ subunits are required in order to mimic the full repertoire of native receptor responses to drugs. Knowledge of the complex pharmacology of GABA_ARs might eventually enable site-directed drug design to further our understanding of

Received: 28 Sep 2008, Accepted: 20 Dec 2008

* Corresponding Address: P.O.Box: 817467354, Cell, Molecular and Developmental Division, Biology Department, Faculty of Science, The University of Isfahan, Isfahan, Iran
Email: aesmaeili@biol.ui.ac.ir



Royan Institute
International Journal of Fertility and Sterility
Vol 2, No 3, Nov-Dec 2008, Pages: 121-124

GABA-related disorders and of the complex interaction of excitatory and inhibitory mechanisms in neuronal processing.

GABA is a major inhibitory neurotransmitter in the mammalian brain. GABA_ARs are constructed from a family of around 21 different subunits including six alpha (α 1-6), four beta (β 1-4), four gamma (γ 1-4), one delta (δ), one epsilon (ϵ), one pi (π), one theta (θ), and three rho (ρ 1-3) subunits, all of which are products of separate genes.

GABA has been discovered in the epididymis, seminal vesicle, and testicle of the adult rat (6). Recently, it has been shown that GABA_ARs subunits are expressed outside of the CNS in the adrenal glands, ovaries, testis, placenta, and uterus in a tissue-specific manner (7). It is suggested that GABA might regulate sperm functions such as capacitation and acrosome reaction via its interaction with the receptors that were originally found in the CNS (8). In the present study, we demonstrate the presence of GABA_ARs subunits in rat testis as well as in sperm.

Materials and Methods

All experiments were performed on adult Wistar rats in accordance with the Biology Department for the Care and Use of Animals for Scientific Purposes and approved by the Animal Ethics Committee in the University of Isfahan. Animals were housed in cages with free access to standard rodent chow and water. The colony room was maintained at 22±2°C with a 12 hour: 12 hour light: dark cycle. Animals were anesthetized with halothane and decapitated. Testis were quickly isolated and kept ice-cold. Tissues from testis were excised. Sperm contained in the caudal epididymis of male rats were released into PBS and washed twice with PBS.

Total cellular RNA was extracted using the RNeasy kit (Qiagen, Valencia, CA) following the manufacturer's specifications. Total RNA was assessed by optical density measurements. DNase treated total RNA (2-3µg) was reverse transcribed using random hexamer primers and Superscript III RT invitrogen according to the manufacturer's instructions.

PCR reactions were performed in a 20µl volume containing 1U Taq (Fisher-Biotech), 1.5mM MgCl₂, 0.2mM dNTP and 10pM of specific GABA_AR subunits (Table 1) forward and reverse primers with 5 ng of template. Samples were made up to 20ml with RNase-, DNase-free water. Reactions were carried out using a BioRad (Bio-Rad Laboratories, Inc., California, USA) PCR machine as follows: an initial 95°C denaturation step for 30 seconds, annealing at 55°C for 30 seconds and extension at 72°C for 30 seconds, repeated

40 times. The amplification products were analyzed on 2% agarose gels stained with ethidium bromide. Products sizes were as expected for all subunits and specificity of the PCR products were confirmed by sequencing.

Results

RT-PCR in rat testis

The expression of 17 GABA_ARs subunits (Table 1) in rat testis assessed by RT-PCR analysis. Specific fragments from rat testis RT samples were obtained using selective primers. Products of corresponding size from rat brain RT samples via PCR with the same primers were also obtained. Subsequently, the amplified products were confirmed by sequencing analysis. It was found that the products from rat testis were identical to those from the rat brain. 18S cDNA served as a control of RT samples (Fig 1). In addition, no specific PCR product from the RNA samples that were omitted in reverse transcription could be observed over background, which verified the absence of genomic DNA contamination (Fig 1).




Fig 1: RT-PCR results of GABA_ARs. (A-C) PCR amplification of GABA_AR subunits in rat brain, sperm and testis respectively. M: marker; GABA_AR subunits are indicated in the figure.

The data suggested that α 5, β 1 α 5, β 1, β 3 and γ 1 subunits were always expressed, which provided evidence that those genes existed in rat testis.

Table 1. Primer sequences for real-time PCR of GABA_AR subunits

Submit	Direction	Primer sequence 5'-3'	Product size(bp)
$\alpha 1$	Forward	AGCTATACCCCTAACTTAGCCAGG	304
	Reverse	AGAAAGCGATTCTCAGTGCAGAGG	
$\alpha 2$	Forward	ACAAGAAGCCAGAGAACAACAAGCCAG	333
	Reverse	GAGGTCTACTGGTAAGCTCTACCAG	
$\alpha 3$	Forward	CAACATAGTGAACCACCTATCC	351
	Reverse	GGGGTTGGGGATTTGGATGCTTC	
$\alpha 4$	Forward	TGCCAGATCCAGAAGGTGGT	437
	Reverse	AAGATATCAAAACCTCCTCC	
$\alpha 5$	Forward	CAAGAAGGCCTTGAAGCAGCTAA	338
	Reverse	GGTTTCCTGTCTACTTTGGAGAG	
$\alpha 6$	Forward	TTCCTGGCTGCAAACTACTCGACA	348
	Reverse	AAGCCCCGGTAGCAAAGTCAAAA	
$\beta 1$	Forward	CCTGGAAATCAGGAATGAGACCAG	341
	Reverse	GGAGTCTAAACCGAACCATGAGAC	
$\beta 2$	Forward	TGAGATGCCACATCAGAAGC	317
	Reverse	TCATGGGAGGCTGGAGTTTAGTTC	
$\beta 3$	Forward	GAAATGAATGAGGTTGCAGGCAGC	355
	Reverse	GCAGGGTAATATTTCACTCAG	
$\gamma 1$	Forward	CAGAGACAGGAAGCTGAAAAGCAA	360
	Reverse	CGAAGTGATTATATTGGACTAAGC	
$\gamma 2$	Forward	TGTGAGCAACCGAAAACCAAGCAA	374
	Reverse	CGTGTGATTCAGCGAAGACCC	
$\gamma 3$	Forward	CATCCAGATTCAACAAGATG	255
	Reverse	AGCTCAGACGTCAATG	
δ	Forward	TGAGGAACGCCATTGTCTCTTCT	333
	Reverse	ACCACCGCACGTGGTACATGTAAA	
$\rho 1$	Forward	GAGAAAATTGACCAGTATA	348
	Reverse	GAGAAGATCTCTGCACCTGT	
$\rho 2$	Forward	GACAAATTTCCATGCACGTGT	210
	Reverse	GGAAAACACTGACCAATAAAT	
$\rho 3$	Forward	CACATATATACCCAGTAAA	350
	Reverse	GGAAAGATATCTGGAATGTA	
ϵ	Forward	CAAGTTAAGGCAAACAAGCCAGT	390
	Reverse	GTGACTTCATGGTCATGCGATTTC	
18S	Forward	GTAACCCGTTGAACCCATT	150
	Reverse	CCATCCAATCGGTAGTAGCG	

RT-PCR in rat sperm

Because the GABA_ARs were present in rat testis, we wondered if they also existed in rat sperm. Results revealed that $\alpha 5$, $\beta 1$, $\beta 2$, $\beta 3$ and $\gamma 1$ were detected in rat sperm (Fig 1). Testis-specific 18S proved the origin of RNA which was absent of genomic DNA contamination (Fig 1). The data suggested that GABA_ARs were always expressed providing evidence of the existence of those genes

in rat sperm.

Discussion

Several laboratories reported the presence of GABA receptors and transporters in testis and sperm (7, 9-11). An important function of GABA in the process of sperm maturation was demonstrated. GABA_ARs have been detected in pig, ram and human sperm membrane preparations by radiolig-

and-binding experiments (12, 13). It is suggested that GABA_AR subunits are localized primarily in the plasma membrane overlying the equatorial segment of motile or fixed, permeabilized capacitated and incapacitated ejaculated human sperm (14). The fact that immunoreactivity was also detected on motile sperm supports the expected plasma membrane location for the receptor. Western blot studies of epididymal rat sperm demonstrated the presence of the $\alpha 5$ or β ($\beta 1$ or $\beta 3$) subunits, and immunofluorescence experiments detected those subunits in the plasma membrane overlying the 'ventral surface' of the acrosome in fixed epididymal sperm (15).

Conclusion

In conclusion our results are in agreement with previous studies. Collectively these results show that sperm GABA_ARs may be involved in important fertilization events, such as depolarization as seen during progesterone-initiated AR capacitating and hyper-activated motility.

We propose that GABA_ARs can be therapeutically targeted for contraception or dysgenesis treatment. Additionally we are awaiting a deeper understanding of the mechanisms and functions of GABA_ARs in male reproductive tissues. We propose to study the expression of GABA_ARs in fertile and infertile male sperm. This may lead to a better understanding of infertility and may result in novel treatments.

Acknowledgments

There is no conflict of interest in this article.

References

1. Olsen RW, Tobin AJ. Molecular biology of GABAA receptors. *FASEB J.* 1990; 4(5):1469-1480.
2. Barnard EA, Skolnick P, Olsen RW, Mohler H, Sieghart W, Biggio G, et al. International Union of Pharmacology. XV. Subtypes of gamma-aminobutyric

acidA receptors: classification on the basis of subunit structure and receptor function. *Pharmacol Rev.* 1998; 50(2): 291-313.

3. Whiting PJ. The GABA-A receptor gene family: new targets for therapeutic intervention. *Neurochem Int.* 1999; 34(5): 387-390.

4. Korpi ER, Grunder G, Luddens H. Drug interactions at GABA(A) receptors. *Prog Neurobiol.* 2002; 67(2): 113-159.

5. Johnston GA. GABAA receptor pharmacology. *Pharmacol Ther.* 1996; 69(3): 173-198.

6. Erdö SL NL, Szporny L. The occurrence of GABA in vas deferens, prostate, epididymis, seminal vesicle and testicle of the rat. *Acta Biol Hung.* 1983; 34(4): 435-437.

7. Akinci MK, Schofield PR. Widespread expression of GABA(A) receptor subunits in peripheral tissues. *Neurosci Res.* 1999; 35(2): 145-153.

8. de las Heras MA VA, Perez LJ. In vitro capacitating effect of gamma-aminobutyric acid in ram spermatozoa. *Biol Reprod.* 1997; 56(4): 964-968.

9. Hu JH HX, Yan YC. Identification of gamma-aminobutyric acid transporter (GAT1) on the rat sperm. *Cell Res.* 2000; 10(1): 51-58.

10. Ma YH HJ, Zhou XG, Mei ZT, Fei J, Guo LH. Gamma-aminobutyric acid transporter (GAT1) overexpression in mouse affects the testicular morphology. *Cell Res.* 2000; 10(1): 59-69.

11. He XB HJ, Wu Q, Yan YC, Koide SS. Identification of GABA(B) receptor in rat testis and sperm. *Biochem Biophys Res Commun.* 2001; 283(1): 243-247.

12. Erdö SL WL. GABAA type binding sites on membranes of spermatozoa. *Life Sci.* 1990; 47(13): 1147-1151.

13. Ritta MN CJ, Bas DE. Occurrence of GABA and GABA receptors in human spermatozoa. *Mol Hum Reprod.* 1998; 4(8): 769-773.

14. Wistrom CA MS. Evidence suggesting involvement of a unique human sperm steroid receptor/Cl-channel complex in the progesterone-initiated acrosome reaction. *Dev Biol.* 1993; 159(2): 679-690.

15. Hu JH HX, Wu Q, Yan YC, Koide SS. Biphasic effect of GABA on rat sperm acrosome reaction: involvement of GABA(A) and GABA(B) receptors. *Arch Androl.* 2002; 48(5): 369-378.