

Correlation of Serum Lipid Profile with Histological and Seminal Parameters of Testis in The Goat

Ali Louei Monfared, Ph.D.

Department of Basic Sciences, Faculty of Veterinary Para-Medicine, Ilam University, Ilam, Iran

Abstract

Background: The lipid composition of a mammal's spermatozoa and seminal plasma vary in both structure and function. Evidence exists to suggest that dietary supplementation with the appropriate polyunsaturated fatty acids (PUFAs) affects spermatogenesis, semen quality and sperm motility. Therefore, this study has been conducted to evaluate the correlations between serum lipid profile and histological, anatomical and seminal parameters of testes in clinically healthy goats.

Materials and Methods: In this analytic, cross-sectional study, we chose a total of ten mature Iranian male goats that comprised a homogenous group through simple random sampling. Blood samples were taken from the jugular vein; the sera were separated and subsequently used for measurement of serum lipids, lipoproteins and testosterone levels. In addition, we collected semen from the animals and evaluated the seminal characteristics. We also performed histological and anatomical assessments of the testes.

Results: The findings demonstrated that serum levels of high density lipoprotein (HDL-c) had a significant positive correlation with interstitial testicular tissue area ($r=0.73$; $p<0.001$), seminiferous tubule area ($r=0.61$; $p<0.01$), the number of Leydig cells ($r=0.53$; $p<0.05$), the diameter of the Leydig cell nuclei ($r=0.54$; $p<0.05$), scrotal circumference ($r=0.83$; $p<0.001$), testis weight ($r=0.72$; $p<0.001$), the number of live, normal sperm ($r=0.94$; $p<0.001$) and serum testosterone levels ($r=0.88$; $p<0.001$). Significant but negative correlations were found between serum triglyceride concentration and seminiferous tubule area ($r=-0.53$; $p<0.05$), the diameter of the Leydig cell nuclei ($r=-0.55$; $p<0.05$), testis weight ($r=-0.64$; $p<0.01$), total sperm number ($r=-0.82$; $p<0.001$), number of live, normal sperm ($r=-0.55$; $p<0.05$) and serum testosterone levels ($r=-0.79$; $p<0.001$). In addition, a significant negative correlation was observed between serum very low density lipoprotein (VLDL-c) concentration and the percent of live sperm ($r=-0.67$; $p<0.01$), and serum testosterone levels ($r=-0.65$; $p<0.01$).

Conclusion: The present results indicated that among serum lipids only the levels of HDL-c positively correlated with testicular parameters. High serum triglyceride levels exerted direct adverse effects at the testicular level, which was mainly observed in the seminiferous tubules (STs), characterization of Leydig cells and semen quality.

Keywords: Lipid, Testis, Goat, Histology

Citation: Louei Monfared A. Correlation of serum lipid profile with histological and seminal parameters of testis in the goat. *Int J Fertil Steril.* 2013; 7(2): 122-129.



Introduction

It is well known that the lipid composition of a mammal's spermatozoa and seminal plasma differ in both structure and function. Evidence exists to suggest that dietary supplementation with appropriate polyunsaturated fatty acids (PUFAs) affects spermatogenesis, semen quality and sperm motility (1, 2). In addition, Esmaeili et al. (3) have reported that dietary fatty acid sources affect the PUFA composition of the ram's spermatozoa.

Lipoproteins transfer various lipids such as cholesterol from the tissue of origin to target sites, where the lipid complex is delivered via lipoprotein receptor-mediated uptake. Also, the supply of the Steroids required for cellular activities, including membrane formation, steroid hormone secretion, and the post-translational modification of proteins is regulated by lipoproteins (4). Lipoprotein-derived cholesterol is a major source of substrate for steroidogenic tissues, including the Leydig cells of some species (5-8).

The ratios of triglycerides and cholesterol to a polipoprotein B in the plasma are inversely related to testosterone levels (9). Also, plasma testosterone levels are related to the serum lipid profile in normal and infertile men (10). It has been suggested that high serum levels of cholesterol and triglyceride are associated with poor semen quality (11). In another study, Sebastian et al. reported that infertility might be associated with the changed lipid metabolism in seminal plasma in male individuals (12). Seminal parameters have been shown to be negatively correlated with serum very low density lipoprotein (VLDL-c) and total triglyceride values (13). Hypercholesterolemia and hypertriglyceridemia in rabbits possibly produce a decreased sperm capacity in an acrosome reaction (14). Cholesterol-enriched diets in rabbits have adverse effects on leydig and sertoli cells secretory functions, initiation and maintenance of spermatogenesis, sperm cytoskeleton formation, the sperm maturation process in the epididymis, and sperm ability to initiate embryonic development (15).

Although goats are economically a very important domestic mammal, there are very few

data in the literature concerning the testicular cell populations and endocrine secretion in this species (16). Because sperm lipids contain extremely high proportions of fatty acids (1), it is necessary to establish a link between serum lipid profile to both testicular structure and function. Accordingly, this study seeks to evaluate correlations between serum lipid profile and histological, anatomical and seminal parameters of testes in clinically healthy goats.

Materials and Methods

Experimental animals and design

In this analytic, cross-sectional study, a total of ten mature Iranian male goats that comprised a homogenous group (aged 28.7 ± 2.2 months and 43.7 ± 3.1 kg average body weight) were selected from local intensively managed goat flocks (Ilam, Iran) through simple random sampling. The experiments were carried out from late May to July 2009, at the research farm of Ilam University. All animals were in good health and of proven fertility since they were being used as sires in a specialized goat farm. All goats received a daily diet of 702 g alfalfa hay, 69.9 g barley and 141 g of a commercial concentrate. The diet contained 2.4 Mcal ME/kg, 12.1% crude protein, 1.1% calcium, and 0.31% phosphorus. All goats had free access to mineral blocks and fresh water. This unit was supervised by an experienced veterinarian and the prevention of internal and external parasites were a routine practice.

Blood sampling and analysis

Blood samples (15 ml) were taken from the jugular vein between 09:00 and 10:00 am after overnight fasting and collected in vacutainers (Becton Dickinson Co., USA). Serum was separated by centrifugation at 2500 rpm for 20 minutes and stored at -20°C until analysis. The sera were analyzed for total cholesterol and triglyceride levels using enzymatic kits (Pars Azmoon, Iran). Lipoproteins were isolated from sera by a combination of precipitation and ultracentrifugation. The precipitation method was used for measuring high density lipoprotein (HDL-c) particles. In this method, after addition of sodium phosphotungstate-magnesium to the serum, the non-HDL lipo-

proteins that sedimented by centrifugation (10000 g for 5 minutes) were aggregated. Then, the residual cholesterol was enzymatically measured (17, 18). Low density lipoprotein (LDL-c) was calculated as the difference between cholesterol measured in the precipitate and in the HDL-c fraction, while VLDL-c was estimated as one-fifth of the concentration of triglycerides (19). In addition, serum testosterone level was measured by a radioimmunoassay method that employed a diagnostic kit (Immunotech, SA, France, PI-1119).

Semen collection and quality evaluation

Semen was collected in the morning by using an artificial vagina and transported to the laboratory (at 37°C) within 10-15 minutes, then placed in a water bath at 37°C. All ejaculates were evaluated for volume, sperm concentration (photometer, IMV, L'Aigle, France), percent of live sperm (eosin-nigrosin method), and percentage of normal sperm in the stained smear (20).

Anatomical measurements

Scrotal circumference was measured using a tape-measure. The combined testis width, then testis length in the scrotum was determined by the use of a caliper (Mitutoyo, Kanagawa, Japan).

Histomorphometrical assessment

The right testis from each animal was removed from the body under local anesthesia. Before surgery, all animals received intramuscular (IM) injections of 1 ml of Rompum (Bayer) per 100 kg of body weight. All surgical procedures were performed by a licensed veterinarian and the obtained testes were trimmed out from the scrotum and epididymis. Then, testes were weighed and cut along the longitudinal axis to obtain transverse sections of the seminiferous tubules (STs). Tissue samples, 1-3 mm thick, were taken near the tunica albuginea. Sections of 4 µm thickness were stained with hematoxylin and eosin (H & E) and viewed by a light microscope (Nikon Eclipse E800). Fields with sections of STs in circular transverse section were selected at 100x magnification; photographs were taken with a digital camera (Coolpix 950, Nikon, China) and stored. Histometric and cytometric studies were conducted using Image

Tool® 3.0 software (UTHSCSA, San Antonio, USA). For this purpose, the boundary demarcation of STs was performed using the software and the area of the STs in each field was deducted from the total area of the field. Thus, the area of the STs and the area of the interstitial tissue were deducted. To measure the tubular diameter and the height of STs epithelium, at least thirty round or nearly round ST cross-sections were chosen at random and measured for each animal. In addition, the number of Leydig cells per 10³ mm of the interstitial area was counted in five fields from each animal, at 1000x magnification. Finally, the diameters of the Leydig cell nuclei were measured according to the Faridha et al. (21) method.

Statistical analysis

Results were presented as mean ± standard error of mean (SEM) for serum lipid profile, histological, anatomical and seminal parameters and the levels of serum testosterone. The data were statistically analyzed using the SPSS statistical software package, version 10.0.1 (SPSS Inc., Chicago, IL, USA). Prior to analysis, the Kolmogorov-Smirnova test was used to check normal distribution of the data. Then, to evaluate the correlation of the serum lipid profile with testicular cell populations and endocrine secretion, Pearson's partial correlation coefficient analyses were conducted. A p value <0.05 was considered statistically significant.

Results

Mean ± SEM and correlation coefficients between serum lipids and lipoproteins with histological measurements of testis in the Iranian male goat are presented in tables 1-3. In the present work, the level of serum HDL-c showed a positive, highly significant correlation with interstitial testicular tissue area ($r=0.73$; $p<0.001$) and STs area ($r=0.61$; $p<0.01$). There was a weak positive correlation of these attributes with the number of Leydig cells ($r=0.53$; $p<0.05$) and the nuclei diameters of the Leydig cells ($r=0.54$; $p<0.05$). In addition, significant but negative correlations were found between the level of triglycerides in the serum and STs area ($r=-0.53$; $p<0.05$) or diameter of the Leydig cell nuclei ($r=-0.55$; $p<0.05$).

Table 1: Mean \pm standard error (SEM) of serum lipids and lipoproteins in the Iranian male goat (n=10)

| Total cholesterol (mg/dl) | Total triglycerides (mg/dl) | Very low density lipoprotein (mg/dl) | Low density lipoprotein (LDL-c) (mg/dl) | High density lipoprotein (HDL-c) (mg/dl) |
|---------------------------|-----------------------------|--------------------------------------|---|--|
| 47.25 \pm 2.9 | 46.6 \pm 1.5 | 10.3 \pm 2.8 | 17.8 \pm 1.1 | 18.1 \pm 0.8 |

Table 2: Mean \pm standard error (SEM) of the histological measurements of testis in the Iranian male goat (n=10)

| Interstitial testicular tissue area (%) | Area of seminiferous tubules (STs) (%) | Height of STs epithelium (μ m) | STs diameter (μ m) | Number of Leydig cells ($\times 10^3$ mm area of the testis) | Diameter of Leydig cell nuclei (μ m) |
|---|--|-------------------------------------|-------------------------|---|---|
| 9.3 \pm 1.3 | 88.8 \pm 0.88 | 72.90 \pm 1.9 | 215.7 \pm 3.7 | 19.5 \pm 1.6 | 3.2 \pm 0.1 |

Table 3: Correlation coefficients between serum lipids and lipoproteins with histological measurements of testis in the Iranian male goat (n=10)

| Serum lipid profile/histological measurements of testes | Total cholesterol (mg/dl) | Total triglycerides (mg/dl) | Very low density lipoprotein (VLDL-c) (mg/dl) | Low density lipoprotein (LDL-c) (mg/dl) | High density lipoprotein (HDL-c) (mg/dl) |
|---|---------------------------|-----------------------------|---|---|--|
| Interstitial testicular tissue area (%) | 0.16 | 0.19 | 0.21 | 0.11 | 0.73*** |
| Seminiferous tubules (STs) area (%) | 0.23 | -0.53* | 0.18 | 0.18 | 0.61** |
| STs epithelium height (μ m) | 0.12 | 0.29 | -0.21 | 0.15 | 0.18 |
| STs diameter (μ m) | 0.11 | 0.13 | 0.25 | 0.12 | 0.15 |
| Number of Leydig cells ($\times 10^3$ mm area of testis) | 0.15 | 0.24 | 0.31 | -0.23 | 0.53* |
| Diameter of the Leydig cell nuclei (μ m) | 0.15 | -0.55* | 0.13 | 0.27 | 0.54* |

*, Significant at $p < 0.05$, **, Significant at $p < 0.01$ and ***; Significant at $p < 0.001$.

The mean \pm SEM of anatomical measurements of the testes, live body weight of the animals and their correlation coefficients with the serum lipid profile are presented in tables 4 and 5. Correlation coefficients of HDL-c with the scrotal circumfer-

ence ($r=0.83$; $p < 0.001$) and testis weight ($r=0.72$; $p < 0.001$) were highly significant. There were significant negative correlations observed between total serum triglyceride level and testis weight ($r=-0.64$; $p < 0.01$).

Table 4: Mean \pm standard error (SEM) of anatomical measurements of the testes and live body weight of the Iranian male goat (n=10)

| Scrotal circumference (cm) | Testis width (cm) | Testis length (cm) | Testis weight (g) | Live body weight (kg) |
|----------------------------|-------------------|--------------------|-------------------|-----------------------|
| 42.23 \pm 2.5 | 265.17 \pm 5.3 | 7.38 \pm 1.6 | 3.59 \pm 1.2 | 29.61 \pm 1.6 |

Table 5: Correlation coefficients between serum lipid profile with anatomical measurements of the testis and live body weight of Iranian male goat (n=10)

| Serum lipid profile/histological measurements of testes | Total cholesterol (mg/dl) | Total triglycerides (mg/dl) | Very low density lipoprotein (VLDL-c) (mg/dl) | Low density lipoprotein (LDL-c) (mg/dl) | High density lipoprotein (HDL-c) (mg/dl) |
|---|---------------------------|-----------------------------|---|---|--|
| Scrotal circumference (cm) | 0.09 | 0.04 | 0.02 | 0.21 | 0.83*** |
| Testis width (cm) | 0.06 | 0.01 | 0.03 | 0.02 | 0.11 |
| Testis length (cm) | 0.21 | 0.11 | 0.31 | 0.08 | 0.06 |
| Testis weight (g) | 0.07 | -0.64** | 0.38 | 0.12 | 0.72*** |
| Live body weight (kg) | 0.05 | 0.19 | 0.01 | 0.17 | 0.03 |

*, Significant at $p < 0.05$, **, Significant at $p < 0.01$ and ***, Significant at $p < 0.001$.

Mean \pm SEM of the semen characteristics and serum testosterone levels and their correlation coefficients with the serum lipid profiles are shown in tables 6 and 7. The correlation coefficients of HDL-c with the number of live, normal sperm ($r = 0.94$; $p < 0.001$) or serum testosterone levels ($r = 0.88$; $p < 0.001$) was highly significant. However there

were negative significant correlation coefficients between serum VLDL-c concentration and the percent of live sperm ($r = -0.67$; $p < 0.01$) or serum testosterone levels ($r = -0.65$; $p < 0.01$). Negative correlations were found between serum triglyceride concentration and total sperm number ($r = -0.82$; $p < 0.001$), number of live, normal sperm ($r = -0.55$; $p < 0.05$) or serum testosterone levels ($r = -0.79$; $p < 0.001$).

Table 6: Mean \pm standard error (SEM) of semen characteristics and serum testosterone concentration in the Iranian male goat (n=10)

| Semen volume (mL) | Sperm concentration (10^9 /mL) | Percent of live sperm (%) | Percent of abnormal sperm (%) | Total sperm number (10^9) | Number of live sperm (10^9) | Number of live and normal sperm (10^9) | Serum testosterone concentration (ng/ml) |
|-------------------|-----------------------------------|---------------------------|-------------------------------|-------------------------------|---------------------------------|--|--|
| 1.20 \pm 0.32 | 2.85 \pm 0.53 | 76.5 \pm 0.34 | 7.5 \pm 2.6 | 2.89 \pm 0.4 | 1.34 \pm 0.2 | 1.06 \pm 0.8 | 7.3 \pm 0.7 |

Table 7: Correlation coefficients between serum lipid profile with semen characteristics and serum testosterone concentration of the Iranian male goat (n=10)

| Serum lipid profile/ histological measurements of testes | Total cholesterol (mg/dl) | Total triglycerides (mg/dl) | Very low density lipoprotein (VLDL-c) (mg/dl) | Low density lipoprotein (LDL-c) (mg/dl) | High density lipoprotein (HDL-c) (mg/dl) |
|--|---------------------------------|-----------------------------------|---|---|--|
| Semen volume (ml) | 0.13 | 0.27 | 0.17 | 0.42 | 0.21 |
| Sperm concentration (10 ⁹ /ml) | 0.19 | 0.07 | 0.16 | 0.11 | 0.26 |
| Live sperm (%) | 0.04 | 0.07 | -0.67** | 0.03 | 0.29 |
| Abnormal sperm (%) | 0.37 | 0.15 | 0.22 | 0.05 | 0.39 |
| Total sperm number (10 ⁹) | 0.24 | -0.82*** | 0.08 | 0.09 | 0.33 |
| Number of live sperm (10 ⁹) | 0.06 | 0.09 | 0.34 | 0.07 | 0.19 |
| Number of live, normal sperm (10 ⁹) | 0.01 | -0.55* | 0.14 | 0.26 | 0.94*** |
| Serum testosterone concentration (ng/ml) | 0.13 | -0.79*** | -0.65** | 0.42 | 0.88*** |

*; Significant at $p < 0.05$, **; Significant at $p < 0.01$ and ***; Significant at $p < 0.001$.

Discussion

The findings demonstrated that among serum lipids, only the HDL-c levels had a positive correlation with testicular histomorphology, serum testosterone values, and seminal parameters. The steroidogenic capacity of Leydig cells in the goat's testis and its changes during development of cellular organelles including the mitochondria and smooth endoplasmic reticulum responsible for steroid biosynthesis has been mentioned previously (22, 23). Also, it has been suggested that increased serum testosterone concentration resulted from increased Leydig cell numbers and also the amount of smooth endoplasmic reticulum within the Leydig cells (24, 25).

The mechanism(s) of beneficial effects of HDL-c on testis are unknown but it seems that its cholesterol content is a major source of substrate for steroidogenic process in the Leydig cells of the testis. The main arguments for this case are as follows. In

human (26), equine (8) and rodents (5), HDL-c is a major substrate source for the steroidogenic process. Sertoli cells play an important role in delivering the necessary nutrients, including lipid substrates to the sperm throughout its differentiation. In addition, despite permeation of HDL-c across the blood-testis barrier and delivery of cholesterol to Sertoli cells, LDL-c and VLDL-c particles cannot pass through this barrier, thus their access to the germ cells is excluded (27). There is evidence that HDL-c is the major plasma lipoprotein in the pre-ruminant and ruminant (28) animals; on the other hands, the plasma levels of LDL-c and VLDL-c are minimal in this species. This phenomenon may be lead to utilize HDL-c for androgen synthesis by testicular cells.

Receptor-mediated endocytosis of HDL-c and the amount of its affinity for binding to the receptor sites have been reported by other researchers as a probable mechanism for its functions in cattle fertility (29, 30). In this regard, the high number

of HDL-c receptors in the testicular tissue may be mainly related to its role in the testosterone synthesis. Padron et al. (11) have reported positive correlations between serum HDL-c levels, sperm density and viability in males.

It has been shown that normal Sertoli and Leydig cell secretory functions play a key role in the initiation and maintenance of spermatogenesis (31), maintenance of the sperm maturation process in the epididymis and stimulation of optimal sperm movements such as rapid forward progression (32). Thus, in the present study, the significant positive correlation between serum HDL-c levels with the number of live and normal sperm, testosterone profile and testis weight might be attributed to normality of the secretory function of the Sertoli and Leydig cells that resulted in normal spermatogenesis and the epididymal sperm maturation process.

The present data demonstrated a significant negative correlation in serum levels of triglycerides and VLDL-c with total sperm number, percent of live sperm, number of live and normal sperm, and serum testosterone concentration. In this regard, Ergün et al. reported that increased serum total triglyceride and VLDL-c values had deleterious effects on spermatogenesis and were significantly correlated with decreased sperm motion characteristics (13). The authors suggested that compromised fertility might be the result of a decreased testosterone concentration secondary to impaired testicular function. These findings were comparable with the present results.

Additional studies are necessary to clarify whether there is a high concentration of VLDL-c in the interstitial testicular tissue, since harmful consequences within the STs may be secondary to the secretory deficiency of Leydig cells. Alternatively other compounds induced by high serum triglyceride levels may exert a direct detrimental action outside of the seminiferous tubule (i.e. on Leydig cells).

Conclusion

The present results seem to indicate that among serum lipids only the levels of HDL-c positively correlate with testicular parameters. The high serum triglyceride levels exert direct adverse effects at the testicular level, which are observed mainly in STs, Leydig cell characterization and semen

quality. Obviously, further studies are required to elucidate the underlying mechanism(s) and to identify the sites of lipids action on the goat testis.

Acknowledgements

The present research was financially supported by a grant from Ilam University (Ilam, Iran). The author expresses his appreciation to Drs. Reza Khani, Amir Parviz Salati and Ahmad Molla Bahrami for their assistance in revising the manuscript. The author is thankful for Mr. Abdollah Jaafarzadeh's assistance with animal collection. There is no conflict of interest in this article.

References

1. Speake BK, Surai PF, Rooke JA. Regulation of avian and mammalian sperm production by dietary manipulation. In: De Vriese SR, Christophe AB. editors. Male fertility and lipid metabolism. USA: AOCS Press; 2003; 96-117.
2. Gholami H, Chamani M, Towhidi A, Fazeli MH. Improvement of semen quality in holstein bulls during heat stress by dietary supplementation of omega-3 fatty acids. *Int J Fertil Steril*. 2011; 4(4): 160-167.
3. Esmaili V, Shahverdi A, Alizadeh AR, Alipour H, Towhidi A, Zarrabi M. Fatty acid profiles of ram's sperm after removing some fatty acid sources from the diets and persistency of fatty acids in sperm. *Int J Fertil Steril*. 2012; 5(4): 211-216.
4. Havel RJ, Kane JP. Structure and metabolism of plasma lipoproteins. In: Scriver CR, Beaudet AL, Valle D, Sly WS, Childs B, Kinzler KW, Vogelstein B. editors. The metabolic and molecular bases of inherited diseases, 8th ed. New York: McGraw-Hill; 2001; 2705-2716.
5. Freeman DA. Regulation of the cholesterol ester cycle of cultured Leydig tumor cells. *Eur J Biochem*. 1987; 164(2): 351-356.
6. Schreiber JR, Weinstein DB, Hsueh AJ. Lipoproteins stimulate androgen production by cultured rat testis cells. *J Steroid Biochem*. 1982; 16(1): 39-43.
7. Benahmed M, Dellamonica C, Haour F, Saez TM. Specific low density lipoprotein receptors in pig Leydig cells. Role of this lipoprotein in cultured Leydig cells steroidogenesis. *Biochem Biophys Res Commun*. 1981; 99(4): 1123-1130.
8. Eisenhauer KM, Roser JF. Effects of lipoprotein, equine luteinizing hormone equine follicle-stimulating hormone, and equine prolactin on equine testicular steroidogenesis in vitro. *J Androl*. 1995; 16(1): 18-27.
9. Perova NV, Gerasimova EN, Chernysheva NP, Nikitina NA, Shcherbakova IA. Change in the apoproteins of very low density lipoproteins in the blood in hypertriglyceridemia. *Vopr Med Khim*. 1979; 25(2): 185-192.
10. Mendoza SG, Osuna A, Zerpa A, Gartside PS, Glueck CJ. Hypertriglyceridemia and hypoalphalipoproteinaemia in azoospermic and oligospermic young men: relationships of endogenous testosterone to triglyceride and high density lipoprotein cholesterol metabolism. *Metabolism*. 1981; 30(5): 481-486.
11. Padron RS, Más J, Zamora R, Riverol F, Licea M, Mallea L, et al. Lipids and testicular function. *Int Urol Nephrol*. 1989; 21(5): 515-519.
12. Sebastian SM, Selvaraj S, Aruldas MM, Govindarajulu P. Patterns of neural and phospholipids in the semen of

- normospermic, oligospermic and azospermic men. *J Reprod Fertil.* 1987; 79(2): 373-378.
13. Ergün A, Köse SK, Aydos K, Ata A, Avci A. Correlation of seminal parameters with serum lipid profile and sex hormones. *Arch Androl.* 2007; 53(1): 21-23.
 14. Diaz-Fontdevilla M, Bustos-Obregon E. Cholesterol and polyunsaturated acid enriched diet: effects on kinetics of the acrosome reaction in rabbit spermatozoa. *Mol Reprod Dev.* 1993; 35(2): 176-180.
 15. Yamamoto Y, Shimamoto K, Sofikitis N, Miyagawa I. Effects of hypercholesterolaemia on Leydig and Sertoli cell secretory function and the overall sperm fertilizing capacity in the rabbit. *Hum Reprod.* 1999; 14 (6): 1516-1521.
 16. Leal MC, Becker-silva SC, Chiarini-Garcia H, Franca LR. Sertoli cell deficiency and daily sperm production in goats (*capra hircus*). *Anim Reprod.* 2004; 1: 122-128.
 17. Burtis CA, Ashwood ER. *Tietz textbook of clinical chemistry.* 2nd ed. USA: W.B. Saunders Co;1994; 508-525.
 18. Durrington PN. A comparison of three methods of measuring serum high density lipoprotein cholesterol in diabetics and non-diabetics. *Ann Clin Biochem.* 1980; 17(4): 199-204.
 19. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972; 18(6): 499-502.
 20. Sorensen AM. *Applied animal reproduction principles and practices.* New York: McGraw-Hill Book Co; 1979; 231-234.
 21. Faridha A, Faisal K, Akbarsha MA. Duration-dependent histopathological and histometric changes in the testis of aflatoxin-treated mice. *J Endocrinol Reprod.* 2006; 10(2): 117-133.
 22. Lunstra DD, Schanbacher BD. Testicular function and Leydig cell ultrastructure in long-term bilaterally cryptorchid rams. *Biol Reprod.* 1988; 38(1): 211-220.
 23. Lincoln GA. Seasonal aspects of testicular function. In: Burger H, de Kretser D. editors. *The testis.* New York: Raven Press; 1989; 329-386.
 24. Amann RP. Endocrine changes associated with onset of spermatogenesis in Holstein bulls. *J Dairy Sci.* 1983; 66(12): 2606-2622.
 25. Zirkin BR, Ewing LL, Kkromann N, Cochran RC. Testosterone secretion by rat, rabbit, guinea pig, dog and hamster tests perfused in vitro: correlation with Leydig cell ultrastructure. *Endocrinology.* 1980; 107(6): 1867-1874.
 26. Tureck RW, Strauss JF 3rd. Progesterone synthesis by luteinized human granulosa cells in culture: the role of de novo sterol synthesis and lipoprotein-carried sterol. *J Clin Endocrinol Metab.* 1982; 54(2): 367-373.
 27. Maboundou JC, Fofana M, Fresnel J, Bocquet J, Le Goff D. Effect of lipoproteins on cholesterol synthesis in rat Sertoli cells. *Biochem Cell Biol.* 1995; 73(1-2): 67-72.
 28. Forte TM, Bell-Qumt JJ, Cheng F. Lipoproteins of fetal and new born calves and adult steers: a study of developmental changes. *Lipids* 1981; 16(4):240-245.
 29. Mendel CM, Kunitake ST, Kane JP. Discrimination between subclasses of human high-density lipoproteins by the HDL binding sites of bovine liver. *Biochim Biophys Acta.* 1986; 875(1):59-68.
 30. Grummer RR, Carroll DJ. A review of lipoprotein cholesterol metabolism: importance to ovarian function. *J Anim Sci.* 1988; 66(12): 3160-3173.
 31. Steinberger E, Root A, Ficher M, Smith KD. The role of androgens in the initiation of spermatogenesis in man. *J Clin Endocrinol Metab.* 1973; 37(5): 746-751.
 32. Turner TT. On the epididymis and its function. *Invest Urol.* 1979; 16(5): 311-321.