

## Effects of *In Vitro* Zinc Sulphate Additive to The Semen Extender on Water Buffalo (*Bubalus bubalis*) Spermatozoa before and after Freezing

Kamran Dorostkar, D.V.M., Ph.D.<sup>1</sup>, Sayed Mortaza Alavi Shoushtari, D.V.Sc., M.Sc.<sup>2\*</sup>, Amir Khaki, D.V.M., Ph.D.<sup>2</sup>

1. Department of Clinical Sciences, Faculty of Veterinary Medicine, Azad University, Sanandaj Branch, Sanandaj, Iran

2. Division of Theriogenology, Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

### Abstract

**Background:** The objective of the study was to investigate the effects of *in vitro* zinc sulphate additive to semen extender on sperm parameters (progressive motility, viability, membrane integrity and DNA stability) after cryopreservation.

**Materials and Methods:** In this Prospective longitudinal laboratory study, semen samples of 5 buffalo bulls of 3-5 years old were collected at 5 different occasions from Iran, Urmia during summer and autumn 2011, 25 samples were used in each treatment. Sperm progressive motility, viability and abnormal morphology were measured before and at 0.5 ( $T_0$ ), 1( $T_1$ ) and 2( $T_2$ ) hours after diluting semen (1:10 v/v) in Tris-citric acid based extender (without egg yolk and glycerol) at 37°C containing none (control group), 0.072, 0.144, 0.288, 0.576 and 1.152 mg/L zinc sulphate to investigate dose and time effects. Next, a Tris-citric acid-egg yolk-glycerol extender (20% egg yolk and 7% glycerol) containing the same amount of zinc sulphate was prepared, diluted semen (1:10 v/v) was cooled and kept into a refrigerated chamber (4°C) for 4 hours to equilibrate. Sperm progressive motility, viability, abnormal morphology, membrane integrity and DNA damage were estimated. The equilibrated semen was loaded in 0.5 ml French straws and frozen in liquid nitrogen. Later, the frozen semen was thawed and the same parameters as well as total antioxidant capacity (TAC) of the frozen-thawed semen were determined.

**Results:** The results showed that zinc sulphate additive at the rate of 0.288 mg/L gave a higher protection of sperm progressive motility ( $53.7 \pm 1.8\%$  vs.  $40.5 \pm 1.7\%$ ), viability ( $70.8 \pm 1.8\%$  vs.  $60.1 \pm 1.5\%$ ), membrane integrity ( $67.3 \pm 1.6\%$  vs.  $56.6 \pm 1.7\%$ ), DNA stability ( $10.1 \pm 0.47\%$  vs.  $11.8 \pm 0.33\%$  damaged DNA) through the process of dilution, equilibration and freeze-thawing and caused a higher TAC level ( $81 \pm 3.3\%$  vs.  $63 \pm 3.2 \mu\text{mol/L}$ ) after freeze-thawing compared to the control group. Adding 0.576 and 1.152 mg/L zinc sulphate, however, was deleterious to the sperm and significantly reduced the studied sperm parameters.

**Conclusion:** Adding 0.288 mg/L zinc sulphate to the extender, compared to the control group, gives a better sperm preservation upon freezing processes which in turn, may results in higher semen fertility. But, addition of higher zinc sulphate concentrations (0.576 and 1.152 mg/L) are detrimental to buffalo spermatozoa.

**Keywords:** Semen, Zinc, Buffalo

**Citation:** Dorostkar K, Alavi Shoushtari SM, Khaki A. Effects of *in vitro* zinc sulphate additive to the semen extender on water buffalo (*bubalus bubalis*) spermatozoa before and after freezing . Int J Fertil Steril. 2014; 8(3): 325-332.

Received: 9 May 2012, Accepted: 21 Jul 2013

\* Corresponding Address: P.O.Box: 57153-1177, Division of Theriogenology, Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran  
Email: m.alavi@urmia.ac.ir



Royan Institute  
International Journal of Fertility and Sterility  
Vol 8, No 3, Oct-Dec 2014, Pages: 325-332

## Introduction

Semen cryopreservation is an important section of artificial insemination programs (1), it allows preservation of semen fertility for a long time. In this procedure, however, many spermatozoa lose their motility and other parameters which lead to a low fertility. A number of studies have demonstrated membrane lipid peroxidation as one of the causes of defective sperm function in liquid semen preserved at 4°C (2) and in cryopreserved semen (3, 4). Some attempts (5-7) have been made to preserve sperm parameters, particularly sperm motility, by adding some elements (zinc, copper and selenium) (8) and materials (vitamins, antioxidants, amino acids and coenzymes) to the semen before freezing.

Zinc is one of the important trace elements in the body, while its deficiency causes infertility in most animals due to disorders of testes development, spermatogenesis (9), steroidogenesis through gonadotropic hormones secretion (10), genetic expression of steroid receptors (11), testosterone synthesis and function through Zn-dependent metalloenzyme 5 $\alpha$ -reductase (12) and serum cholesterol level adjustment (13). The total zinc content in semen from mammals is high and zinc has been found to be critical to spermatogenesis (14, 15) and sperm concentration (16).

Zinc ions affect the expression of some germ-cell specific-genes during the course of spermatogenesis in sheep (17). Zinc supplementation leads to improved fertility in zinc deficient animals by increasing concentration and motility of spermatozoa (18) and sperm membrane integrity and reducing sperm DNA damage in human subjects (19). Zinc also contributes to the stability of sperm chromatin and repair of DNA damage (20). Mechanisms responsible for DNA strand break and changes in the histone to protamine ratio in DNA of zinc deficient men has been described in detail (20). Zinc influences the fluidity of lipids, and thus the stability of biological membranes. It is involved in the formation of free oxygen radicals and may play a regulatory role in the process of capacitation and the acrosome reaction (21), sperm nuclear chromatin condensation and acrosin activity (22). Zinc, at high concentrations in particular, may depress oxygen uptake (cell respiration) in the sperm cell and influence the sperm motility (23), zinc antioxidant capacity has been reported by many authors (24, 25). Alavi-Shoushtari et al.

observed that buffalo semen samples with higher seminal plasma zinc content ( $161.07 \pm 1.36$  vs.  $136.42 \pm 4.96$  mg/L) had higher sperm motility, viability as well as lower abnormal morphology (26) and total antioxidant capacity ( $1.57 \pm 0.01$  vs.  $1.23 \pm 0.05$  mmol/L) (27).

Information on the *in vitro* effects of zinc sulphate on buffalo bull spermatozoa is scarce. This prospective longitudinal laboratory study was conducted to investigate the effects of *in vitro* zinc supplementation on the progressive motility, viability, sperm membrane integrity and DNA stability of the spermatozoa in buffalo bulls' semen before and after semen equilibration and freezing with the aim of finding a practical protocol to have an improved semen quality after freeze-thawing.

## Materials and Methods

### *Semen collection and processing*

In this prospective longitudinal laboratory study, semen samples of 5 buffalo bulls of 3-5 years old kept in Buffalo Breeding Center of North-West of Iran, Urmia (37° 33' N, 45° 4' E), were collected by artificial vagina at 5 different occasions at weekly intervals during the late summer and autumn 2011. A total number of 25 samples were used in each examination. Semen samples were immediately transferred into a 37°C water bath after a physical (appearance, homogeneity, density, color and volume) examination. Sperm gross and progressive motility, viability, and abnormal morphology of semen were evaluated. Thin, watery specimens with low quality were discarded. These parameters (except gross motility) were measured within 0.5(T<sub>0</sub>), 1(T<sub>1</sub>) and 2(T<sub>2</sub>) hours after diluting semen (1:10 v/v) in the Tris-citric acid extender traditionally used in the center [(without egg yolk and glycerol), pH=7.1, osmotic pressure=300 mosmolkg<sup>-1</sup> and all the chemicals were purchased from Merck Co., Germany]. The extender consisting of Tris 2.66 g, glucose 1.2 g, citric acid 1.39 g, cysteine 0.139 g and distilled water in total volume of 100 ml that contained none (control, without zinc sulphate), 0.072, 0.144, 0.288, 0.576 and 1.152 mg/L zinc sulphate (ZnSO<sub>4</sub> · 7H<sub>2</sub>O, Scharlau Chemie S.A., Sentimental, Spain). Next, a Tris-citric acid-egg yolk-glycerol extender (above mentioned extender with 20% egg yolk and 7% glycerol, added in one step, as protectant and without antimicrobial agents) containing the same amounts of zinc

sulphate was prepared at room temperature. The semen samples were diluted at a rate of 1:10 v/v, (an approximate count of  $12 \times 10^6$  sperm cells/mL), cooled to 4°C within 2 hours, transferred to the equilibrium chamber of 4°C and left for 4 extra hours to equilibrate. Sperm parameters (progressive motility, viability, membrane integrity and DNA damage) were estimated in equilibrated semen and semen was then loaded in 0.5 ml French straws and frozen over liquid nitrogen in static vapor up to -120°C within 25 minutes before being plunged in the liquid nitrogen (28) and stored until further analysis. Later, the frozen semen was thawed in 37°C water bath for 30 seconds, and the same parameters, as well as total antioxidant capacity (TAC) of the frozen-thawed semen, were determined.

### ***Sperm functional assays***

#### ***Sperm progressive motility, viability and abnormal morphology***

Sperm progressive motility was estimated using a computer assisted system of analysis (CASA) [Hoshmand Fannavar (HF) CASA, version 6, Amirkabir Medical Engineering Co, Tehran]. After the samples were stained using eosin-nigrosin staining method (29), sperm viability was estimated by counting live and dead spermatozoa using a light microscope (Olympus BX41, Japan), at least 200 sperm cells were examined on each slide. Sperm cell abnormal morphology was also estimated.

#### ***Plasma membrane integrity***

Hypo-osmotic swelling test (HOST) was used to examine membrane integrity of spermatozoa before and after freezing according to a method of Jeyendran et al. (30) as described by El-Sisy et al. (31). In brief, the hypo-osmotic solution (osmotic pressure  $\equiv 190$  mosmol  $\text{kg}^{-1}$ , Osmomat 030, Nr. 951211, Gonotec, Germany) was prepared by dissolving 0.73 g sodium citrate and 1.35 g fructose in 100 ml of distilled water. Hypo-osmotic solution (500  $\mu\text{l}$ ) was mixed with 50  $\mu\text{l}$  of semen and incubated at 37°C for 40 minutes. After incubation, a drop of semen sample was examined under a phase-contrast microscope ( $\times 400$ ) (Olympus BX31, Olympus Optical Co., Japan) and 200 spermatozoa were counted in at least 5 different fields for their swelling characterized by coiled tail

indicating intact plasma membrane, which could be differentiated from abnormally coiled tails previously estimated.

#### ***Sperm DNA damage***

DNA damage was detected using acridine orange staining technique, according to the method of Katayose et al. (32). Briefly, first, spermatozoa were smeared on the glass slide. After being air dried, the samples were treated with acid alcohol (methyl alcohol-glacial acetic acid 3:1, vol/vol) for 2 hours. Immediately after air drying, approximately 1 mL working solution including 0.019% acridine orange [3, 6-bis (dimethylamino) acridine and hemi (zinc chloride) salt (Sigma Chemical Co., St. Louis, MO, USA)] was mounted on each glass slide for 5 minutes at room temperature and the samples were then washed with distilled water. The samples were observed under a fluorescence microscope (Model GS7, Nikon Co., Japan) immediately after placing a coverslip on the slide. A total of 100 to 200 spermatozoa were observed and classified, like green (intact DNA) or red (damaged DNA), based on differences in their fluorescent color.

#### ***Total antioxidant capacity (TAC)***

TAC of the frozen-thawed semen was assayed in duplicates using a commercial kit (Antioxidant Capacity Assay Kit, Cayman Chemical Co., Ann Arbor, MI, USA).

#### ***Statistical analysis***

The data obtained from 25 semen samples from 5 bulls (total number of 125 assays for each estimation) were analyzed using SAS (Version 9.2, SAS Institute Inc., Cary NC.). Prior to analyses, data were checked using box plots to examine for errors and outliers. No outlier was detected. The parameters were analyzed using a linear mixed model (PROC MIXED) with the REPEATED command. The Kenward-Roger procedure was used to approximate the denominator degrees of freedom. The residuals were assessed visually by quantile-quantile plots for testing of the normality and also predicted values were plotted versus residuals to assess the homogeneity of variance. To meet the assumptions of the tests, the data for all parameters, except than that for TAC, were transformed by applying square root transformation. The initial

model for each parameter included treatment, stage and their interactions as fixed effects and buffaloes nested in treatment were considered as random effect. When a non-significant interaction term was detected, the model was re-run with the interaction effect excluded from the model. For each parameter several covariance structure between repeated measures were examined and the model which provided better fit according to minimum Akaike's Information Criterion (AIC) was used. Differences between least squares means were determined using the DIFF option and Bonferroni's correction was applied to pair-wise comparisons. All reported values are least squares means and statistical mean and standard error of mean (SEM). Effects were declared significant at  $p < 0.05$ . This study was approved by the Ethic Committee of Urmia University.

## Results

### Freshly diluted semen

The results are summarized in tables 1 and 2. Adding 0.288 mg/L zinc sulphate to the extender preserved the sperm motility at  $T_1$  and  $T_2$  compared to controls ( $85.4 \pm 1.2\%$  vs.  $84.8 \pm 1.5\%$  at  $T_0$ ,  $85.5 \pm 1.1\%$  vs.  $80.8 \pm 1.3\%$  at  $T_1$  and  $85.7 \pm 1.1\%$  vs.  $77.4 \pm 1.5\%$  at  $T_2$ ) which was significant

( $p < 0.05$ ) at  $T_1$  and  $T_2$  indicating that the highest value belongs to only at  $T_2$ , whereas shows a significant effect on sperm viability ( $86.3 \pm 0.9\%$ ,  $p < 0.05$ , Table 1). Immediately after adding 0.576 and 1.152 mg/L zinc sulphate ( $T_0$ ) to the extender, the sperm motility (from  $84.8 \pm 1.5\%$  in control to  $80.7 \pm 1.7\%$  and  $75.2 \pm 1.2\%$  respectively) and viability (from  $86.1 \pm 1.2\%$  in control to  $83.5 \pm 1.6\%$  and  $77.8 \pm 1.2\%$  respectively) were significantly ( $p < 0.05$ ) decreased as compared with the controls and continued decreasing as time passed ( $p < 0.001$ , Table 1). Sperm abnormal morphology did not show significant variation.

### Equilibrated diluted semen

In equilibrated diluted semen the highest sperm progressive motility ( $80.7 \pm 1.9\%$ ), viability ( $84.1 \pm 1.0\%$ ) and sperm membrane integrity ( $83.3 \pm 0.9\%$ ) values were observed in 0.288 mg/L zinc sulphate, the number of intact DNA cells (green cells) in this diluted solution was not statistically different from control. Adding 1.152 mg/L zinc sulphate to the extender, however, significantly ( $p < 0.001$ ) reduced motility, viability and membrane integrity while increasing sperm cell DNA damage rate (Table 2).

**Table 1: Effects of different zinc sulphate concentrations (mg/L) on the motility and viability (LS mean  $\pm$  SEM) of spermatozoa in freshly diluted semen**

Parameter	Time	Treatment						SE	P value		
		Control	0.072	0.144	0.288	0.576	1.152		TRT	STG	TRT $\times$ STG
Motility	$T_0$	$84.8 \pm 1.5^{ab}$	$84.2 \pm 1.4^{ab}$	$85.0 \pm 1.8^{ab}$	$85.4 \pm 1.2^a$	$80.7 \pm 1.7^b$	$75.2 \pm 1.2^c$	0.0025	<0.0001	<0.0001	<0.0001
	$T_1$	$80.8 \pm 1.3^{ac}$	$81.0 \pm 1.4^{ac}$	$83.5 \pm 1.6^{ab}$	$85.5 \pm 1.1^b$	$78.2 \pm 1.7^c$	$71.7 \pm 1.2^d$	0.0025			
	$T_2$	$77.4 \pm 1.5^{***}$	$78.9 \pm 1.2^{bc**}$	$81.6 \pm 1.4^b$	$85.7 \pm 1.1^c$	$72.7 \pm 1.0^{d****}$	$67.9 \pm 1.2^{e****}$	0.0025			
Viability	$T_0$	$86.1 \pm 1.2^{ab}$	$86.3 \pm 1.2^{ab}$	$87.3 \pm 1.5^a$	$87.2 \pm 1.0^{ab}$	$83.5 \pm 1.6^b$	$77.8 \pm 1.2^c$	0.0016	<0.0001	<0.0001	0.0220
	$T_1$	$83.8 \pm 1.5^{ab}$	$84.2 \pm 1.2^a$	$85.6 \pm 1.4^a$	$86.8 \pm 1.0^a$	$80.2 \pm 1.6^b$	$74.0 \pm 1.3^{c*}$	0.0016			
	$T_2$	$81.7 \pm 1.1^{ac*}$	$80.2 \pm 1.4^{ac***}$	$83.5 \pm 1.2^{ab}$	$86.3 \pm 0.9^b$	$78.7 \pm 1.2^{c*}$	$71.7 \pm 0.9^{d****}$	0.0016			

$T_0$ : 0,  $T_1$ : 60,  $T_2$ : 120 minutes after diluting semen,  $a, b, c, d, e$ ; Signify a significant difference ( $p < 0.05$ ) within the same row, \*; Signifies a significant difference ( $p < 0.05$ ) with  $T_0$  values, \*\*; Signifies a significant difference ( $p < 0.01$ ) with  $T_0$  values, \*\*\*; Signifies a significant difference ( $p < 0.001$ ) with  $T_0$  values, +; Denotes a significant difference ( $p < 0.05$ ) with  $T_1$  values, LS; Least square, TRT; Treatment and STG; Stage.



**Table 2: Effect of different zinc sulphate concentrations (mg/L) on sperm parameters (LS mean  $\pm$  SEM) after equilibrium and thawing\* time**

Parameter	Stage	Treatment						SE	P value		
		Control	0.072	0.144	0.288	0.576	1.152		TRT	STG	TRT $\times$ STG
<b>Motility (%)</b>	EQ	71.4 $\pm$ 1.9 <sup>a</sup>	73.5 $\pm$ 1.8 <sup>ab</sup>	78.3 $\pm$ 1.6 <sup>bc</sup>	80.7 $\pm$ 1.9 <sup>c</sup>	64.1 $\pm$ 1.6 <sup>d</sup>	56.9 $\pm$ 1.3 <sup>e</sup>	0.0081	<0.0001	<0.0001	<0.0001
	TW	40.5 $\pm$ 1.7 <sup>a</sup>	43.2 $\pm$ 2.0 <sup>ab</sup>	47.8 $\pm$ 1.9 <sup>b</sup>	53.7 $\pm$ 1.8 <sup>c</sup>	32.9 $\pm$ 1.7 <sup>d</sup>	27.6 $\pm$ 1.6 <sup>e</sup>	0.0081			
<b>Viability (%)</b>	EQ	78.0 $\pm$ 1.5 <sup>a</sup>	79.1 $\pm$ 1.3 <sup>a</sup>	80.4 $\pm$ 1.2 <sup>a</sup>	84.1 $\pm$ 1.0 <sup>b</sup>	77.4 $\pm$ 1.1 <sup>a</sup>	73.5 $\pm$ 1.3 <sup>c</sup>	0.0036	<0.0001	<0.0001	<0.0001
	TW	60.1 $\pm$ 1.5 <sup>a</sup>	62.0 $\pm$ 1.6 <sup>ab</sup>	65.7 $\pm$ 1.8 <sup>b</sup>	70.8 $\pm$ 1.8 <sup>c</sup>	59.6 $\pm$ 1.8 <sup>a</sup>	53.7 $\pm$ 1.9 <sup>d</sup>	0.0036			
<b>Membrane Integrity (%)</b>	EQ	79.0 $\pm$ 1.2 <sup>ac</sup>	79.4 $\pm$ 1.3 <sup>ac</sup>	81.7 $\pm$ 0.9 <sup>ab</sup>	83.3 $\pm$ 0.9 <sup>b</sup>	77.8 $\pm$ 1.1 <sup>c</sup>	73.8 $\pm$ 1.5 <sup>d</sup>	0.0036	<0.0001	<0.0001	<0.0001
	TW	56.6 $\pm$ 1.7 <sup>a</sup>	58.9 $\pm$ 1.7 <sup>a</sup>	62.1 $\pm$ 1.8 <sup>b</sup>	67.3 $\pm$ 1.6 <sup>c</sup>	54.9 $\pm$ 1.7 <sup>a</sup>	48.8 $\pm$ 2.2 <sup>d</sup>	0.0036			
<b>Damaged DNA (%)</b>	EQ	3.0 $\pm$ 0.25 <sup>a</sup>	3.4 $\pm$ 0.21 <sup>a</sup>	3.4 $\pm$ 0.23 <sup>a</sup>	3.0 $\pm$ 0.25 <sup>a</sup>	3.5 $\pm$ 0.24 <sup>ab</sup>	4.14 $\pm$ 0.21 <sup>b</sup>	0.0025			
	TW	11.8 $\pm$ 0.33 <sup>a</sup>	11.6 $\pm$ 0.43 <sup>a</sup>	10.3 $\pm$ 0.49 <sup>b</sup>	10.1 $\pm$ 0.47 <sup>b</sup>	12.2 $\pm$ 0.33 <sup>ac</sup>	13.3 $\pm$ 0.26 <sup>c</sup>	0.0025	<0.0001	<0.0001	0.3084
<b>TAC (<math>\mu</math>mol/L)</b>	TW	63 $\pm$ 3.2 <sup>a</sup>	64 $\pm$ 3.7 <sup>a</sup>	77 $\pm$ 3.4 <sup>b</sup>	81 $\pm$ 3.3 <sup>b</sup>	57 $\pm$ 2.7 <sup>ac</sup>	49 $\pm$ 2.4 <sup>c</sup>				

\*; 30 seconds in a 37°C water bath, EQ; After equilibrium, TW; After thawing, TAC; Total antioxidant capacity, <sup>a, b, c, d, e</sup>; Signify a significant difference ( $p < 0.05$ ) within the same row, LS; Least square, TRT; Treatment and STG; Stage.

### Frozen-thawed semen

In frozen-thawed semen, the highest sperm motility (53.7  $\pm$  1.8%,  $p = 0.000$ ), viability (70.8  $\pm$  1.8%,  $p = 0.001$ ) and sperm membrane integrity (67.3  $\pm$  1.6%,  $p < 0.05$ ) values, were observed in 0.288 mg/L zinc sulphate, indicating that this concentration is significantly higher than the other treatments. Furthermore, the least number of DNA damaged cells ( $p < 0.01$ ) was obtained in this treatment (Table 2). Moreover, adding 0.144 and 0.288 mg/L zinc sulphate to the freezing extender, significantly ( $p < 0.05$ ) increased the total antioxidant capacity of the frozen-thawed semen as compared to the control (77  $\pm$  3.4 and 81  $\pm$  3.3  $\mu$ mol/L vs. 63  $\pm$  3.2  $\mu$ mol/L, respectively).

### Discussion

Sperm motility and fertility is reduced by sperm processing and cryopreservation involved in the semen preservation in artificial insemination programs (33-35), while the antioxidant defense capacity of the semen is also decreased (36-38). Furthermore, antioxidants supplementation of semen extenders during liquid storage or cryopreservation of the bull (5) and the boar (39) spermatozoa and its beneficial effects have been reported. Therefore, this study was designed to investigate the effects of different doses of zinc sulphate supplementation added to the extender on the quality of the sperm

during freezing.

The extender used in this study was Tris-citric acid base extender that had been used in the Buffalo Breeding Center for some years but the egg yolk, glycerol and antibiotics were not added to it in the first stage, in order not to interfere with the examination and in the view that they might have some effect on the sperm parameters. Andrabi (40) in a review of several studies concluded that Tris-citric acid might provide the best satisfactory buffering system to improve the post-thaw freezability and motion characteristics of buffalo spermatozoa. Akhter et al. (41) compared effects of soya lecithin based extender (Bioxcell®), Tris-citric acid egg yolk, egg yolk-citrate and skim milk extenders on buffalo spermatozoa stored at 5°C for 5 days and observed that motility, viability and plasma membrane integrity of buffalo bulls were similar in Tris-citric acid egg yolk, milk and Bioxcell on the first and third day of storage but on the fifth day they were better in Bioxcell. However, Rasul et al. (42) reported that Tris-citrate was better than Tris-citric acid, Tris-Tes and Tris-HEPES buffers in terms of improving post-thaw motion characteristics of buffalo spermatozoa.

In this study a CASA was used for the sperm progressive motility evaluation, but since the system was not calibrated for buffalo sperm, progressive motility of 25 semen samples were examined

visually coincident with the system evaluation and the results were compared. The difference of two readings was less than 3%. So, only this parameter of motility was used in the study. The other parameters of motility were not used because their reliability for buffalo semen was not assured.

Sperm progressive motility, viability and abnormal morphology were evaluated immediately after the semen being diluted with the extender containing different doses of zinc sulphate at different time intervals in order to assess the dose and time effect of zinc sulphate on spermatozoa quality. In all the examinations, all the treated samples were compared with their own control which had no zinc sulphate in its extender.

In selecting zinc sulphate dose two points were considered: 1. molecular weight (MW) of "zinc sulphate,  $7H_2O$ " is nearly 288 (287.57) grams with 65 gram zinc content, 2. Storage media, commercially prepared for *in vitro* fertilization (IVF) (Ham's F-10), usually contains 0.0288 mg/L (1:10000 MW) zinc sulphate,  $7H_2O$  in their composition (43). The dose of 0.288 mg/L (1:1000 MW) zinc sulphate,  $7H_2O$ , 10 times more than that in storage media, was selected as a base to create a zinc sulphate dose high enough to affect the spermatozoa; it was two times divided into halves [0.288:2(0.144) and 0.288:4(0.072) mg/L] or doubled 0.288×2(0.576) and 0.288×4 (1.152) mg/L for dose selection in this study to gain a reasonable range of zinc sulphate concentrations in the extender. This selection worked well in the study since low doses (0.072 and 0.144 mg/L) were not as effective as 0.288 mg/L dose and high doses (0.576 and 1.152 mg/L) showed detrimental effects.

The present study revealed that supplementation of zinc sulphate improved the quality of freshly diluted and frozen semen of buffalo bulls as compared to the non-supplemented control group. An earlier study on buffalo semen by Alavi-Shoushtari et al. (26) showed significant correlations between zinc content of seminal plasma and sperm parameters including progressive motility and viability, suggesting that zinc ions may be positively correlated with semen quality. Some authors have suggested that zinc, as an antioxidant agent, particularly as a co-factor of copper/zinc superoxide dismutase (Cu/Zn SOD), plays a major role in the protection of spermatozoa against peroxidative damages of and reactive oxygen species (ROS) (44). This is shown by adding Zn to the testicular cell

culture and increased enzymatic activity of Cu/Zn SOD (45). Our results revealed that adding zinc sulphate to semen extender improves semen TAC after freezing, in a dose dependant manner. These results were in consistent with the study of Omu et al. (19), who reported zinc therapy-associated improvement in sperm parameters includes an increase in the seminal antioxidant capacity and reduction of oxidative stress status. Alvarez and Storey (46) demonstrated that cryopreservation enhanced lipid peroxidation due to the loss of SOD activity during the freezing process. A better preserved sperm progressive motility, viability and membrane integrity after semen equilibration and freezing observed in this study could be attributed to increased antioxidant capacity of zinc ions, as reported by Hidiroglou et al. (47). Furthermore, this antioxidant activity might have been responsible for lower cases of DNA damage observed in this study. This could also be the effect of zinc antioxidant capacity on ROS released from the sperm mitochondria or on preventing production of lipid peroxidation products during freezing, which could cause DNA damage (DNA strand breakage) (20).

As most of sperm functions originated from the membrane, zinc may improve sperm functional capacity by creating a favorable environment (19, 48). In this study, the higher membrane integrity rate after adding 0.288 mg/L of the extender before and after semen freezing may be due to membrane stabilizing effect of zinc which is the result of interacting with some functional group of the intrinsic component of sperm membrane that has been reported by Sorensen et al. (14). We also observed a high percentage of membrane integrity, progressive motility and viability in test group of 0.288 mg/L, before and after freezing. As reported by Kumar et al. (49), sperm motility and viability are dependent upon membrane transport rate affected by zinc concentration.

A low percentage of motile sperms after adding high levels of zinc sulphate (0.576 and 1.152 mg/L) to the extender observed in this study could be due to an elevated free zinc fraction and its subsequent uptake by spermatozoa (50) and reduction of oxygen consumption, since high levels of zinc in semen impairs the oxygen consumption of sperms (14). This elevated free zinc fraction may be accounted for lower viability and membrane integrity rate and for higher

cases of DNA damage as recorded in this study.

In this study zinc is believed to be important for membrane and chromatin stability and sperm motility (47). The high sperm viability values obtained by adding 0.144 and 0.288 mg/L of zinc sulphate may be due to the membrane stabilizing action of zinc to prevent leakage of enzymes, proteins and other vital components of the sperm in order to extend the functional life of sperm (49). Betteger and O'Dell (51) have reported that zinc stabilizes ribosomes, lysosomes, DNA and RNA that result in survival and normal functioning of the sperm. They further found that zinc protects sperm from free radicals that induce damages through their scavenging properties.

Storage time affected quality of the freshly diluted semen by deteriorating sperm progressive motility and viability which was more evident in the control group and low doses of zinc sulphate supplementation groups, while adding 0.288 mg/ml zinc sulphate to the extender showed significantly better result. In equilibrated semen some spermatozoa did not tolerate the process. This was evident by lower semen quality (Table 2). Many sperms were damaged during the process of freezing which led to lower percentage of sperm progressive motility, viability, and membrane integrity, lower total antioxidant capacity and higher DNA damaged cells. Adding 0.288 mg/ml zinc sulphate to the extender significantly reduced the cell damage during freezing process, while the lower doses were not so effective and higher doses had adverse effect.

## Conclusion

The results showed that 0.288 mg/L zinc sulphate improve to the sperm quality (progressive motility, viability, membrane integrity and total antioxidant capacity) preservation upon freezing processes, however, we suggest a bigger sample population to achieve a definite statement. Our finding also revealed that zinc affects the cell membrane and leads to a lower degree of sperm DNA damage after semen freeze-thawing, which in turn, results in higher semen fertility. However, addition of higher zinc concentrations (0.576 and 1.152 mg/L) are detrimental to spermatozoa.

## Acknowledgements

This project was financially supported by the

grant of Research Department of the Urmia University, Urmia, Iran. We wish to thank the authorities and personnel of the Buffalo Breeding Center of North-West of Iran, Urmia, for their cooperation and providing semen samples and access to their facilities. We are grateful to Dr. B. Naghdeh for his help in data analysis and table lay outs. There is no conflict of interest in this study.

## References

1. Watson PF. The causes of reduced fertility with cryopreserved semen. *Anim Reprod Sci.* 2000; 60-61: 481-492.
2. Vishwanath R, Shannon P. Storage of bovine semen in liquid and frozen state. *Anim Reprod Sci.* 2000; 62(1-3): 23-53.
3. Chatterjee S, Gagnon C. Production of reactive oxygen species by spermatozoa undergoing cooling, freezing and thawing. *Mol Reprod Dev.* 2001; 59(4): 451-458.
4. Nair SJ, Brar AS, Ahuja CS, Sangha SPS, Chaudhary KC. A comparative study on lipid peroxidation, activities of antioxidant enzymes and viability of cattle and buffalo bull spermatozoa during storage at refrigeration temperature. *Anim Reprod Sci.* 2006; 96: 21-29.
5. Foote RH, Brockett CC, Kaproth MT. Motility and fertility of bull sperm in whole milk extender containing antioxidants. *Anim Reprod Sci.* 2002; 71(1-2): 13-23.
6. Agarwal A, Prabakaran SA, Said TM. Prevention of oxidative stress injury to sperm. *J Androl.* 2005; 26(6): 654-660.
7. Shiva Shankar Reddy N, Jagan Mohanarao G, Atreja SK. Effects of adding taurine and trehalose to a tris-based egg yolk extender on buffalo (*Bubalus bubalis*) sperm quality following cryopreservation. *Anim Reprod Sci.* 2010; 119(3-4): 183-190.
8. Dorostkar K, Alavi-Shoushtari SM, Mokarizadeh A. Effects of in vitro selenium addition to the semen extender on the spermatozoa characteristics before and after freezing in water buffaloes (*Bubalus bubalis*). *Vet Res Forum (VRF).* 2012; 3(4): 263-268.
9. Massanyi P, Trandzic J, Nad P, Korenekova B, Skalicka M, Toman R, et al. Concentration of copper, iron, zinc, cadmium, lead, and nickel in bull and ram semen and relation to the occurrence of pathological spermatozoa. *J Environ Sci Health A Tox Hazard Subst Environ Eng.* 2004; 39(11-12): 3005-3014.
10. Hurley WL, Doan RM. Recent developments in the roles of vitamins and minerals in reproduction. *J Dairy Sci.* 1989; 72(3): 784-804.
11. Freedman LP. Anatomy of the steroid receptor zinc finger region. *Endocr Rev.* 1992; 13(2): 129-145.
12. Prasad AS. Discovery of human zinc deficiency and studies in an experimental human model. *Am J Clin Nutr.* 1991; 53(2): 403-412.
13. Tajik J, Nazifi S. Correlation of serum lipids and lipoproteins with trace elements in water buffalo (*Bubalus bubalis*). *Vet Res Forum.* 2010; 1(2): 91-97.
14. Sorensen MB, Bergdahl IA, Hjollund NH, Bonde JP, Stoltenberg M, Ernst E. Zinc, magnesium and calcium in human seminal fluid: relations to other semen parameters and fertility. *Mol Hum Reprod.* 1999; 5(4): 331-337.
15. Slivkova J, Popelkova M, Massanyi P, Toporcerova S, Stawarz R, Formicki G, et al. Concentration of trace elements in human semen and relation to spermatozoa quality. *J Environ Sci Health A Tox Hazard Subst Environ Eng.* 2009; 44(4): 370-375.



16. Wong WY, Flik G, Groenen PM, Swinkels DW, Thomas CM, Copius-Peereboom JH, et al. The impact of calcium, magnesium, zinc, and copper in blood and seminal plasma on semen parameters in men. *Reprod Toxicol*. 2001; 15(2): 131-136.
17. Ghasemzadeh-Hasankolai M, Batavani R, Eslaminejad MB, Sedighi-Gilani M. Effects of zinc ions on differentiation of bone marrow-driven mesenchymal stem cells to male germ cells and some germ cell-specific gene expression in rams. *Biol Trace Elem Res*. 2012; 150(1-3): 137-146.
18. Cupic Z, Sinovec Z, Veselinovic S, Ivkov O, Veselinovic S, Medic D, et al., editors. The effect of dietary zinc, on semen quality in holstein-friesian bulls. *Proceedings of the 4<sup>th</sup> International Symposium on Animal Reproduction*; 1998; Ohrid, Macedonia.
19. Omu AE, Al-Azemi MK, Kehinde EO, Anim JT, Oriowo MA, Mathew TC. Indications of the mechanisms involved in improved sperm parameters by zinc therapy. *Med Princ Pract*. 2008; 17(2): 108-116.
20. Barratt CL, Aitken RJ, Bjorndahl L, Carrell DT, De Boer P, Kvist U, et al. Sperm DNA: organization, protection and vulnerability: from basic science to clinical applications-a position report. *Hum Reprod*. 2010; 25(4): 824-838.
21. Andrews JC, Nolan JP, Hammerstedt RH, Bavister BD. Role of zinc during hamster sperm capacitation. *Biol Reprod*. 1994; 51(6): 1238-1247.
22. Steven FS, Griffin MM, Chantler EN. Inhibition of human and bovine sperm acrosin by divalent metal ions. Possible role of zinc as a regulator of acrosin activity. *Int J Androl*. 1982; 5(4): 401-412.
23. Colagar AH, Marzony ET, Chaichi MJ. Zinc levels in seminal plasma are associated with sperm quality in fertile and infertile men. *Nutr Res*. 2009; 29(2): 82-88.
24. Powell SR. The antioxidant properties of zinc. *J Nutr*. 2000; 130 Suppl 5S: 1447S-1454S.
25. Dissanayake DMA, Wijesinghe PS, Ratnasooriya WD, Wimalasena S, Paliawadana TS. Effect of zinc levels in the sperm culture medium on sperm recovery and quality of sperms in the swim up procedure for sperm processing. *Ceylon J Med Sci*. 2006; 49: 21-27.
26. Alavi-Shoushtari SM, Rezaei SA, Ansari MH, Khaki A. Effects of the seminal plasma zinc content and catalase activity on the semen quality of water buffalo (*Bubalus bubalis*) bulls. *Pak J Biol Sci*. 2009; 12(2): 134-139.
27. Eghbali M, Alavi-Shoushtari SM, Asri-Rezaei S, Khadem Ansari MH. Calcium, magnesium and total antioxidant capacity (TAC) in seminal plasma of water buffalo (*Bubalus bubalis*) bull and their relationships with semen characteristics. *Vet Res Forum*. 2010; 1(1): 12-20.
28. Sukhato P, Thongsodseang S, Uthra A, Songsasen N. Effects of cooling and warming conditions on post-thawed motility and fertility of cryopreserved buffalo spermatozoa. *Anim Reprod Sci*. 2001; 67(1-2): 69-77.
29. Barth AD. Evaluation of potential breeding soundness of the Bull. In: Youngquist RS, Threlfall WR, editors. *Current therapy in large animal theriogenology*. 2<sup>nd</sup> ed. Philadelphia: Saunders Elsevier; 2007; 235-239.
30. Jeyendran RS, Van der Ven HH, Perez-Pelaez M, Crabo BG, Zaneveld LJ. Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to other semen characteristics. *J Reprod Fertil*. 1984; 70(1): 219-228.
31. El-Sisy GA, El-Sheshtawy RI, Mohamed AA, El-Nattat WS. Correlation between semen parameters and conception rate in buffaloes. *Glob Vet*. 2010; 5(1): 15-21.
32. Katayose H, Yanagida K, Hashimoto S, Yamada H, Sato A. Use of diamide-acridine orange fluorescence staining to detect aberrant protamination of human-ejaculated sperm nuclei. *Fertil Steril*. 2003; 79 Suppl 1: 670-676.
33. Roca J, Rodriguez MJ, Gil MA, Carvajal G, Garcia EM, Cuello C, et al. Survival and in vitro fertility of boar spermatozoa frozen in the presence of superoxide dismutase and/or catalase. *J Androl*. 2005; 26(1): 15-24.
34. Guthrie HD, Welch GR. Determination of intracellular reactive oxygen species and high mitochondrial membrane potential in percoll-treated viable boar sperm using fluorescence activated flow cytometry. *J Anim Sci*. 2006; 84(8): 2089-2100.
35. Wathes DC, Abayasekara DR, Aitken RJ. Polyunsaturated fatty acids in male and female reproduction. *Biol Reprod*. 2007; 77(2): 190-201.
36. Bilodeau JF, Chatterjee S, Sirard MA, Gagnon C. Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Mol Reprod Dev*. 2000; 55(3): 282-288.
37. Jiang ZL, Li QW, Li WY, Hu JH, Zhao HW, Zhang SS. Effect of low density lipoprotein on DNA integrity of freezing-thawing boar sperm by neutral comet assay. *Anim Reprod Sci*. 2007; 99(3-4): 401-407.
38. Kadirvel G, Kumar S, Kumaresan A. Lipid peroxidation, mitochondrial membrane potential and DNA integrity of spermatozoa in relation to intracellular reactive oxygen species in liquid and frozen-thawed buffalo semen. *Anim Reprod Sci*. 2009; 114(1-3): 125-134.
39. Funahashi H, Sano T. Select antioxidants improve the function of extended boar semen stored at 10 degrees C. *Theriogenology*. 2005; 63(6): 1605-1616.
40. Andrabi SM. Factors affecting the quality of cryopreserved buffalo (*Bubalus bubalis*) bull spermatozoa. *Reprod Domest Anim*. 2009; 44(3): 552-569.
41. Akhter S, Ansari MS, Rakha BA, Ullah N, Andrabi SM, Khalid M. In vitro evaluation of liquid-stored buffalo semen at 5°C diluted in soya lecithin based extender (Bioxcell®), tris-citric acid egg yolk, skim-milk and egg yolk-citrate extenders. *Reprod Domest Anim*. 2011; 46(1): 45-49.
42. Rasul Z, Anzar M, Jalali S, Ahmad N. Effect of buffering system on post-thaw motion characteristics, plasma membrane integrity and acrosome morphology of buffalo spermatozoa. *Anim Reprod Sci*. 2000; 59(1-2): 31-41.
43. Wright RW Jr, Bodioli KR. Aspects of in vitro fertilization and embryo culture in domestic animals. *J Anim Sci*. 1981; 53(3): 702-729.
44. Sikka SC. Relative impact of oxidative stress on male reproductive function. *Curr Med Chem*. 2001; 8(7): 851-862.
45. Celino FT, Yamaguchi S, Miura C, Ohta T, Tozawa Y, Iwai T, et al. Tolerance of spermatogonia to oxidative stress is due to high levels of Zn and Cu/Zn superoxide dismutase. *PLoS One*. 2011; 6(2): e16938.
46. Alvarez JG, Storey BT. Evidence for increased lipid peroxidative damage and loss of superoxide dismutase activity as a mode of sublethal cryodamage to human sperm during cryopreservation. *J Androl*. 1992; 13(3): 232-241.
47. Hidioglou M, Knipfel JE. Zinc in mammalian sperm: a review. *J Dairy Sci*. 1984; 67(6): 1147-1156.
48. Chia SE, Ong CN, Chua LH, Ho LM, Tay SK. Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men. *J Androl*. 2000; 21(1): 53-57.
49. Kumar N, Verma RP, Singha LP, Varshney VP, Dass RS. Effect of different levels and sources of zinc supplementation on quantitative and qualitative semen attributes and serum testosterone level in crossbred cattle (*Bos indicus* × *Bos taurus*) bulls. *Reprod Nutr Dev*. 2006; 46(6): 663-675.
50. Carpino A, Siciliano L, Petrone M, Stefano C, Aquila S, Ando S. Low seminal zinc bound to high molecular weight proteins in asthenozoospermic patients: evidence of increased sperm Zn content in oligoasthenozoospermic patients. *Hum Reprod*. 1998; 13: 111-114.
51. Bettger WJ, O'Dell B. A Critical physiological role of zinc in the structure and function of biomembranes. *Life Sci*. 1981; 28(13): 1425-1438.