

Assessment of *DPY19L2* Deletion in Familial and Non-Familial Individuals with Globozoospermia and *DPY19L2* Genotyping

Parastoo Modarres, M.Sc.¹, Somayeh Tanhaei, M.Sc.¹, Marziyeh Tavalaei, M.Sc.², Kamran Ghaedi, Ph.D.^{1,3*},
Mohammad Reza Deemeh, M.Sc.², Mohammad Hossein Nasr-Esfahani, Ph.D.^{1,2,4*}

1. Department of Cellular Biotechnology, Cell Science Research Center, Royan Institute for Biotechnology, ACECR, Isfahan, Iran
2. Department of Reproductive Biotechnology, Reproductive Biomedicine Research Center, Royan Institute for Biotechnology, ACECR, Isfahan, Iran
3. Department of Biology, Faculty of Sciences, University of Isfahan, Isfahan, Iran
4. Isfahan Fertility and Infertility Center, Isfahan, Iran

Abstract

Background: Globozoospermia is a rare syndrome with an incidence of less than 0.1% among infertile men. Researchers have recently identified a large deletion, about 200 kbp, encompassing the whole length of *DPY19L2* or mutations in *SPATA16* and *PICK1* genes associated with globozoospermia. The aim of this study was to analyze the *DPY19L2* gene deletion using polymerase chain reaction technique for the exons 1, 4- 8, 11 and 22 as well as break point (BP) “a” in globozoospermic men.

Materials and Methods: In this experimental study, genome samples were collected from 27 men with globozoospermia (cases) and 36 fertile individuals (controls), and genomic analysis was carried out on each sample.

Results: Deletion of *DPY19L2* gene accounted for 74% of individuals with globozoospermia. *DPY19L2* gene deletion was considered as the molecular pathogenic factor for the onset of globozoospermia in infertile men. By quantitative real-time polymerase chain reaction (qPCR), we genotyped *DPY19L2* deletion and identified carriers within the population.

Conclusion: This technique may be considered as a method for family counseling and has the potential to be used as a pre-implantation genetic diagnosis, especially in ethnic community with high rate of consanguineous marriages.

Keywords: Gene Expression, Genotyping, Globozoospermia

Citation: Modarres P, Tanhaei S, Tavalaei M, Ghaedi K, Deemeh MR, Nasr-Esfahani MH. Assessment of *DPY19L2* deletion in familial and non-familial individuals with globozoospermia and *DPY19L2* genotyping. *Int J Fertil Steril*. 2016; 10(2): 196-207.

Introduction

Globozoospermia is a rare autosomal recessive genetic syndrome with an incidence of less than 0.1%. In this syndrome, due to defect in the process of acrosome biogenesis, the sperm contains a round head shape, consequently leading to no penetration into the oocyte during fertilization. Thus, direct intracytoplasmic sperm insemination (ICSI) along

with artificial oocyte activation is the only solution to gain pregnancy at couples suffering this abnormality (1). Genetic pedigree assessment of these individuals indicates the congenital origin of globozoospermia. To define molecular defects involved in this disorder, several autosomal genes have been identified in knockout mice models including: *Csnk2a2*, *Hrb*, *Gopc*, *Pick1*, *Gba2*, *Vps54*, *Zpbp1* and

Received: 27 Sep 2015, Accepted: 25 Nov 2015

*Corresponding Addresses: Department of Cellular Biotechnology, Cell Science Research Center, Royan Institute for Biotechnology, ACECR, Isfahan, Iran
Department of Reproductive Biotechnology, Reproductive Biomedicine Research Center, Royan Institute for Biotechnology, ACECR, Isfahan, Iran
Emails: kamranghaedi@royaninstitute.org, mh.nasr-esfahani@royaninstitute.org



Royan Institute
International Journal of Fertility and Sterility
Vol 10, No 2, Jul-Sep 2016, Pages: 196-207

Hsp90b1 (2-9). Defect of these genes in mouse models represented phenotypically similar abnormalities to human globozoospermia. However, among the aforementioned genes, only *PICK1* mutation was yet detected in human. *PICK1* protein is involved in subcellular trafficking in brain, pancreas and testis. The respective gene is located on human chromosome 22, and contains 13 exons. In spermatogenesis, *PICK1* is involved in trafficking of pro-acrosomal vesicles from golgi apparatus to acrosome. Liu et al. (10) showed a homozygous missense mutation (G198A) at the C-terminal domain of *PICK1* which disrupted *PvuII* site, culminating in formation of sperms with round head shape in human. Other human autosomal genes involved in globozoospermia are *SPATA16* and *DPY19L2* (11-13). *SPATA16* is a testis specific gene, translating a protein which is localized in the golgi apparatus and plays a role in the transportation of pro-acrosomal vesicles from golgi to the acrosome in the round and elongated spermatids (14). Dam et al. (11) found a homozygous sequence variation in the last nucleotide of exon 4 (G848A) of this gene which impaired *NciI* or *HpaII* recognition site, in three infertile brothers of a Jewish family with globozoospermia.

However, the most likely considered gene to have a pivotal role in globozoospermia is *DPY19L2*. This gene is expressed primarily in spermatids with a specific localization limited to the inner nuclear membrane, facing the acrosomal vesicle. Lack of the relevant protein causes instability of acrosome vesicles and thereby loss of acrosome (15). It has been demonstrated that complete deletion of *DPY19L2* by non-allelic homologous recombination (NAHR) results in globozoospermia (12, 13). Recent studies have revealed that *DPY19L2* gene function could be eliminated at nine possible breakpoints covering three regions, known as breakpoints "a, b and c" in two low copy flanking repeats (LCRs) of *DPY19L2* gene. High incidence (96.5%) of LCR sequences facilitates the occurrence of NAHR in this region (16).

Considering the role of aforementioned genes in globozoospermia and in line with our previous study (16), the aim of this study was to

evaluate the prevalence of missense mutations, G848A, in exon 4 of *SPATA16* gene and G198A in exon 13 of *PICK1*, as well as *DPY19L2* deletion in Iranian infertile individuals with globozoospermia referring to Isfahan fertility and infertility center (IFIC). Herein, we observed complete deletion of *DPY19L2* gene in 20 out of 27 globozoospermic individuals, but no mutation was detected in *SPATA16* or *PICK1* gene. We also performed quantitative real-time polymerase chain reaction (qPCR) assay to identify individuals with homo/hemizygous deletion of *DPY19L2* gene.

Materials and Methods

Mutational analysis of *SPATA16*, *PICK1* and *DPY19L2* genes

This experimental study was approved by Institutional Review Board (IRB) of Royan Institute. In this case-control study, 27 male with globozoospermia from Iranian population were contributed. An arbitrary number was assigned to each globozoospermic individual (G1 to G29), out of whole two individuals, G11 and G17, were omitted due to missing. We assessed the mutations for *SPATA16* and *PICK1* genes and provided pedigrees for two families with complete deletion of *DPY19L2* and one family with deletion of exon 5, 6 and 7 in *DPY19L2* gene.

In this process, blood samples were taken from 27 individuals, with globozoospermia with round-headed spermatozoa who referred to IFIC, as well as their family members after completing a consent form. Two out of 27 persons with more than 50% acrosomeless spermatozoa in their normal and round-headed sperm samples were considered to have partial globozoospermia, while the rest of individuals were suffering from total globozoospermia. Peripheral blood samples were also taken from 30 fertile men as well as the parents of three individuals with globozoospermia (G8, 14 and 21). In the sample group, except three brothers (G21, 22, and 23) and two cases of five (G5, 6, 20, 26, and 27) and two (G9 and 29) cousin subjects, the remaining 17 individuals with globozoospermia belonged to unrelated families (Table 1).

Table 1: Features of 27 individuals with globozoospermia

Patient	Type of globozoospermia		Consanguinity	Deficiency in <i>DPY19L2</i> gene			Reference	
	Complete	Partial		No deletion	Complete deletion			Partial deletion (exons 5, 6, 7)
					Unknown break point	Break point "a"		
G1, 4, 7, 12, 13, 15	✓		Non-familial		✓		(16)	
G2, 8, 10, 14, 16	✓		Non-familial			✓	(16)	
G3, 18	✓		Non-familial	✓			(16)	
G19		✓	Non-familial	✓			(16)	
G5, 6, 20	✓		Familial			✓	(16)	
G9	✓		Familial (G29)			✓	(16)	
G21, 22, 23	✓		Familial			✓	Current study	
G24	✓		Non-familial		✓		Current study	
G25, 28		✓	Non-familial	✓			Current study	
G26, 27	✓		Familial (G5)			✓	Current study	
G29	✓		Familial (G9)			✓	Current study	

Genomic DNA was extracted from individuals' peripheral blood samples using standard salting out procedure and kept at -20°C until usage (17). Specific primers for identification of G848A, in exon 4 of *SPATA16* gene and G198A in exon 13 of *PICK1* gene were designed by oligo7 primer designing software (Molecular Biology Insights, CO, USA) according to the respective sequences obtained from National Center for Biotechnology Information (NCBI) database, whereas primer sequences (Table 2) for assessment of *DPY19L2* deletion were ordered according to previous report (16). Missense mutations of *SPATA16* and *PICK1* genes were assessed using Restriction Fragment Length Polymorphism PCR (RFLP-PCR) assay, due to ability of their PCR products digestion by *NciI* and *PvuII* restriction enzymes, respectively. Indeed, G848A nucleotide variation in *SPATA16* gene causes disruption of *NciI* site in this location. Thus, a partial PCR product (635 bp) of this gene encompassing G848A could not be cut to produce 283 and 352 bp fragments. Similarly, mutation of G198A region in *PICK1* gene disrupts one of two *PvuII* restric-

tion sites located in this 548 bp PCR product. Thus, G198A mutation produces two bands after *PvuII* cut, lack of which could cause production of three bands after *PvuII* digestion. In this study, we did not evaluate the other mutations in these two genes.

Following identification of three exons (5, 6 and 7) deletion in one of the affected Iranian individual (G9) which was previously reported by Elinati et al. (16), and due to the history of infertility in his family, blood samples of several volunteer family members were obtained and the target of interest was analyzed in their DNA samples.

For detection of *DPY19L2* deletion, a multiplex PCR assay was performed for exons 1, 5, 6, 7, 11 and 22 of this gene, together with a part of *β-ACTIN* or *PROTAMIN 1* genes, as internal control using specific primers (Table 2). Lack of amplification for all or some *DPY19L2* exons indicates respectively total or partial deletion of this gene in the studied cases. To confirm complete deletion of this gene, specific breakpoint "a" amplification was performed in the samples with lack of amplification for all *DPY19L2* exons.

Table 2: List of primers used for polymerase chain reaction and real time PCR analysis

	Genes	Amplified sequences	Primer sequence (5'→3')	Annealing temperature (°C)	Product length (bp)	
Conventional PCR	<i>β-ACTIN</i>	-	F: CGTGACATTAAGGAGAAGCTGTGC R: CTCAGGAGGAGCAATGATCTTGAT	55	375	
	<i>DPY19L2</i>	Exon 1	F: GGCCAACTTCTTTCTACTCGGAC R: GACCCAGCTCCACCATACTCCTT	65	504	
		Exon 4	F: CAAAATAGCGAGAAGTGATTAG R: TTCTACTCAACTATAAGGATACAC	54	414	
		Exon 5	F: AGCTTCATCCATGTCCTAT R: AGCCTTCTCAGAAAACATTTTT	60	432	
		Exon 6	F: GGGTAAATAATTAACACAGCA R: AAACAACAGAATAAAAGGGAT	57	462	
		Exon 7	F: AATTTATACGTACTACTTTTGTAGAATTA R: ATTTAAACATTTCAATCAACATGC	55	420	
		Exon 8	F: TGGACATGGTAGTTAATTGCTG R: TCCCAAAGTGCTGAATTGAA	55	371	
		Exon 11	F: AACCTCCTCAAGTGACTTAG R: TTGGCCAAGAGTCATT	53	516	
		Exon 22	F: GTGTCTGTTATTAAGCTTGTG R: ATTGTCTCTAGACAGCAATACAT	59	313	
		Break point "a"	-	F: ATGCCATGTTGCCTGCT R: TCTTCTGGGAAAGGTATTATCGTAG	62	1700
		<i>SPATA16</i>	Exon 4	F: AATTCTTTGCCATGTGCATATC R: GGTCAAGCGCATTTCTATTAC	58	635
	<i>PICK1</i>	Exon 13	F: TGGGCTGCCATCCATGATC R: GCTCCCAGGCTCCGTCCTC	66	568	
	<i>PROTAMINE1</i>	-	F: CCCCTGGCATCTATAACAGGCCGC R: TCAAGAACAAGGAGAGAAGAGTGG	60	530	
	Real-time PCR	<i>β-ACTIN</i>	-	F: AGATGCGTTGTTACAGGAAG R: TGTGTGGACTTGGGAGAG	60	92
<i>DPY19L2</i>		-	F: GACCCAGCTCCACCATACTCCTT R: TTCCATCTCCTCTACCTCCG	60	144	

Quantitative assessment of mutated *DPY19L2* alleles

qPCR was implemented by two alternative methods, to analyze the genotyping of *DPY19L2* gene for the family members of three cases (G8, 14 and 21) in terms of homo/hemizyosity deletion or normal state of *DPY19L2* gene. Specific *DPY19L2* and *β-ACTIN* primer pairs were designed to quantify both the target and reference genes (Table 2). Of note, primer efficiencies for target gene (*DPY19L2*) and reference gene (*β-ACTIN*) were almost equal (Fig.1).

In the first method, samples were quantified absolutely, using a control blood sample obtained from a healthy fertile donor, who voluntarily par-

ticipated in this study. After genomic DNA extraction by standard salting out procedure, 60 ng of standard genomic DNA was used as a template for further serial dilution preparations. Different amounts of DNA (60, 12, 2.5, 0.5, 0.1 ng) from this fertile donor were used as template in each PCR reaction in three set of PCR to draw a standard threshold cycle (Ct) curve (red squares shown in the Fig.1). Then, 60 ng of sample tests were subjected to PCR reactions (blue squares shown in the Fig.1). The quantity of the target gene (*DPY19L2*, lower curve shown in the Fig.1) and the reference gene (*β-ACTIN*, upper curve shown in the Fig.1) of each subject was calculated based on their Ct in the standard curve which was drawn

with different amounts of DNA from the fertile (control) sample in ABI step one plus real-time PCR system (Life Technologies, CA, USA). Proportion of PCR products of *DPY19L2* to β -*ACTIN* quantities was considered for further analyses. This proportion for fertile was considered between 0.8-1, for carrier and patient cases was approximately 0.5 (ranged 0.3-0.7) and 0 respectively, as reported earlier (18). Additionally, to assess the accuracy of this method, equal volume of DNA extracts from blood samples taken from the fertile individual (control sample) and a patient with globozoospermia (G14) were mixed and the resulting mixture was used as a heterozygous (hetero) sample.

In the second method, conventional relative quantification (RQ, using $2^{-\Delta\Delta C_t}$ equation) method was used with the same samples, utilizing 60 ng of DNA templates to quantify PCR-products of *DPY19L2* relative to β -*ACTIN*. In this study, RQ level was considered 0.8-1 for normal cases, while this level was approximately 0.5 (ranged

0.3-0.7) and 0 in carrier and patients respectively, as previously reported (18). All PCR reactions contained 5 μ l SYBR Green (TaKaRa, Japan), 0.2 μ l Rox and 5 μ M of each specific primer (0.2 μ l) for *DPY19L2* or for β -*ACTIN* (0.5 μ l) in a 10 μ l final volume of PCR reaction.

Results

Clinical characteristics of the patients with globozoospermia

Clinical parameters of the patients who participated in this study are depicted in the Table 3. Analyses showed lower sperm motility of the patients, compared to the highlighted standard criteria by World Health Organization (WHO). Regarding the round-headed shape of the sperms, in this study, ICSI technique was used to obtain successful fertilization culminated in three healthy births (Table 3). In this survey, three pedigree members that suffered from globozoospermia were further studied.




Fig.1: A standard threshold cycle (Ct) curve was drawn to calculate the allele quantities of *DPY19L2* and β -*ACTIN* for individuals who are suspected to be carrier for pathogenic allele of *DPY19L2*. As described in materials and methods, a standard Ct curve was drawn using different amounts of DNA from a fertile donor (60, 12, 2.5, 0.5 and 0.1 ng, red squares) through qPCR. Then, the quantity of the target gene (*DPY19L2*, lower curve) and the reference gene (β -*ACTIN*, upper curve) of each tested sample, for individuals who were suspected carriers of the pathogenic allele of *DPY19L2*, was calculated based on their Ct on the standard curve. The primer efficiency for both genes was almost similar. Meanwhile, regression coefficients (R²) and the slope of Ct curves were mostly equal (approximately 0.99, and -3.8 respectively).

Table 3: Clinical parameters of patients with globozoospermia

Patient	Consanguinity of the parents	Sperm parameters				ICSI attempts and results		
		Round-headed sperm (%)	Volume (mL)	Sperm concentration (10 ⁶ /mL)	Progressive motility (%)	Number of ICSI (ET cycles)	Clinical pregnancy (Abortion)	Live delivery (Sexuality)
G1	Non-familial	100	3	80	10	2	No (-) No (-)	-
G2	NA	100	3	80	30	1	No (-)	-
G3	NA	100	4	64	15	ND	-	-
G4	Familial	100	3	20	5	ND	-	-
G5	Familial	100	3.5	40	2	ND	-	-
G6	Familial	100	4	66	10	ND	-	-
G7	Non-familial	100	1	65	25	ND	-	-
G8	Non-familial	100	4	66	10	1	Yes (-)	Ongoing
G9	Familial	100	4	50	20	1	No (-)	-
G10	Familial	100	2.5	40	20	3	No (-) Yes(+) Yes (-)	1 Singleton (Girl)
G12	Non-familial	100	1.5	70	20	1	Yes (-)	1 Singleton (Boy)
G13	Non-familial	100	0.5	67	25	2	No (-) Yes (-)	1 Singleton (Girl)
G14	Familial	100	1	2	25	ND	-	-
G15	Non-familial	100	2	74	0	ND	-	-
G16	Non-familial	100	3	30	15	1 (1)	No (-) No (-)	-
G18	NA	100	4	80	10	1 (1)	No (-) No (-)	-
G19	NA	98	4	80	0	1	Yes (-)	Ongoing
G20	Familial	100	3	40	15	1 (1)	No (-) No (-)	-
G21	Non-familial	100	3.1	10	5	ND	-	-
G22	Non-familial	100	6.7	60	35	2	No (-) No (-)	-
G23*	Non-familial	100	2.1	10	5	ND	-	-
G24	Familial	100	2	18	5	1	No (-)	-
G25	Familial	96	2.9	90	40	1	No (-)	-
G26	Familial	100	2.3	40	10	ND	-	-
G27	Familial	100	1	40	15	1 (1)	No (-) No (-)	-
G28	Familial	98	2.3	28	10	ND	-	-
G29	Familial	100	3	45	40	1	No (-)	-

Two samples of G11 and G17 were lost, thus they were deleted. ET; Freeze-thawed embryo transfer, ICSI; Intra - cytoplasmic sperm insemination, NA; Not assigned, ND; Not done, +; Stands for successful pregnancy, -; Stands for abortion, and *; Globo 23 is single and not married.

Mutational analysis in *SPATA16* and *PICK1* genes

In this study, 27 cases with globozoospermia and 30 fertile men as control group were analyzed for detection of nucleotide variation (Table 1). In our first screening, regarding that digestion of the PCR products resulted in similar pattern to the fertile cases (data not shown), we did identify missense mutations of neither G848A in exon 4 of *SPATA16* gene (Fig.2, left panel) nor G198A in exon 13 of *PICK1* gene in the studied cases (Fig.2, right panel).




Fig.2: Assessment of missense mutations of G848A in the exon 4 of *SPATA16* (left panel) and G198A in the exon 13 of *PICK1* (right panel) genes using *NciI* and *PvuII* restriction endonuclease enzymes, respectively. No mutation was observed due to complete digestion of amplified fragments as described in materials and methods. M; 50 bp DNA ladder.

Analysis of *DPY19L2* deletion in the cases with globozoospermia

We have previously reported that *DPY19L2* gene deletion leads to globozoospermia (16). In this study, further to 14 (out of 18) individuals who had shown some deletion in *DPY19L2* gene, six (out of

nine) new cases with globozoospermia, missed the entire length of *DPY19L2* gene (G21, 22, 23, 24, 26 and 27, Table 1). One of these six individuals was unrelated (G24), while the remaining individuals were originated from two different pedigrees (Table 1).

This experiment was carried out by several multiplex PCR on exon 1 (Fig.3A), exon 11 (Fig.3B) and exon 22 of *DPY19L2* gene (Fig.3C) with a part of β -*ACTIN* or *PROTAMINE1* gene. Two new individuals (G25 and 28), with partially globozoospermia demonstrations, showed a wild-type condition for *DPY19L2* gene (Table 1). Furthermore, data indicated the presence of breakpoint “a” (BP_a) in most of the new cases (five out of six) with entire *DPY19L2* gene deletion (Fig.3D, respective lanes for G21 and 26).

Moreover, *DPY19L2* gene hemizyosity (complete deletion of one *DPY19L2* gene allele) was evaluated in parents of one case (G8), who has previously been recognized to suffer from deletion of entire length of *DPY19L2* gene (16). Here we confirmed hemizyosity of *DPY19L2* for both parents of G8, by amplification of BP_a (Fig.4A) and exon 11 (Fig.4B). Of note that other siblings of this family were fertile.




Fig.3: Analysis of *DPY19L2* gene deletion in exons 1, 11 and 22 as well as identification of breakpoint “a” (BP_a) in a number of individuals with globozoospermia (G#). Multiplex PCR products of A. Exon 1(504 bp, upper band), B. Multiplex PCR products of exon 22 of *DPY19L2* gene (313 bp, lower band) and part of *PROTAMINE1* gene (530 bp, upper band). Co+ or positive control in A, B and C is a fertile specimen and Co- or negative control is no template sample, C. Exon 11 (516 bp, upper band) of *DPY19L2* gene together with a part of β -*ACTIN* gene (375 bp, lower band) and D. PCR analysis of BP_a. P1 and P2 are parents of globozoospermia patient (G21) and B1 is his fertile brother and negative control is a fertile specimen, F, and positive control is a case with globozoospermia, which has been confirmed to have BP_a. M; 50 bp DNA ladder in panel A and 100 bp DNA ladder for the rest of the panels and PCR; Polymerase chain reaction.




Fig.4: Detection of exon 11 and BPA in parents of one case with globozoospermia (G8). **A.** Amplification of BPA implied hemizyosity state for *DPY19L2* gene in the parents. DNA sample of fertile individual (F) was not amplified for BPA as expected, Co+; Globozoospermia who previously proved to have BPA, P1, P2; Parents of G8, Co-; No template sample and **B.** Multiplex PCR products for exon 11 of *DPY19L2* gene (516 bp, upper band) together with a part of β -ACTIN gene (375 bp, lower band), Co+; Sample from a fertile man, Co-; No DNA template, M; 100 bp DNA ladder and P1, P2; Parents of G8.

Evaluation of familial globozoospermia

In this experiment study, two cases (G5 and 9) were selected for sibling analysis. As shown in the Figure 5, the genetic pedigree belongs to family of G5 (with the history of reproductive failure and miscarriage) revealed that all of five members (G5, 6, 20, 26, and 27) had globozoospermia associated with complete deletion of *DPY19L2* gene. We have previously demonstrated (16) a partial

deletion of *DPY19L2* including exons 5, 6 and 7 in one case (G9, Table 1). Due to infertility history of his family (Fig.6) and access to DNA samples of all family members, multiplex PCR of the aforementioned exons was performed. Curiously, we determined similar mutations pattern of *DPY19L2* gene to G9 patient, in the cousin with complete globozoospermia (G29). Indeed, detection of exons 4 and 8 by PCR confirmed this partial deletion (data not shown).




Fig.5: Pedigree of one case with globozoospermia (G5) and repeated pregnancy loss. There is more consanguineous marriage in this family but for simplicity detailed data are not depicted in the pedigree. Polymerase chain reaction (PCR) analysis showed G26, 27, 5, 6 and 20 suffering from globozoospermia due to complete deletion of *DPY19L2* gene. II2, III11 and III15 are infertile individuals with performing no genetic analysis. The inset numbers which are shown in the squares/circles represent the numbers of healthy (fertile) siblings who were not shown in this pedigree.




Fig.6: Consanguineous pedigree of the G9 family, with partial deletion of *DPY19L2* gene. Only one cousin (G29) who was also suffering from globozoospermia had deletion of exons 5, 6 and 7. The inset numbers which are shown in the squares/circles represent the numbers of healthy (fertile) siblings who were not shown in this pedigree. n; The sexuality and numbers of siblings were not determined.

DPY19L2 genotyping analysis

To set a reliable method for homo/hemizyosity state of *DPY19L2* deletion, we performed qPCR based genotyping analysis for family members of one case who showed whole *DPY19L2* gene deletion, G21. The *DPY19L2* deletion consanguineous pedigree for G21 patient is shown in the Figure 7. Data analyses demonstrated whole *DPY19L2* gene deletion of one allele for the G21 parents. Quantity proportion values of *DPY19L2* to β -*ACTIN* for carriers of *DPY19L2* deletion

were almost between 0.4- 0.6, while in normal cases, it ranged between 0.8-1 based on two calculated methods (quantities ratio and $2^{-\Delta\Delta Ct}$ ratio presented in the Table 3). In addition, no amplification plot was detected for G21, 22 and 23 indicating lack of the mentioned gene in these subjects. In addition, detection of BPa in hemizygot individuals of the family confirmed the outcomes of qPCR assay (Fig.3D, where P1 and P2 are parents of G21 patient, B1 is the hemizygot fertile brother. G21 is proband person) (Table 4).




Fig.7: Consanguineous pedigree of G21 with complete deletion of *DPY19L2* gene in BPa. *DPY19L2* genotyping analysis in this pedigree indicated that B2 and wives of G21 and B2 are hemizygot. Three pedigree members (B1 wife, G21 sister and B2 daughter) did not participate in the analysis and their zygosity state remained undefined (NA).

Table 4: *DPY19L2* genotype achieved by quantitative real-time PCR for the family members of Globo 21 and Globo 22

Subjects	Quantities ratio	$2^{-\Delta\Delta Ct}$ ratio
Father (P1)	0.522924	0.4
Mother (P2)	0.408959	0.4
Brother-1 (B1)*	1.006982	1
B1 sibling	0.957183	0.9
Brother-2 (B2)*	0.499332	0.4
B2 wife	0.414017	0.4
Globo22 wife	1.037303	1
Globo21 wife	0.488268	0.4

*; B1 and B2 are 2 fertile patient's (Globo21) brothers, Ct; Threshold cycle, and PCR; Polymerase chain reaction.

Table 5: DPY19L2 genotype achieved by quantitative real-time PCR

Subject	Quantities ratio	2 ^{-ΔΔCt} ratio
Globo 14 father	0.58377	0.5
Globo 14 mother	0.40623	0.4
Globo 8 father (P1)	0.47476	0.5
Globo 8 mother (P2)	0.424836	0.4
Hetero	0.560374	0.4
Control-1	0.973175	0.9
Control-2	1.381863	1.3

Ct; Threshold cycle and PCR; Polymerase chain reaction.

To extend the application of previously suggested method (quantities ratio) for identification of gene homo/hemizyosity at different individuals, we performed further analyses on the G8 and 14 patients' parents, besides of the hemizygot sample (hetero) as notified in materials and methods. Data affirmed the hemizygot status of the parents and hetero case by two alternatively implicated calculation methods (Table 5).

Discussion

In the recent years, there have been an increasing amounts of literatures proposing the molecular mechanisms of globozoospermia (7, 9-13, 15, 16, 19-21). Our previous studies have described DPY19L2 gene as a basic factor required for development of normal acrosome biogenesis. Partial or complete deletion of the DPY19L2 gene is pivotal factor in globozoospermia (16).

Therefore, we investigated complete deletion of DPY19L2 gene effects to reaffirm the potential association of DPY19L2 gene and globozoospermia. In addition, deletion of this gene was evaluated in the family members of three globozoospermic individuals. Thus, deletion analysis of DPY19L2 gene (12q14.2) was carried out in three exons 1, 11, 22 of DPY19L2 gene, using multiplex PCR, compared to β -ACTIN or PROTAMINE1 genes as internal controls. Briefly, all of three assessed exons of DPY19L2 gene (1, 11, 22) were missed in 20 out of 27 cases (74%) suggesting total absence of DPY19L2 gene in these cases.

It should be noted that identification of total deletion of DPY19L2 gene with BPa in 18 cases, out of 27, has previously been reported by Elinati et

al. (16). Overall, six out of nine new individuals showed complete deletion of DPY19L2 gene, five of whom carried BPa and the remaining may have unknown BP. Also, one new patient (G29) harbored a partial deletion of this gene and two others (G25 and G28) with partial globozoospermia had two wild type alleles. Previous studies have also demonstrated molecular mutations in DPY19L2 gene (19, 21). Deletion of the DPY19L2 gene is a common genomic rearrangement that occurs due to LCRs flanking the gene by NAHR. Concurrent with the cases, family members of three globozoospermic patients were investigated in this study. In this regard, two pedigrees (G5 and 9 pedigrees) from different geographically accommodation regions, similar ethnicity and high rate of consanguineous marriages showed the history of reproductive failure due to globozoospermia. Regarding high incidence of this rare abnormality among tribal races, diagnosis of carrier individuals could help them, in terms of genetic management, for future family planning.

Several studies have previously detected heterozygosity of the other genes, like SMN1 and DYSTROPHIN, through quantitative real-time PCR based on comparative Ct method (18, 22, 23). In this article, we identified the carriers in one pedigree (G21 pedigree) by this method and also proposed a modified method, quantities ratio. Thus, we designed qPCR assay for family members of G21. Analyses were performed based on proportion of DPY19L2 to β -ACTIN quantities. After providing the standard curve based on serial diluted DNA samples of a fertile man, quantities of the reference and target gene were estimated. Quantitative analysis of DPY19L2 gene for G21

family members led us to identify individuals with hemizyosity at this gene. We determined that parents with a quantity ratio ranging between 0.4-0.6 are carrier. One of the fertile brothers (B2) as well as partners of G21 and B2, were hemizygote for deletion of *DPY19L2* gene. Quantity ratio for normal cases, consisting one of the fertile brother, the grand daughter and partner of G22, were ranging from 0.95-1.3. These results were similar to previously reported threshold cycle method verifying our conclusion to determine the individuals with no gene deletion or carriers (18, 22). Considering non-consanguinity of parents, the incidence of the abnormality in this family could be attributed to their accommodation in the same geographical area. To validate our calculation method on the allele hemizyosity, we extended experiments on the G8 and G14 patients' parents who kindly accepted to participate voluntarily in this survey.

These findings are in agreement with previous studies, indicating a strong relationship between *DPY19L2* gene and globozoospermia. However, molecular cause of few cases remains yet unclear, requiring further investigations to identify genetic defect(s) in the other gene(s) affecting globozoospermia.

Regarding the other genes, in the present study mutation screening of the *SPATA16* and *PICK1* genes were also carried out on 27 cases with globozoospermia and 30 fertile men. Our data revealed that *PICK1* and *SPATA16* genes were intact in all of studied individuals.

Conclusion

Our result revealed that qPCR analysis can be used for genotyping of *DPY19L2* deletion and this may help genetic counselors in family planning. In future, it might also help prevent occurrence of this syndrome in carrier families through pre-implantation genetic diagnosis, especially in ethnic community with high consanguineous marriages.

Acknowledgements

This study was supported by Royan Institute and we would like to express our gratitude to staff of IFIC for their fully supports. There is no conflict of interest in this study.

References

1. Dam AH, Feenstra I, Westphal JR, Ramos L, van Golde RJ, Kremer JA. Globozoospermia revisited. *Hum Reprod Update*. 2007; 13(1): 63-75.
2. Xu X, Toselli PA, Russell LD, Seldin DC. Globozoospermia in mice lacking the casein kinase II alpha catalytic subunit. *Nat Genet*. 1999; 23(1): 118-121.
3. Kang-Decker N, Mantchev GT, Juneja SC, McNiven MA, Van Deursen JM. Lack of acrosome formation in Hrb-deficient mice. *Science*. 2001; 294(5546): 1531-1533.
4. Yao R, Ito C, Natsume Y, Sugitani Y, Yamanaka H, Kuretake S, et al. Lack of acrosome formation in mice lacking a Golgi protein, GOPC. *Proc Natl Acad Sci USA*. 2002; 99(17): 11211-11216.
5. Xiao N, Kam C, Shen C, Jin W, Wang J, Lee KM, et al. PICK1 deficiency causes male infertility in mice by disrupting acrosome formation. *J Clin Invest*. 2009; 119(4): 802-812.
6. Yildiz Y, Matern H, Thompson B, Allegood JC, Warren RL, Ramirez DM, et al. Mutation of beta-glucosidase 2 causes glycolipid storage disease and impaired male fertility. *J Clin Invest*. 2006; 116(11): 2985-2994.
7. Paiardi C, Pasini ME, Gloria M, Berruti G. Failure of acrosome formation and globozoospermia in the wobbler mouse, a *Vps54* spontaneous recessive mutant. *Spermatogenesis*. 2011; 1(1): 52-62.
8. Lin YN, Roy A, Yan W, Burns KH, Matzuk MM. Loss of zona pellucida binding proteins in the acrosomal matrix disrupts acrosome biogenesis and sperm morphogenesis. *Mol Cell Biol*. 2007; 27(19): 6794-6805.
9. Audouard C, Christians E. Hsp90beta1 knockout targeted to male germline: a mouse model for globozoospermia. *Fertil Steril*. 2011; 95(4): 1475-1477.
10. Liu G, Shi QW, Lu GX. A newly discovered mutation in PICK1 in a human with globozoospermia. *Asian J Androl*. 2010; 12(4): 556-560.
11. Dam AH, Kosciński I, Kremer JA, Moutou C, Jaeger AS, Oudakker AR, et al. Homozygous mutation in *SPATA16* is associated with male infertility in human globozoospermia. *Am J Hum Genet*. 2007; 81(4): 813-820.
12. Harbuz R, Zouari R, Pierre V, Ben Khelifa M, Kharouf M, Coutton C, et al. A Recurrent deletion of *DPY19L2* causes infertility in man by blocking sperm head elongation and acrosome formation. *Am J Hum Genet*. 2011; 88(3): 351-361.
13. Kosciński I, Elinati E, Fossard C, Redin C, Muller J, Velez de la Calle J, et al. *DPY19L2* deletion as a major cause of globozoospermia. *Am J Hum Genet*. 2011; 88(3): 344-350.
14. Xu M, Xiao J, Chen J, Li J, Yin L, Zhu H, et al. Identification and characterization of a novel human testis-specific Golgi protein, NYD-SP12. *Mol Hum Reprod*. 2003; 9(1): 9-17.
15. Pierre V, Martinez G, Coutton C, Delaroche J, Yassine S, Novella C, et al. Absence of *Dpy19l2*, a new inner nuclear membrane protein, causes globozoospermia in mice by preventing the anchoring of the acrosome to the nucleus. *Development*. 2012; 139(16): 2955-2965.
16. Elinati E, Kuentz P, Redin C, Jaber S, Vanden Meerschaut F, Makarian J, et al. Globozoospermia is mainly due to *DPY19L2* deletion via non-allelic homologous recombination involving two recombination hotspots. *Hum Mol Genet*. 2012; 21(16): 3695-3702.
17. Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res*. 1988; 16(3): 1215.
18. Lee TM, Kim SW, Lee KS, Jin HS, Koo SK, Jo I, et al.

- Quantitative analysis of SMN1 gene and estimation of SMN1 deletion carrier frequency in Korean population based on real-time PCR. *J Korean Med Sci.* 2004; 19(6): 870-873.
19. Coutton C, Zouari R, Abada F, Ben Khelifa M, Merdassi G, Triki C, et al. MLPA and sequence analysis of DPY19L2 reveals point mutations causing globozoospermia. *Hum Reprod.* 2012; 27(8): 2549-2558.
 20. Kuentz P, Vanden Meerschaut F, Elinati E, Nasr-Esfahani MH, Gurgan T, Iqbal N, et al. Assisted oocyte activation overcomes fertilization failure in globozoospermic patients regardless of the DPY19L2 status. *Hum Reprod.* 2013; 28(4): 1054-1061.
 21. Zhu F, Gong F, Lin G, Lu G. DPY19L2 gene mutations are a major cause of globozoospermia: identification of three novel point mutations. *Mol Hum Reprod.* 2013; 19(6): 395-404.
 22. Akbari MT, Noruzinia M, Mozdarani H, Hamid M. Determination of exon 7 SMN1 deletion in Iranian patients and heterozygous carriers by quantitative real-time PCR. *J Genet.* 2011; 90(1): 133-136.
 23. Maksimovic N, Andjelkovic A, Rasic VM, Stojanovic VR, Kotlica BK, Brankovic S, et al. Quantitative analysis of the dystrophin gene by real-time PCR. *Arch Biol Sci.* 2012; 64(2): 787-792.
-

Archive of SID