

The Frequency of *Staphylococcus aureus* Isolated from Endocervix of Infertile Women in Northwest Iran

Mohammad Taghi Akhi, Ph.D.^{1,2}, Aylin Esmailkhani, M.Sc.^{2*}, Javid Sadeghi, Ph.D.², Behrooz Niknafs, Ph.D.³, Laya Farzadi, M.D.⁴, Aydin Akhi, Pharm.D.⁵, Elmira Najafi Nasab, M.Sc.²

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran
2. Department of Bacteriology and Virology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran
3. Department of Anatomical Sciences, Medical Faculty, Tabriz University of Medical Sciences, Tabriz, Iran
4. Women's Reproductive Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran
5. Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran

Abstract

Background: Infertility is one of the major social issues. Due to the asymptomatic cervical infection associated with *Staphylococcus aureus* (*S. aureus*), the majority of patients remain undiagnosed. The present study intended to assess the frequency of *S. aureus* isolated from infertile women's endocervix in northwest Iran.

Materials and Methods: In a descriptive cross sectional study, specimens were randomly collected during vagina examination using a sterile speculum and swabbing. After performance of antibiotic susceptibility testing, polymerase chain reaction (PCR) was used to identify methicillin-resistance *S. aureus* (MRSA) and toxic shock syndrome toxin-1 (TSST-1).

Results: About 26 (26%) and 9 (9%) women's urogenital tracts were colonized by *S. aureus* and *Candida* spp., respectively, of which three (11.5%) patients were infected with fungi and *S. aureus*, simultaneously. Antibiotic susceptibility results showed high activity of vancomycin and co-trimoxazole on isolates. Regarding PCR results, *mecA* sequences were detected in 7 (26.9%) strains, whilst the *tst* gene encoding TSST-1 was not detected in any of clinical strains.

Conclusion: The prevalence of *S. aureus* was very high in infertile women. Therefore, it demands all patients undergoing infertility treatment to be investigated thoroughly for this type of infection.

Keywords: Infertility, *S.taphylococcus aureus*, Endocervix, *mecA*

Citation: Akhi MT, Esmailkhani A, Sadeghi J, Niknafs B, Farzadi L, Akhi A, Najafi Nasab E. The frequency of staphylococcus aureus isolated from endocervix of infertile women in northwest Iran. Int J Fertil Steril. 2017; 11(1): 28-32.

Introduction

One of the most common reproductive health issues in developing countries is the high rate of infertility (1). It becomes a globally important subject for clinical research and practice because infertility affects 60 to 168 million individuals, both women and men, in reproductive age. It has been reported that of 10 couples, one couple incurs the early or secondary stages infertility (2, 3). Infertility is characterized as inability

to conceive during one year, despite normal cohabitation (4), as indicated by the European Society of Human Reproduction and Embryology (ESHRE), which is evidenced in 10-15% of all couples (5).

There are a number of factors which are responsible for infertility in females. According to a study by Vayena et al. (3), the rates of primary infertility are generally between 1 and 8% with rates of secondary infertility reaching as high as

Received: 3 Jul 2016, Accepted: 24 Aug 2016

*Corresponding Address: P.O.Box: 5165665931, Department of Bacteriology and Virology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran
Email: a.esmailkhani@gmail.com



Royan Institute
International Journal of Fertility and Sterility
Vol 11, No 1, Apr-Jun 2017, Pages: 28-32

35%. Infection of reproductive organs is one of the most important factors affecting infertility, while the determination of the type of infection has a significant contribution to the treatment of this issue. However, the significance of these infections in the genital tracts is not well known. Many microorganisms, including bacteria, viruses, parasites and fungi, seem to be able to interfere with the reproductive function in both genders. Bacterial vaginosis is a prevalent issue among women with changing the balance of normal vaginal flora such as lactobacilli (6, 7). However, some related pathogenic impacts have been evidenced such as increased rates of premature rupture of the membranes (PROM), late miscarriage in first trimester, preterm labor, endometriosis and delivery (6). Bacterial vaginosis is the most common lower genital tract disorder among reproductive age of women (pregnant and non-pregnant) and the most common cause of vaginal odor and malodorous discharge (8).

The bacteria encountered in the female genital tract can be divided into aerobic and anaerobic organisms. Among the aerobic Gram-positive organisms, there are several species of *Streptococcus* (*S.*), such as group B *S.* (GBS), and among Gram-positive facultative anaerobic organisms, there are several species of *Enterococcus* (*E.*) (9). *S. aureus* is an infrequent but one of the most successful human pathogens. It has the ability to cause a number of infections in various environmental corners within the host. *S. aureus* has additionally been reported to be commonly isolated microorganism from cervical specimens (10). It is found in the genital tract of approximately 9 to 10% of asymptomatic women, approximately 10% of patients with postoperative wound abscess after gynecologic or obstetric procedures, and 5 to 20% of genital tract cultures of women with pelvic infection. It is also detected in virtually 100% of women who have toxic shock syndrome toxin-1 (TSST-1) (9). This study was, therefore, designed to assess the frequency of *S. aureus* isolated from infertile women's endocervix in northwest Iran.

Materials and Methods

Sampling

Descriptive cross sectional study was carried

out at the Tabriz University of Medical Science (TUMS), Tabriz, Iran, between April and July 2015. The cervical samples were randomly taken from 100 women who attended the Department of Obstetrics and Gynecology of Milad Infertility Center affiliated to TUMS for unexplained infertility during the mentioned-period. Patients with infertility due to unrelated reasons to *S. aureus* infections, such as ovarian cancer, lazy ovary, ovarian cysts and polycystic ovary syndrome (PCOS), were excluded from the study.

The Ethic Commission of TUMS approved this study (Number: 5/412912-2015). After obtaining a written consent form from all patients, the specimens were collected using a sterile vaginal speculum and swab.

Isolation and identification

Samples were streaked on blood agar and manitol salt agar (Merck, Germany) plates and incubated aerobically at 37°C for 24-48 hours. After overnight incubation, the isolates were examined by Gram staining technique using catalase, DNase and coagulase tests (11).

Antibiotic susceptibility testing

The susceptibility of *S. aureus* isolates to antimicrobial agents was measured *in vitro* using the disc diffusion method according to Clinical and Laboratory Standards Institute (CLSI) protocols (12). The tested antibiotics included penicillin, gentamicin, ciprofloxacin, vancomycin, trimethoprim/sulfamethoxazole and ceftiofloxacin (MAST Diagnostics, UK).

Detection of *mecA* and *tst* genes

Bacterial DNA was extracted from the isolates according to tissue buffer boiling method (13). Firstly, 20 µl tissue buffer [0.25% sodium dodecyl sulfate (SDS)+0.05M NaOH (CinnaClon, Iran)] were mixed with one colony of bacterial isolate, the combination was incubated for ten minutes in 95°C, the mixture was centrifuged for one minute in 13000 g, 180 µl Milli-Q water (CinnaClon, Iran) were slowly added, and finally extracted DNA was frozen in -20°C for durable storage.

DNA isolates with the concentration of 0.1 ng/µl

were used as the templates for polymerase chain reaction (PCR) analysis. Multiplex PCR was carried out by CinnaGen MsterMix (CinnaClon, Iran) using the *mecA* and *tst* primers, as described previously (14).

The sequences of the *mecA* primers used were: 5'-ACTGCTATCCACCCTCAAAC-3' and 5'-CTGGTGAAGTTGTAATCTGG-3', while the sequences of the *tst* primers used were: 5'-ACCCCTGTTCCCTTATCATC-3' and 5'-TTTTTCAGTATTTGTAACGCC-3' (synthesized at the CinnaClon, Iran).

The strains 92-S-1344 (*tst*) and 95-S-739 (*mecA*) were used as positive control in this study. Amplification was carried out using a thermocycler (Eppendorf, Germany) as follows: i. Initial denaturation at 94°C for 5 minutes, ii. 35 cycles of denaturation at 94°C for 2 minutes, annealing at 57°C for 2 minutes, and primer extension at 72°C for 1 minutes, and iii. Terminal extension at 72°C for 7 minutes. Electrophoresis of PCR products was performed on 1% agarose gel (CinnaClon, Iran). The gel staining was performed in ethidium bromide for 20 minutes and visualized using a gel documentation system (UVP, USA).

Results

During the study period, a total of 100 infertile women were included in this study. All participants underwent intrauterine insemination (IUI). About 26 (26%) and 9 (9%) women's urogenital tracts were colonized by *S. aureus* and *Candida* spp., respectively, which were identified by mycology methods. Among them, three (8.5%) patients were infected with fungus and *S. aureus*, simultaneously. The average age of these patients was 30.96 years, ranging between 18 and 56 years. Of the patients who were colonized by *S. aureus*, 11 (42.1%) were under 30 years and 14 (53.7%) were over 30 years of age. After examining the wives of the patients who were colonized by *S. aureus*, the semen samples of four (15.38%) couples were positive for *S. aureus*. In addition, penicillin-resistant strains of *S. aureus* were colonized in the reproductive system of a couple. Of those *Candida* spp. carriers, three (3 of 9) wives were also infected with *Candida* spp. The results of antibiotic susceptibility testing are summarized in Table 1.

Table 1: The results of antibiotic susceptibility testing

Antibiotic	Sensitive (%)	Intermediate (%)	Resistant (%)
Penicillin	3 (11.53)	0 (0)	23 (88.46)
Gentamicin	19 (73.07)	0 (0)	7 (26.09)
Ciprofloxacin	19 (73.07)	0 (0)	7 (26.09)
Vancomycin	25 (96.15)	1 (3.84)	0 (0)
Co-trimoxazole	25 (96.15)	0 (0)	1 (3.84)
Cefoxitin	19 (73.07)	0 (0)	7 (26.09)

The data were presented as N (%).

In general, vancomycin and co-trimoxazole showed high activity against the isolates. Regarding PCR results, *mecA* sequences were detected in 7 (26.9%) isolates, whilst the *tst* gene encoding TSST-1 was not detected in any of clinical isolates.




Fig.1: Multiplex PCR product for the *mecA* and *tst* genes. M; 100 bp molecular weight marker, Lane 1; *tst* gene positive control (ATCC 92-S-1344), Lane 2; *mecA* gene positive control (ATCC 95-S-739), Lane 3; Negative control, and Lane 4-9; *mecA* positive .

Discussion

The study investigated the prevalence of *S. aureus* in infertile women. An in-depth study on these genera has not yet been conducted in Iran. Generally, infectious vaginitis is a prevalent disorder with noteworthy clinical results if left untreated. Infertility is an important health issue with far-reaching consequences on couple, family planning program,

health system and spread of sexually transmitted diseases (STD) and acquired immunodeficiency syndrome (AIDS). It can be characterized as the lack of a conception after at least one year of constant, unprotected sexual intercourse (15).

In a study by Okonofua et al. (10), *Candida albicans* (25%), *S. aureus* (21.7%) and *Neisseria gonorrhoeae* (17.4%) were the most commonly isolated microorganisms; however, there was no difference between fertile and infertile women in the rates of isolation of these pathogens. In most cases, resistance to penicillin is attributable to β -lactamase production. We found that the most antibiotic resistance was against penicillin, which is not supported by a study conducted by Ghiasi et al. (16), in which they have indicated 100% of *S. aureus* were sensitive. Our findings also indicated that none of the *S. aureus* strains were resistant to vancomycin. In a study by El-Ghodban et al. (17), they have reported that less than 50% of *S. aureus* strains were β -lactamase producers and resistant to penicillin. However, almost 75% of the strains originating from food were positive for β -lactamase and resistant to penicillin. In our study, all isolates were positive for *mecA* genes, while they were resistant to the gentamicin, ciprofloxacin and cefoxitin. de A Trindade et al. (18) have also reported that among methicillin-resistant *S. aureus* (MRSA) isolated from blood samples, twenty (13%) individuals were susceptible to four or more antimicrobials.

The incidence of TSST-1-producing strains has been registered worldwide (19). Colonization with *S. aureus* is generally highest (20 to 30%) in the oropharynx or nose of non-healthcare workers. Vaginal colonization with *S. aureus* has been determined to be lower (10 to 20%) in the United States, Europe, and Asia (20). Similarly, TSST-1-producing strains of *S. aureus* have been isolated vaginally from 1 to 4% of healthy, menstruating women in the general population (19). Due to a higher immune response to TSST-1, *S.* infections are more common in developing countries than developed countries (21). We decided to use tst in the present study because of limited reports from many developing countries. In a study by Parsonnet et al. (19) carried out on Japanese women, of the 159 *S. aureus* isolates recovered, 14 (9%) were TSST-1 positive, suggesting that only 47% of women had positive titers of anti-

TSST-1 antibody, which is significantly lower than the reported seropositivity rates in the Europe and United States (20). In the same study by El-Ghodban et al. (17), TSST-1 was detected in only three (7.5%) of 40 *S. aureus* clinical strains and in none of the food strains. In another study by Tsen et al. (22), they have firmly reported the comparative discoveries, in which only three (4.8%) of 62 strains of *S. aureus* from clinical sources as tst-carrying strains were identified using PCR, but none of their food strains carried this gene.

In the etiologies of infertility, the most contributed factors are related to female (40 to 55%) followed by male factors (30 to 40%), both partners (10%) and unexplained factors (10%) (1). There are several factors which increase risks for acquisition of bacterial vaginosis, while bacterial vaginosis is more prevalent in women who smoke (23), black women (24), women who are sexually active compared with virginal women (25), and women who utilize vaginal douches (26).

The infertility leads to decreased levels of personal well-being, while for many individuals, it causes more serious consequences (27). Subsequently, it appears that screening is a reasonable approach that is likely to be cost effective. However, all physicians must have a high index of suspicion and utilize promptly accessible screening methods to detect and treat the patients with infectious vaginitis adequately (28). A limitation of the present study was the lack of evaluation of *S. aureus* in fertile women. However, it is suggested that further studies will be conducted on a larger sample.

Conclusion

The exact knowledge of *S. aureus* colonization rate in infertile women has a great importance. Therefore, it demands all patients undergoing infertility treatment to be investigated thoroughly. Such screening and treatment during the course of infertility treatment increase the pregnancy rate extensively. However, randomized studies with larger number of participants are needed to reach more validated conclusions.

Acknowledgements

This research was financially supported by a grant from Immunology Research Center of Tabriz

University of Medical Science. We would like to thank the staff of Milad Infertility Center and the Microbiology Department for their help. The authors announce that there is no conflict of interests in this article.

References

1. Devi A, Ranjani A, Dhanasekaran D, Thajuddin N, Ramanidevi T. Surveillance of multidrug resistant bacteria pathogens from female infertility cases. *Afr J Biotechnol*. 2013; 12(26): 4129-4134.
2. Evens EM. A global perspective on infertility: an under recognized public health issue. The University of North Carolina at Chapel Hill. 2004; 18: 1-42.
3. Vayena E, Rowe PJ, Peterson HB. Assisted reproductive technology in developing countries: why should we care? *Fertil Steril*. 2002; 78(1): 13-15.
4. Zabrek EM. "Can I get pregnant?": the basic infertility workup. *Clin Obstet Gynecol*. 1996; 39(1): 223-230.
5. Forti G, Krausz C. Clinical review 100: evaluation and treatment of the infertile couple. *J Clin Endocrinol Metab*. 1998; 83(12): 4177-4188.
6. Hay PE, Lamont RF, Taylor-Robinson D, Morgan DJ, Ison C, Pearson J. Abnormal bacterial colonisation of the genital tract and subsequent preterm delivery and late miscarriage. *BMJ*. 1994; 308(6924): 295-298.
7. Yudin MH, Money DM. Infectious Diseases Committee. Screening and management of bacterial vaginosis in pregnancy. *J Obstet Gynaecol Can*. 2008; 30(8): 702-716.
8. Fleury FJ. Adult vaginitis. *Clin Obstet Gynecol*. 1981; 24(2): 407-438.
9. Gibbs RS. Microbiology of the female genital tract. *Am J Obstet Gynecol*. 1987; 156(2): 491-495.
10. Okonofua FE, Ako-Nai KA, Dighitoghi MD. Lower genital tract infections in infertile Nigerian women compared with controls. *Genitourin Med*. 1995; 71(3): 163-168.
11. Duguid JP. *Staphylococcus: cluster-forming gram-positive cocci*. In: Colle JG, Duguid JP, Fraser AG, Marmion BP, editors. *Mackie and McCartney practical medical microbiology*. 13th ed. New York: Churchill Livingstone; 1989; 303-316.
12. Patel JB, Cockerill FR, Bradford PA, Eliopoulos GM, Hindler JA, Jenkins SG, et al. Performance standards for antimicrobial susceptibility testing: twenty-fifth informational supplement. *Clinical and Laboratory Standards Institute (CLSI)*. 2015; 35(3): 29-50.
13. Asgharzadeh M, Yousef S, Samadi Kafil H, Nahaei MR, Ansarin K, Akhi MT. Comparing transmission of *Mycobacterium tuberculosis* in East Azarbaijan and West Azarbaijan provinces of Iran by using IS6110-RFLP method. *Biotechnology*. 2007; 6(2): 273-277.
14. Mehrotra M, Wang G, Johnson WM. Multiplex PCR for detection of genes for *Staphylococcus aureus* enterotoxins, exfoliative toxins, toxic shock syndrome toxin 1, and methicillin resistance. *J Clin Microbiol*. 2000; 38(3): 1032-1035.
15. Vander H, Prabha V. Evaluation of fertility outcome as a consequence of intravaginal inoculation with sperm-impairing micro-organisms in a mouse model. *J Med Microbiol*. 2015; 64(Pt 4): 344-347.
16. Ghiasi M, Fazaeli H, Kalhor N, Sheykh-Hasan M, Tabatabaei-Qomi R. Assessing the prevalence of bacterial vaginosis among infertile women of Qom city. *Iran J Microbiol*. 2014; 6(6): 404-408.
17. El-Ghodban A, Ghenghesh KS, Márialigeti K, Esahli H, Tawil A. PCR detection of toxic shock syndrome toxin of *Staphylococcus aureus* from Tripoli, Libya. *J Med Microbiol*. 2006; 55(Pt 2): 179-182.
18. de A Trindade P, Pacheco RL, Costa SF, Rossi F, Barone AA, Mamizuka EM, et al. Prevalence of SCCmec type IV in nosocomial bloodstream isolates of methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol*. 2005; 43(7): 3435-3437.
19. Parsonnet J, Goering RV, Hansmann MA, Jones MB, Ohtagaki K, Davis CC, et al. Prevalence of toxic shock syndrome toxin 1 (TSST-1)-producing strains of *Staphylococcus aureus* and antibody to TSST-1 among healthy Japanese women. *J Clin Microbiol*. 2008; 46(8): 2731-2738.
20. Parsonnet J, Hansmann MA, Delaney ML, Modern PA, DuBois AM, Wieland-Alter W, et al. Prevalence of toxic shock syndrome toxin 1-producing *Staphylococcus aureus* and the presence of antibodies to this superantigen in menstruating women. *J Clin Microbiol*. 2005; 43(9): 4628-4634.
21. Vergeront JM, Stolz SJ, Crass BA, Nelson DB, Davis JP, Bergdoll MS. Prevalence of serum antibody to staphylococcal enterotoxin F among Wisconsin residents: implications for toxic-shock syndrome. *J Infect Dis*. 1983; 148(4): 692-698.
22. Tsen HY, Yu GK, Wang KC, Lin LY. Comparison of the enterotoxigenic types, toxic shock syndrome toxin I (TSST-1) strains and antibiotic susceptibilities for enterotoxigenic *Staphylococcus aureus* strains isolated from food and clinical samples. *Food Microbiol*. 1998; 15(1): 33-41.
23. Jonsson M, Karlsson R, Rylander E, Gustavsson A, Wadell G. The associations between risk behaviour and reported history of sexually transmitted diseases, among young women: a population-based study. *Int J STD AIDS*. 1997; 8(8): 501-505.
24. Goldenberg RL, Klebanoff MA, Nugent R, Krohn MA, Hillier S, Andrews WW. Bacterial colonization of the vagina during pregnancy in four ethnic groups. *Vaginal infections and prematurity study group*. *Am J Obstet Gynecol*. 1996; 174(5): 1618-1621.
25. Bump RC, Buesching WJ 3rd. Bacterial vaginosis in virginal and sexually active adolescent females: evidence against exclusive sexual transmission. *Am J Obstet Gynecol*. 1988; 158(4): 935-939.
26. Hawes SE, Hillier SL, Benedetti J, Stevens CE, Koutsky LA, Wolner-Hanssen P, et al. Hydrogen peroxide-producing lactobacilli and acquisition of vaginal infections. *J Infect Dis*. 1996; 174(5): 1058-1063.
27. Fidler AT, Bernstein J. Infertility: from a personal to a public health problem. *Public Health Rep*. 1999; 114(6): 494-511.
28. Rein MF, Liang BA. Diagnosis and treatment of infectious vaginitis. *Hospital Physician*. 1999; 35(10): 46-58.