

The Effects of *Calligonum* Extract and Low-Intensity Ultrasound on Motility, Viability, and DNA Fragmentation of Human Frozen-Thawed Semen Samples

Hamid Ekrami, M.Sc.^{1,2}, Mansoureh Movahedin, Ph.D.^{1*}, Fereshteh Koosha, Ph.D.³, Zohreh Mazaheri, Ph.D.², Manijhe Mokhtari-Dizaji, Ph.D.⁴

1. Department of Anatomical Sciences, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Basic Medical Science Research Center, Histogenotech Company, Tehran, Iran

3. Department of Radiology Technology, Faculty of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Medical Physics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Abstract

Background: The study aimed to evaluate the impact of *Calligonum* extract and US radiation on sperm parameters of cryopreserved human semen samples.

Materials and Methods: In this experimental study, twenty-five semen specimens were obtained from healthy semen donors and incubated in human tubal fluid (HTF) medium supplemented with 10% human serum albumin (HSA) for 45 minutes. Samples were treated with *Calligonum* extract (10 µg/ml) alone (CGM group) and US radiation (LIPUS-exposed group) alone or a combination of both treatments (CGM+LIPUS). The US group received US stimulation (in both continuous and pulsed wave modes) at a frequency of 1 MHz and intensity of 200 mW/cm² for 200 seconds. Sperm morphology was assessed by Diff-Quik staining. The DNA fragmentation was evaluated the Halo sperm kit. Sperm parameters was analyzed by a computer-assisted semen analysis system. Reactive oxygen species (ROS) was assessed by flow cytometry.

Results: The results showed that the treatment with *Calligonum* extract significantly ($P < 0.05$) increased the progressive motility of spermatozoa in the CGM group as compared with the control group. The application of low-intensity US significantly ($P < 0.05$) decreased the motility and viability of spermatozoa in the US group when compared with the control group. Our findings also indicated that the use of both low-intensity US in continuous mode and *Calligonum* extract slightly increased progressive motility; however, such an increase was not statistically significant. The rate of DNA fragmentation was considerably higher ($P < 0.05$) in control and LIPUS-exposed groups than the other groups.

Conclusion: Treatment of spermatozoa with *Calligonum* extract slightly improved the sperm parameters due to its antioxidant activity, on the other hand, according to our results, US radiation did not improve sperm parameters which may be due to interference with the motility of sperm, as well as its physical effects on spermatozoa.

Keywords: Antioxidants, *Calligonum*, Cryopreservation, Low-Intensity Ultrasound, Spermatozoa

Citation: Ekrami H, Movahedin M, Koosha F, Mazaheri Z, Mokhtari-Dizaji M. The effects of calligonum extract and low-intensity ultrasound on motility, viability, and DNA fragmentation of human frozen-thawed semen samples. *Int J Fertil Steril.* 2020; 14(2): 84-90. doi: 10.22074/ijfs.2020.5896.

This open-access article has been published under the terms of the Creative Commons Attribution Non-Commercial 3.0 (CC BY-NC 3.0).

Introduction

Cryopreservation of human semen is considered one of the most vital and essential strategies for the preservation and maintenance of spermatozoa, and it is broadly applied in malignancies or other therapies which could damage to the functionality of testicles (1). During the cryopreservation process, the viability, motility and the integrity of chromatin might be adversely influenced, usually accompanied by the elevation of oxidative stress and excessive production of reactive oxygen species (ROS) (2). In the presence of increased ROS production, free radical oxygen molecules can target and attack to bis-allylic methylene groups of phospholipid-bound polyunsaturated fatty

acids which are present in the plasma membrane of spermatozoa, leading to lipid peroxidation (3). The impact of lipid peroxidation on the sperm function include irreversible loss of motility, discharge of intracellular enzymes, sperm DNA damage, impairment of oocyte penetration and sperm-oocyte fusion, and apoptosis of spermatozoa in frozen media (4).

Many efforts have been made to minimize the rate of cryodamage in frozen spermatozoa during the cryopreservation process. For instance, the addition of antioxidant agents or cryoprotectants to the extender is one of the promising tools for the increase of sperm quality during the freeze-thaw process (5-7). Various types

Received: 3/March/2019, Accepted: 28/Jan/2020

*Corresponding Address: P.O.Box: 1441633665, Department of Anatomical Sciences, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Email: movahed.m@modares.ac.ir



Royan Institute
International Journal of Fertility and Sterility
Vol 14, No 2, July-September 2020, Pages: 84-90

www.SID.ir

of antioxidants, such as vitamin C or E (8), genistein (5), and resveratrol (9) have been employed for cryopreservation. However, the yield of the cryopreservation procedure for human spermatozoa is not yet satisfactory, and new strategies are still warranted to improve the fidelity of cryopreservation. *Calligonum comosum* (*C. comosum*) is a medicinal plant abundantly found in the Egyptian desert, containing numerous polyphenol antioxidant agents. The beneficial roles of *C. comosum* may stem from its antioxidant activity, which has been extensively assessed *in vitro* (10). Some studies have demonstrated some biological properties of *C. comosum* extract on rodent models, such as the anti-inflammatory and anti-gastric ulcer activities (11).

Additionally, *C. comosum* showed the anticancer potential in mice inoculated with Ehrlich ascites carcinoma cells (12). It has been implicated that low-intensity pulsed ultrasound (LIPUS) at the incident power density from 0.5 to 3000 mW/cm² have been used for bone healing (13). LIPUS can promote the growth of human skin fibroblasts, thereby the activation of the integrin receptors, RhoA (Ras homolog gene family, member A)/ROCK, and Src-ERK signaling pathways (14, 15). The biological action of LIPUS occurs when the mechanical wave is converted into a biochemical signal within the cell as the mechanoreceptors and integrin are thought to be involved in this process. A number of studies have highlighted the formation of focal adhesions on the surface of the cells treated with LIPUS, which is mediated by the activation of integrin-associated signaling pathways (16, 17). Focal adhesions are large multi-protein complexes that could serve as a transmembrane bridge between the extracellular matrix (ECM) and the actin cytoskeleton. They could be identified in specific sites within the cell where clustered integrin receptors can interact with the ECM components on the outside and with the actin cytoskeleton on the inside of the cells. One of the main components of focal adhesion proteins involved in the transduction of the LIPUS signal from a mechanical force to a chemical messenger is focal adhesion kinase (FAK), which is phosphorylated when the cells are exposed to LIPUS (17, 18).

Hence, regarding the above statements, it would be plausible that LIPUS can enhance the penetration of herb extraction into the sperm via an increase in motility of spermatozoa. Therefore, the primary goal of the present study was to determine the impact of *C. comosum* extract alone, LIPUS signal, and the combination of both on the count, viability, total motility, progressive motility, DNA fragmentation, and morphology of spermatozoa during the freeze-thawing process.

Materials and Methods

Study design

In this experimental study, we evaluated the effects of *Calligonum* extract and LIPUS at a frequency of 1 MHz (in pulsed and continues wave modes) on the cryo-

preservation of human spermatozoa. After the preparation of semen samples, each sample was liquated into 5 parts included; washed spermatozoa (control group), frozen-thawed spermatozoa, frozen-thawed spermatozoa treated with *Calligonum* extract at a concentration of 10 µg/ml (CGM group), frozen-thawed spermatozoa exposed to the US radiation (LIPUS-exposed group) at a duty cycle of 40% (pulsed mode, at a frequency of 1 MHz, at the incident power density of 200 mW/cm² for 200 seconds), and frozen-thawed spermatozoa treated with the combination of *Calligonum* extract and the US radiation with continues mode (CGM+LIPUS group). The present research was approved by Ethical Committee of Tarbiat Modares University (No. 52/6757 dated 30.11.92).

Herb extraction

The identification of the herb

The plant (*Calligonum comosum* L.) was collected by Dr. Hosein Batooli from the desert located at the proximity of Kashan, Iran, (33.9850°N, 51.4100°E). The taxonomic identity of the collected plant was confirmed by Dr. Abdoalrasool Haghiri Ebrahim Abadi, Faculty of Science, Kashan University, Iran. Freshly collected plant materials were air-dried in the shade at room temperature. The aerial parts (stem, flowers, and leaves) of this herb were used for further investigations.

The extraction protocol

Two kilograms of the fresh aerial parts of the plant (equal to 50% of the weight of a wildy-growing plant) were air-dried in the shade at room temperature, grounded, and exhaustively extracted by cold maceration with aqueous methanol (70%). The extract was evaporated under reduced pressure at 40°C to yield 80 g residue. The residue was suspended in distilled water and successfully fractionated with n-hexane, CH₂Cl₂, EtOAc, and n-BuOH (Thermo Fisher, USA) saturated with H₂O. Each extract was evaporated under reduced pressure to yield 3, 7, 12 and 22 g residues, respectively.

The antioxidant activity

The activity of free radical-scavenging of the methanolic extract of *C. comosum* was determined concerning the potential to neutralize the free radical-producing [2,2-diphenyl-1-picrylhydrazyl (DPPH)] according to a method, as described previously (19, 20). Concisely, the level of DPPH was calculated by the measurement of the absorbance at the wavelength of 517 nm prior to and after the addition of a specific amount of the herb extract. Afterward, the inhibitory percentage of radical formation was estimated using the following equation: % inhibition = $(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100$, where A_{control} is considered the absorbance of DPPH alone and A_{sample} is regarded as the absorbance of DPPH in combination with the *C. comosum* extract.

Semen sample collection

In this experimental study, semen samples were obtained from twenty-five fertile men with the average age of 34 (range, 25-50 years) who referred to the *in vitro* fertilization (IVF) center of Gandhi Hospital according to the World Health Organization (WHO) criteria. The process of semen collection was carried out by masturbation into a sterile, wide-mouthed, calibrated glass container after five days of sexual abstinence. The collected specimens were allowed to liquefy at 37°C for 30 minutes prior to the semen analysis. The entire samples were collected from individuals who referred for medical procedures during assisted reproductive technology (ART). We used the remaining samples obtained from patients, with their permission, for research purposes.

Sperm preparation by the swim-up method

A fraction of motile spermatozoa was selected to be analyzed by the swim-up method. For this purpose, 1 ml of each semen sample (kept at 37°C and 5% CO₂) was added to 3 ml of human tubal fluid (HTF, Genocell Co., Iran) supplemented with 10% human albumin serum (HAS, Biotest, Germany) and centrifuged at 2000 ×g for 3 minutes. Afterward, 0.5 ml of HTF supplemented with 10% HAS was gently added on the resultant pellets. The samples were then incubated at 37°C, 5% CO₂, and inclined at a 45° angle to incubator for 45 minutes. Consequently, 0.5 ml of the uppermost medium was recovered, and the swim-up method was performed (21).

Sperm freezing and thawing

Each processed semen sample was cryopreserved according to the standard protocol for sperm freezing. According to Li et al. (7) and Ibrahim et al. (22), semen samples were gently mixed with an equal volume of modified cryoprotective medium (Global Media, USA) supplemented with 10 µg/ml of *Calligonum* extract. The samples were kept at 4°C for 30 minutes and then frozen by placing the straws horizontally at 10 cm above the surface of liquid nitrogen (nitrogen vapor) at -80°C for 20 minutes. Finally, all frozen straws were stored in liquid nitrogen until use.

At the thawing stage, the cryo-straws were removed from liquid nitrogen and immediately immersed in a water bath at 37°C for at least 1-2 minutes. The thawed straws containing semen samples were flicked and inverted to mix the contents before sampling thoroughly and then washed with a culture medium HTF supplemented with 10% HAS (The samples were then treated with *Calligonum* extract (10 µg/ml)). At the end step, semen samples were centrifuged at 2000 ×g for 3 minutes to remove any trace of cryoprotectant in the freezing medium. The samples were analyzed for motility, viability, morphology, and DNA fragmentation (9, 23).

Preparation of ultrasound device

The US device (Physiomed, Germany) was set up at the frequency of 1 MHz, incident power density of 200

mW/cm², 200-second time period, and a 45-minute period as the duration of the experiment. These fine-tuned parameters were chosen based on our previous study published in this regard (24). The US stimulation was carried out in which the transducer was put at the focal distance (3.5 cm) of a cell culture plate incubated in 5% CO₂ at 32°C. It was transmitted through the bottom of the well via coupling gel between the transducer and the tube. The device was transmitted through the bottom of the well via coupling the gel between the transducer and cell culture plate. Notably, no temperature change more than 1°C was recorded during the US stimulation.

Sperm stimulation using low-intensity pulsed ultrasound

Sperms maintained in the HTF medium supplemented with 10% HAS (Gibco, Germany). The samples were exposed to low-intensity pulsed ultrasound (LIPUS) (1 MHz, 200 mW/cm² and 40% DC) alone or in combination with *C. comosum* coined as experimental groups. The control group was also cultured in the HTF medium supplemented with 10% HAS. After the US stimulation, spermatozoa were incubated for 45 minutes in 5% CO₂ at 32°C, similar to other experimental groups. To investigate the sperm parameters, the mean number of whole cells per volume, viability, morphology, and motility were examined after the incubation process.

Semen samples analysis

The sperm count, motility, morphology, viability, and DNA fragmentation were evaluated according to the guidelines introduced by the WHO (25). The sperm motility analysis was conducted using light microscopy (at ×400 magnification), combined with a computer-assisted semen analysis system (CASA, Video Test Sperm 1.2, Russia). A Makler chamber was used for the scoring of the sperm motility at room temperature. A minimum of 100 spermatozoa from at least five different fields was analyzed from each aliquot. Sperm morphology was assessed by the Diff-Quik staining method according to WHO guidelines by light microscopy (Labomed, USA). The sperm viability was determined by Eosin B (5% in saline) (Merck, Germany) staining technique at one-step. In this procedure, viable or dead spermatozoa are recognized by white (unstained) and pink (or red) coloration in the head region of the sperm cells, respectively.

Sperm DNA fragmentation

The Halotech DNA G2 kit (Spain) was used to study the DNA fragmentation in frozen-thawed spermatozoa. Based on the sperm chromatin dispersion test (SCD), the intact frozen-thawed spermatozoa were diluted in a culture medium HTF supplemented with 10% HAS to achieve sperm concentration of 15-20 million per milliliter. In this method, 50 µl of the semen sample was added to 100µl of dissolved agarose (0.7%) (Sigma, US). Afterward, 25 µl of the cell suspension was transferred to slides, pre-coated with

Archive of SID

0.65% agarose and covered with a coverslip without any air bubbles, and then incubated at 4°C for 5 minutes. After the removal of coverslips, the slides were horizontally immersed in a tray with freshly prepared acid denaturation solution (0.08 N HCl) for 7 minutes at 22°C in a dark place to create restricted single-stranded DNA from DNA breaks. The slides were then immersed in lysis Solution I [0.4 M Tris (Sigma, USA), 0.4 M 2-Mercaptophenol (Sigma, USA), 1% sodium dodecyl sulfate (SDS, Sigma, USA), and 50 mM Ethylenediaminetetraacetic acid (EDTA, Sigma, USA), pH=7.5 for 20 minutes and lysis Solution II (0.4 M Tris, 2 M NaCl, and 1% SDS, pH=7.5) for 5 minutes to remove nuclear proteins. Slides were then rinsed with distilled water for 5 minutes, followed by dehydration through an ascending gradient of ethanol (70, 90, and 100%) for 2 minutes. The slides were then placed at room temperature to be air-dried.

For Diff-Quik staining, the slides were incubated in eosin solution for 6 min; then, transferred into Azur B solution, for another 6 minutes. The nucleoid of spermatozoa with fragmented DNA did not develop a dispersion halo, or the halo was minimal. From each slide, a minimum of 500 spermatozoa was scored under an oil-immersion objective ($\times 100$ magnification) by light microscopy (Labomed, USA). The sperm cells showing no halo, small halo, medium halo, large halo, or fragmentation were separately scored. Spermatozoa indicating no halo/fragmentation were considered to have damaged chromatin, and the results were expressed as a percentage of sperm cells with DNA damage.

Intracellular reactive oxygen species level measurement

Spermatozoa were rinsed with phosphate-buffered saline (PBS, Atocel, Austria) and incubated with 20 μ M 2, 7-dichlorofluorescein diacetate (DCFH-DA, Life Technologies, USA), diluted in serum-free medium at 37°C for 45 minutes. The intracellular ROS level was immediately analyzed with flow cytometry at an excitation wavelength of 488 nm and an emission wavelength of 530 nm (26).

Statistical analysis

The analysis of the values obtained in this study was performed by the SPSS version 19 (SPSS Inc., IBM company, USA), while the level of significance was set at $P < 0.001$. The difference between the values of each group was analyzed by one-way analysis of variance (one-way ANOVA), followed by Tukey's test. The data were expressed as the means \pm standard deviation (means \pm SD).

Results

The activity of radical scavenging of DPPH

The *C. comosum* extract showed free radical-scavenging activity in a dose-dependent manner when examined by the DPPH assay. The inhibitory concentration (IC_{50}) value (the concentration of a given substance causing 50% inhibition of the DPPH) of the herb extract was

29.2 μ g/ml when compared with ascorbic acid (positive control).

Effect of *Calligonum* extract and/or LIPUS on sperm parameters

Motility assessment

The demographic characteristics of patients are demonstrated in Table 1.

Table 1: The demography of semen samples

Characteristics	Average	SD
Age (Y)	34	3.041
Total motility (%)	80.7795	1.69
Grade A (%)	33.1496	1.74
Grade B (%)	30.7443	1.56
Grade C (%)	16.8856	1.39
Grade D (%)	19.4599	1.58
Grade A+B (%)	64.0651	2.08
Viability (%)	87	1.27
Normal morphology (%)	28.6397	4.26
SDF (%)	10.63	2.15
ROS (%)	11.0804	2.16

Grade A; Linear progressive motility, Grade B; Progressive motility, Grade C; Non-progressive motility, Grade D; Immotile, SDF; Sperm DNA fragmentation, ROS; Reactive oxygen species, and SD; Standard Deviation.

All of the sperm parameters, including viability, motility, and morphology in all experimental groups, are shown in Table 2. Accordingly, the percentage of total motility of spermatozoa in the fresh group was 89.37 ± 3.74 , while this value was 81.30 ± 5.72 in the frozen-thawed group. After the addition of 10 μ g/ml of *Calligonum* extract in the freezing medium increased the total motility; yet, such an increase was not statistically significant when compared with the control group ($P \geq 0.05$). LIPUS (in pulsed and continuous mode waves) alone decreased the motility of spermatozoa, compared with the control group ($P \leq 0.038$), but the combination of *Calligonum* extract and LIPUS increased the motility to (81.90 ± 3.93) and (81.65 ± 5.18) respectively when used in pulsed mode and continuous modes waves. Also, there was an improvement in progressive motility of spermatozoa treated with the combination of *Calligonum* extract and LIPUS, compared with the frozen-thawed group; however, the difference between two groups was not statistically significant ($P \geq 0.05$).

Viability assessment

As shown in Table 2, the viability of spermatozoa in all groups undergone the freeze-thawing process was significantly reduced as compared with the fresh group ($P \leq 0.026$). Such a reduction was more pronounced in the LIPUS-exposed and CGM-treated groups, compared with other groups. There was no significant difference between the CGM+LIPUS and control groups (frozen-thawed spermatozoa) (frozen-thawed spermatozoa, $P \geq 0.05$).

Table 2: Comparison of sperm parameters (\pm SD) between the experimental groups after frozen-thawed and treatment with 10 μ g/ml Calligonum (CGM) extract and LIPUS (pulsed mode and continues wave)

Groups	Control	Frozen - thawed	Frozen thawed+CGM extract	Frozenthawed+ continues wave of ultrasound	Frozenthawed+ continues wave of trasound+CGM extract	Frozenthawed+ pulsed mode of ultrasound	Frozenthawed+ pulsed mode of ultrasound +CGM extract
Viability (%)	94.17 ^c \pm 5.97	87.52 \pm 4.75	90.11 \pm 6.12	82.76 ^b \pm 5.68	86.70 \pm 7.12	81.82 ^b \pm 2.87	87.94 \pm 4.25
Total motility (%)	89.37 \pm 3.74	81.30 \pm 3.74	85.42 \pm 4.28	76.76 ^{ab} \pm 5.29	81.65 \pm 5.18	75.86 ^{ab} \pm 3.47	81.90 \pm 3.93
Progressive motility (%)	38.80 \pm 2.57	34.81 ^a \pm 3.28	37.81 \pm 4.13	33.04 ^{ab} \pm 2.98	37.24 \pm 5.11	32.60 ^{ab} \pm 3.58	36.18 ^a \pm 4.78
Normal morphology (%)	30.11 \pm 3.16	26.58 \pm 3.54	28.41 \pm 4.27	27.58 \pm 3.95	30.82 \pm 2.75	28.35 \pm 4.82	30.05 \pm 3.66

^a; Significant difference with control group in the same row ($P \leq 0.05$), ^b; Significant difference with freeze and thawed group + CGM extract, in the same row $P \leq 0.026$, and ^c; Significant difference with other groups in the same row ($P \leq 0.047$).

Table 3: Assessment of fragmented DNA and free radicals percentage in all experimental groups

Groups	Control	Frozen - thawed	Frozen - thawed + CGM extract	Frozen - thawed + continues wave of ultrasound	Frozen - thawed + continues wave of ultrasound+CGM extract	Frozen - thawed + pulsed mode of ultrasound	Frozen - thawed + pulsed mode of ultrasound+CGM extract
Fragmented DNA (%) \pm SD	24.5 \pm 7.23	46.5 ^a \pm 9.26	37 \pm 4.29	38 \pm 5.37	34 \pm 5.17	51 ^a \pm 11.85	42.5 ^a \pm 7.39
Free radicals(ROS) (%) \pm SD	-	12.46 \pm 7.06	9.18 \pm 2.57	12.52 \pm 8.14	7.77 \pm 2.06	14.75 \pm 4.67	10.05 \pm 1.68

CGM; Calligonum extract, ROS; Reactive oxygen species, and ^a; significant difference with control group in the same row ($P \leq 0.042$).

Morphology assessment

According to Table 2, the normal morphology score was 30.11 ± 3.16 in the fresh group, while the normal morphology score was 26.58 ± 3.54 in the control group. The statistical analysis revealed that there was no significant difference in the score of normal morphology among all treated groups, namely, the CGM-treated, LIPUS-exposed, and CGM+LIPUS groups.

Effect of Calligonum extract and/or LIPUS (pulsed or continuous mode waves) on DNA fragmentation

Table 3 shows the rate of DNA fragmentation in all experimental groups. The percentage of spermatozoa undergone DNA fragmentation was considerably higher in all groups than the fresh group. The rate of DNA fragmentation was significantly ($P \leq 0.042$) elevated in the frozen-thawed (46.5 ± 9.26) and LIPUS-exposed groups (51 ± 11.85) in comparison with the control group.

Effect of Calligonum extract and/or LIPUS on reactive oxygen species level

According to Table 3, the level of ROS in the control group receiving no treatment was 12.46 ± 7.06 , while the level of ROS was decreased to 9.18 ± 2.57 in the CGM group. The statistical analysis demonstrated that the difference in the amount of ROS was not significant in all experimental groups as compared with the control group ($P \geq 0.05$).

Discussion

Cryopreservation of human semen provides a valuable therapeutic opportunity for the management of patients who are at risk of infertility (27). However, during cryopreservation, spermatozoa experience physical and chemical stress that could result in detrimental changes in lipid composition of the cell membrane, leading to the excessive amount of ROS production, as well as a decrease in sperm motility and viability (28, 29). Notably, the osmotic stress and the alterations in the temperature may cause mechanical stress to the cell membrane of spermatozoa. Therefore, the changes mentioned above could reduce the fertilization capability of human spermatozoa after the cryopreservation process (30).

It has been implicated that antioxidant therapy can increase the quality of cryopreserved spermatozoa when employed both *in vitro* and *in vivo* (31). Numerous antioxidant agents have been indicated to have beneficial roles in the protection against cellular damages caused by cryopreservation-induced ROS, affecting sperm motility and viability. Hence, the application of antioxidant compounds to neutralize the deleterious impacts of oxidative damage would be of asset to improve the sperm parameters (32). Therefore, in this study, we examined the effects of *Calligonum* extraction, as an antioxidant agent, to annul the harmful effects of free radical molecules generated during the freeze-thawing process. We also applied LIPUS (in continuous and pulsed mode waves) to induce

physical stimulation to spermatozoa for possible inciting the motility and viability of sperm cells after cryopreservation.

Our data revealed *Calligonum* extract at a concentration of 10 µg/ml slightly increased the sperm motility and viability, whereas it decreased DNA fragmentation and ROS level in human spermatozoa. The combination therapy using LIPUS and *Calligonum* extract could enhance sperm parameters when compared with the LIPUS-exposed group. Our findings were consistent with the previous studies performed in this area. For instance, Martínez-Soto et al. (29) evaluated the effects of extender supplemented with genistein on frozen-thawed human spermatozoa. They found that genistein, known as isoflavone compound, had antioxidant properties on cryopreserved spermatozoa. Their results also showed that the ROS production was decreased and the sperm motility was slightly improved in response to treatment with genistein. In another study, Banihani et al. (33) found that L-carnitine had positive effects on the improvement of sperm motility and viability during cryopreservation but had no effect on the reduction of DNA oxidation. Regarding our results, the addition of *Calligonum* extract to the freezing medium led to a significant increase in progressive motility.

On the other hand, LIPUS at the frequency of 1 MHz and incident power density of 200 mW/cm², in both pulsed and continuous mode waves, had adverse impacts on the sperm parameters. In the LIPUS-exposed group, the viability, as well as total and progressive motility was decreased while the number of non-motile spermatozoa was increased as compared with the fresh group. The US radiation alone increased intracellular ROS level and disrupted the balance of pro-oxidant and antioxidant contents in human spermatozoa, led to the elevated rate of DNA fragmentation. Also, our data demonstrated that LIPUS did not alter the morphology of frozen spermatozoa; however, the combinatory treatment of spermatozoa with *Calligonum* extract and LIPUS was capable of enhancing the sperm parameters during cryopreservation.

Previous studies have highlighted that LIPUS, as mechanical energy, could have therapeutic effects on bone and wound healing (34, 35). Also, it has been reported that it could expedite the process of tissue repair by the stimulation of the proliferation of fibroblasts and osteoblasts (17, 36). Furthermore, Xu et al. (37) have shown that LIPUS can stimulate the viability of freeze-thawed spermatozoa and it can increase the proliferation and differentiation of hematopoietic stem cells, obtained from fresh and cryopreserved peripheral blood leukapheresis product. However, inconsistent with other studies, our findings failed to show the beneficial effects of this therapeutic approach on the sperm parameters.

In the present study, the adverse effects of LIPUS on the sperm parameters may be due to the changes in the frequency of waves as a result of the sperm motility, leading to the reduction in the effectiveness of LIPUS penetration. Since the distribution of the US field is not homog-

enous, and it is susceptible to the reflection and attenuation once the field passes through the boundary separating two different media (38), the potency of US waves might be mitigated. So further examinations on applying other techniques for irradiating ultra sound on motile sperms is needed.

Conclusion

Our results showed that *Calligonum* extract, at a concentration of 10 µg/ml, slightly enhanced the cellular parameters of cryopreserved spermatozoa diminished the rate of DNA fragmentation, and decreased the intracellular ROS level. Contrary to our expectation, LIPUS, in both pulsed and continuous mode waves, had the adverse effects on the sperm parameters which may stem from alterations in the temperature of the medium as a result of LIPUS treatment. It should be noted that the sperm motility might influence the frequency of the US waves, lowering the effectiveness of LIPUS treatment. This phenomenon can weaken the impact of LIPUS on the sperm parameters during cryopreservation.

Acknowledgements

This study was financially supported by Research Deputy of Tarbiat Modares University, Tehran, Iran. There is no conflict of interest in this study.

Authors' Contributions

M.M.; Contributed to the conception and design of the work and acquisition, analysis and interpretation of the data; H.E.; Participated in study design, data collection and evaluation, drafting and statistical analysis. Z.M., K.F.; Contributed extensively in interpretation of the data and the conclusion. M.M.D.; Designed and conducted ultrasound irradiation part and read early draft and gave some corrections. All authors read and approved the final manuscript.

References

1. Meamar M, Zribi N, Cambi M, Tamburrino L, Marchiani S, Filimberti E, et al. Sperm DNA fragmentation induced by cryopreservation: new insights and effect of a natural extract from *Opuntia ficus-indica*. *Fertil Steril*. 2012; 98(2): 326-333.
2. Aitken RJ, Baker MA. Oxidative stress, sperm survival and fertility control. *Mol Cell Endocrinol*. 2006; 250(1-2): 66-69.
3. Agarwal A, Makker K, Sharma R. Clinical relevance of oxidative stress in male factor infertility: an update. *Am J Reprod Immunol*. 2008; 59(1): 2-11.
4. Gadea J, Molla M, Selles E, Marco MA, Garcia-Vazquez FA, Gardon JC. Reduced glutathione content in human sperm is decreased after cryopreservation: effect of the addition of reduced glutathione to the freezing and thawing extenders. *Cryobiology*. 2011; 62(1): 40-46.
5. Thomson LK, Fleming SD, Aitken RJ, De Iulius GN, Zieschang J-A, Clark AM. Cryopreservation-induced human sperm DNA damage is predominantly mediated by oxidative stress rather than apoptosis. *Hum Reprod*. 2009; 24(9): 2061-2070.
6. Kalthur G, Raj S, Thiyagarajan A, Kumar S, Kumar P, Adiga SK. Vitamin E supplementation in semen-freezing medium improves the motility and protects sperm from freeze-thaw-induced DNA damE-

- krami et al. *Int J Fertil Steril*, Vol 14, No 2, July-September 2020 90 age. *Fertil Steril*. 2011; 95(3): 1149-1151.
7. Li Z, Lin Q, Liu R, Xiao W, Liu W. Protective effects of ascorbate and catalase on human spermatozoa during cryopreservation. *J Androl*. 2010; 31(5): 437-444.
 8. Askari HA, Check JH, Peymer N, Bollendorf A. Effect of natural antioxidants tocopherol and ascorbic acids in maintenance of sperm activity during freeze-thaw process. *Arch Androl*. 1994; 33(1): 11-15.
 9. Garcez ME, dos Santos Branco C, Lara LV, Pasqualotto FF, Salvador M. Effects of resveratrol supplementation on cryopreservation medium of human semen. *Fertil Steril*. 2010; 94(6): 2118-2121.
 10. Abdo W, Hirata A, Shukry M, Kamal T, Abdel-Sattar E, Mahrous E, et al. Calligonum comosum extract inhibits diethylnitrosamine-induced hepatocarcinogenesis in rats. *Oncol Lett*. 2015; 10(2): 716-722.
 11. Liu XM, Zakaria MN, Islam MW, Radhakrishnan R, Ismail A, Chen HB, et al. Anti-inflammatory and anti-ulcer activity of Calligonum comosum in rats. *Fitoterapia*. 2001; 72(5): 487-491.
 12. Badria FA, Ameen M, Akl MR. Evaluation of cytotoxic compounds from Calligonum comosum L. growing in Egypt. *Z Naturforsch C J Biosci*. 2007; 62(9-10): 656-660.
 13. Khanna A, Nelmes RT, Gougoulas N, Maffulli N, Gray J. The effects of LIPUS on soft-tissue healing: a review of literature. *Br Med Bull*. 2009; 89: 169-182.
 14. Iwashina T, Mochida J, Miyazaki T, Watanabe T, Iwabuchi S, Ando K, et al. Low-intensity pulsed ultrasound stimulates cell proliferation and proteoglycan production in rabbit intervertebral disc cells cultured in alginate. *Biomaterials*. 2006; 27(3): 354-361.
 15. Kobayashi Y, Sakai D, Iwashina T, Iwabuchi S, Mochida J. Low-intensity pulsed ultrasound stimulates cell proliferation, proteoglycan synthesis and expression of growth factor-related genes in human nucleus pulposus cell line. *Eur Cell Mater*. 2009; 17: 15-22.
 16. Mahoney CM, Morgan MR, Harrison A, Humphries MJ, Bass MD. Therapeutic ultrasound bypasses canonical syndecan-4 signaling to activate rac1. *J Biol Chem*. 2009; 284(13): 8898-8909.
 17. Zhou S, Schmelz A, Seufferlein T, Li Y, Zhao J, Bachem MG. Molecular mechanisms of low intensity pulsed ultrasound in human skin fibroblasts. *J Biol Chem*. 2004; 279(52): 54463-54469.
 18. Tang CH, Yang RS, Huang TH, Lu DY, Chuang WJ, Huang TF, et al. Ultrasound stimulates cyclooxygenase-2 expression and increases bone formation through integrin, focal adhesion kinase, phosphatidylinositol 3-kinase, and Akt pathway in osteoblasts. *Mol Pharmacol*. 2006; 69(6): 2047-2057.
 19. Loganayaki N, Siddhuraju P, Manian S. Antioxidant activity and free radical scavenging capacity of phenolic extracts from *Helicteres isora* L. and *Ceiba pentandra* L. *J Food Sci Technol*. 2013; 50(4): 687-695.
 20. Kedare SB, Singh RP. Genesis and development of DPPH method of antioxidant assay. *J Food Sci Technol*. 2011; 48(4): 412-422.
 21. Saylan A, Duman S. Efficacy of hyaluronic acid in the selection of human spermatozoa with intact DNA by the swim-up method. *Cell J*. 2016; 18(1): 83-88.
 22. Ibrahim SF, Osman K, Das S, Othman AM, Majid NA, Rahman MP. A study of the antioxidant effect of alpha lipoic acids on sperm quality. *Clinics (Sao Paulo)*. 2008; 63(4): 545-550.
 23. Donnelly ET, McClure N, Lewis SE. Cryopreservation of human semen and prepared sperm: effects on motility parameters and DNA integrity. *Fertil Steril*. 2001; 76(5): 892-900.
 24. Mohaqiq M, Movahedin M, Mokhtari Dizchi M, Mazaheri Z. Investigation on the effect of low intensity ultrasound stimulation on mouse spermatogonial stem cell proliferation and colonization. *Anat Sci J*. 2013; 10(3): 119-124.
 25. Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, et al. World Health Organization reference values for human semen characteristics. *Hum Reprod Update*. 2010; 16(3): 231-245.
 26. Chang JC, Lien CF, Lee WS, Chang HR, Hsu YC, Luo YP, et al. Intermittent Hypoxia prevents myocardial mitochondrial Ca²⁺ overload and cell death during ischemia/reperfusion: the role of reactive oxygen species. *Cells*. 2019; 8(6): pii: E564.
 27. Oehninger S, Duru NK, Srisombut C, Morshedi M. Assessment of sperm cryodamage and strategies to improve outcome. *Mol Cell Endocrinol*. 2000; 169(1-2): 3-10.
 28. O'connell M, McClure N, Lewis SE. The effects of cryopreservation on sperm morphology, motility and mitochondrial function. *Hum Reprod*. 2002; 17(3): 704-709.
 29. Martínez-Soto JC, de Dioshourcade J, Gutiérrez-Adán A, Landeras JL, Gadea J. Effect of genistein supplementation of thawing medium on characteristics of frozen human spermatozoa. *Asian J Androl*. 2010; 12(3): 431-441.
 30. Câmara DR, Silva SV, Almeida FC, Nunes JF, Guerra MM. Effects of antioxidants and duration of pre-freezing equilibration on frozen-thawed ram semen. *Theriogenology*. 2011; 76(2): 342-350.
 31. Shi X, Hu H, Ji G, Zhang J, Liu R, Zhang H, et al. Protective Effect of sucrose and antioxidants on cryopreservation of sperm motility and DNA integrity in C57BL/6 mice. *Biopreserv Biobank*. 2018; 16(6): 444-450.
 32. Yousef MI, Abdallah GA, Kamel KI. Effect of ascorbic acid and vitamin E supplementation on semen quality and biochemical parameters of male rabbits. *Anim Reprod Sci*. 2003; 76(1-2): 99-111.
 33. Banihani S, Agarwal A, Sharma R, Bayachou M. Cryoprotective effect of L-carnitine on motility, vitality and DNA oxidation of human spermatozoa. *Andrologia*. 2014; 46(6): 637-641.
 34. Heckman JD, Ryaby JP, McCabe J, Frey JJ, Kilcoyne RF. Acceleration of tibial fracture-healing by non-invasive, low-intensity pulsed ultrasound. *J Bone Joint Surg Am*. 1994; 76(1): 26-34.
 35. El-Bialy T. Therapeutic ultrasound applications in craniofacial growth, healing and tissue engineering. *Rejuvenation Res*. 2007; 10(3): 367-371.
 36. Doan N, Reher P, Meghji S, Harris M. In vitro effects of therapeutic ultrasound on cell proliferation, protein synthesis, and cytokine production by human fibroblasts, osteoblasts, and monocytes. *J Oral Maxillofac Surg*. 1999; 57(4): 409-419.
 37. Xu P, Gul-Uludag H, Ang WT, Yang X, Huang M, Marquez-Curtis L, et al. Low-intensity pulsed ultrasound-mediated stimulation of hematopoietic stem/progenitor cell viability, proliferation and differentiation in vitro. *Biotechnol Lett*. 2012; 34(10): 1965-1973.
 38. Patel US, Ghorayeb SR, Yamashita Y, Atanda F, Walmsley AD, Scheven BA. Ultrasound field characterization and bioeffects in multiwell culture plates. *J Ther Ultrasound*. 2015; 3: 8.