

Response of Four Phytoplankton Species Found in Some Sectors of Nigerian Coastal Waters to Crude Oil in Controlled Ecosystem

Adekunle, I. M.*, Ajijo, M.R., Adeofun, C.O. and Omoniyi, I.T.

Department of Environmental Management and Toxicology, Department of Fisheries and Aquaculture, University of Agriculture, P.M.B. 2240, Abeokuta, Nigeria

Received 12 April 2009;

Revised 17 July 2009;

Accepted 27 July 2009

ABSTRACT: Identification and enumeration of phytoplankton species from Ilaje and Lagos sectors of the Nigerian coastal waters were conducted using standard procedures. Effects of different crude oil concentrations (6 to 50 ppm) on population of *Coscnodiscus centralis*, *Thalassionema frauenfeldii*, *Odontella mobiliensis*, and *Ceratium trichoceros* at different exposure periods (6 to 42 h) via microcosm experiments were then assessed. Results showed that the phytoplankton species consisted of diatoms (83.33%) and dinoflagellates (16.67%) whose abundance ranged from 2 to 516 Cell/mL. Crude oil toxicity varied from 0.06 to 36.43% for *C.centralis*, 1.41 to 35.58% for *C.trichoceros*, 1.71 to 46.11% for *T.frauenfeldii* and 0.66 to 44.90% for *O.mobiliensis* and showed direct relationship ($r = +0.81$ to $+0.97$; $p < 0.001$) with concentration but inverse with exposure period ($r = -0.83$ to -0.90 ; $p < 0.001$). Vulnerability within 24-h contact decreased in the order: *T.frauenfeldii* > *O.mobiliensis* > *C.centralis* > *C.trichoceros*. Study is a contribution to the scarce data bank on crude oil dose-response assessment on plankton species in Nigeria, demonstrating that influx of crude oil into the Nigerian coastal waters is a risk factor to ecological status.

Key words: Crude oil, Coastal waters, Phytoplankton, Toxicity, Environment, Nigeria

INTRODUCTION

Nigeria is one of the leading oil producing countries of the world with an economy heavily dependent on the oil sector, which provides about 20% of gross domestic product, 70 to 90% of foreign exchange earnings and about 65% of budgetary revenues. Crude oil, for several decades, has been the main export product of Nigeria and the oil reserve is put at 36 billion barrels while production is estimated at 2.1 million barrels per day (Eyong *et al.*, 2004; Egberongbe *et al.*, 2006; Adeniyi *et al.*, 2008). The oil reserves are found in relatively simple geological structures along the country's coastal Niger Delta and deeper water offshore.

The Niger delta region consists of the states of Abia, Akwa Ibom, Bayelsa, Cross River, Delta, Edo, Imo and Ondo, all in the southern region of the country. Typical oil wells, pipelines and infrastructure in the Niger Delta are shown in Fig.1.

*Corresponding author E-mail: imkunle@yahoo.com

According to available statistics, in the last 30 years more than 400,000 tons of oil has spilled into the creeks and soils of southern Nigeria (Egberongbe *et al.*, 2006). Oil spills in Nigeria occur due to a number of factors including sabotage, corrosion of pipes and storage tanks, carelessness during production operations and oil tankers' accidents. Oil obviously has brought economic boost to the country but the exploration activities, associated spills in the Niger Delta and attendant environmental issues have been of national and global concern (Powel *et al.*, 1985). Reports on the negative environmental impact ranging from vegetation destruction, water pollution, destruction of some terrestrial and aquatic flora and fauna are documented (Isichei and Sanford, 1979; Powel *et al.*, 1985; Kakulu and Osibanjo, 1986; Snowden and Ekweozor, 1987; Kakulu *et al.*, 1987; Kakulu and Osibanjo, 1988; Adekambi, 1989; Jinadu, 1989; Eyong *et al.*, 2004; Igwebuike *et al.*, 2007).




Fig. 1. Some oil wells and oil infrastructure in the Niger Delta

Studies on crude oil effects on phytoplankton species, which are not just aquatic primary producers, the base of food web structure but also a key factor to global oxygen production in the country, are limited. Previous studies on phytoplankton research in Nigeria were primarily focused on composition, abundance and temporal variation (Khan and Ejike, 1984; Opute, 1990; Kadiri, 1993; Kemdirim, 2001; Ezra and Nwankwo, 2001; Ajayi and Akonai, 2003; Akin-Oriola, 2003; Chindah, 2003; Chindah, 2004; Nyananyo et al., 2006; Essien *et al.*, 2006; Essien *et al.*, 2007; Akinyemi and Nwankwo, 2007; Ajayi *et al.*, 2007). Ekom (2006) investigated the effect of waste engine oil on phytoplankton of the Calabar river estuary while Obire and Anyawu (2009) assessed the effect of various concentrations of crude oil on fungi populations of soil. Identification of microalgal bioindicator for crude oil pollution and natural remediation as well as petroleum hydrocarbon levels in some organisms and sediments from Qua Iboea estuary, Nigeria were also investigated (Essien *et al.*, 2006; Essien *et al.*, 2008; Benson *et al.*, 2008).

Although several decades of research has produced data on oil and hydrocarbon effects on a wide range of phytoplankton species in the developed countries (Gordon and Prouse, 1973; Miller *et al.*, 1978; Kusk, 1980; Karydis and Fogg, 1980; Kelly *et al.*, 1999; Dzierzbicka-Glowacka, 2007), there is paucity of information on risk assessment with regards to a defined dose-

response relationship for Nigerian crude oil to species of phytoplankton found in the Nigerian coastal waters.

Experiments for optimization purposes are better carried out in controlled environments rather than open ecosystems such as the marines due to the fact that species like phytoplankton are passively carried along by water currents and their biomass is affected by processes such as seasonal variation, turbulent diffusion, sinking, primary production, respiration, natural mortality, grazing, cell lyses or sudden appearance of in the habitat, toxic pollutant other than the investigated xenobiotic chemical (Kelly *et al.*, 1999; Onyema, 2007; Adekunle *et al.*, 2007a; Dzierzbicka-Glowacka, 2007).

This study was therefore designed to assess the risk of low level crude oil concentrations on the population of four phytoplankton species (*Thalassionema frauenfeldii*, *Cosnoddiscus centralis*, *Odontella mobiliensis* and *Ceratium trichoceros*) found in the costal waters of Ilaje, Ondo state (oil producing area) and Bar-beach Lagos (non-oil producing area) under controlled laboratory conditions.

MATERIALS & METHODS

Ilaje is located in Ondo state, Niger Delta region of Nigeria and comprises many towns including the study locations namely Aiyetoro, Ilowo and Ilepete communities, located along the shore line (Fig. 2).




Fig. 2. Ilaje communities in Ondo state showing Ayetoro, Ilepete and Awoye

The area is characterized by a difficult terrain owing to the predominantly shoreline/marine nature, silt, mud and sedimentary deposits, freshwater swamps and marshland with few islands in-between. Bar-beach Lagos is one of the two popular beaches in Lagos, Nigeria, selected as a coastal segment found in non-oil producing area of the country.

The physico-chemical characteristics of sea water samples were determined following standard procedures (APHA, 1989; Adekunle et al., 2007b). Temperature, salinity, pH and electrical conductivity were measured using 0 – 100°C mercury in glass thermometer, and HANNA series of meters. The pH meter was calibrated with 4 and 9 buffers. Phosphate was analyzed as orthophosphate with the ammonium molybdate - ascorbic acid spectrophotometric procedure. The resulting blue colored antimony-phosphomolybdate complex was measured at 880 nm using a 1 cm ÷ 3B Perkin Elmer spectrophotometer. A standard calibration curve was prepared using potassium hydrogen phosphate salt (KH₂PO₄) in the range of 0.05 to 0.35 mg/L.

Nitrate was determined via the phenoldisulphonic method. The absorbance values

were recorded on the same spectrophotometer but at the wavelength of 410 nm and a calibration curve was prepared from KNO₃ whose working range varied from 0 to 3 mg/L. Dissolved oxygen was determined by direct probe analysis using dissolved oxygen meter. Biochemical oxygen demand (BOD) was assessed via dilution method and chemical oxygen demand (COD) by the dichromate procedure under reflux.

Phytoplankton samples were collected from Aiyetoro, Ilowo and Ilepete in Ilaje, Ondo State and Bar-beach Lagos, Lagos State, Nigeria using 53µm plankton net and 1.5 L specimen plastic bottles. With the aid of a fishing boat, the net was lowered into the seawater, 5 cm deep, and dragged for five minutes while the boat was still in motion. Only short tows of 5 minutes were taken to reduce error caused by net clogging. The tow replicates were taken at different times of the day at each study station. Each sample was concentrated to a volume of 200 mL. The plankton samples for species identification and enumeration were transferred into sampling bottles and fixed with 5% formalin. Those for toxicity test were transferred into sampling bottles that contained no formalin so as to keep alive. At the chemical

laboratory, phytoplankton species were enumerated using a 1 mL –Sedwick-Rafter counting chamber and each was counted 3 times on a phase contrast compound microscope with an eye piece. The common marine plankton identification key was utilized (Kolb, 1986; Kemdirim, 2001).

Nigerian crude oil with specific gravity (AP) of 41.7 to 43.0 was used and experiment was carried out under illumination (12:12 h light: dark period) provided by white fluorescent lamps of 20 W each at room temperature of $29 \pm 1^\circ\text{C}$. A stock solution (1000 ppm) of the crude oil was prepared by taking the equivalent volume that gave 0.1g crude oil using a micropipette, which was then transferred to a 100 mL volumetric flask, already containing 30 mL of n-hexane and made up to mark with the solvent. Seven different lower concentrations of crude oil (6, 8, 10, 14, 20 and 50 ppm) were thereafter prepared from the stock solution and the effect of each crude oil concentration on the population of the 4 phytoplankton species (*T. frauenfeldii*, *C. centralis*, *O. mobiliensis* and *C. trichoceros*) at different contact periods of 6-h interval (6, 12, 18, 30, 36 and 42 hours) was investigated as subsequently described.

Erlenmeyer 500 mL flasks were used as incubation reactors and triplicates of each treatment and control (consisting only of n-hexane treatments to the phytoplankton species) were set up. Exactly 200 mL sea water was filtered through a 0.22 μm membrane and the initial population of each of the 4 phytoplankton species recorded as previously described. A given concentration of crude oil (6, 8, 10, 14, 20 or 50 ppm) was gradually added and the flask gently swirled for uniform distribution and aerators were connected to all the flasks. The reduction in phytoplankton population (mortality) was used as a measure of response to crude oil induced stress; hence, phytoplankton species cell abundance at the end of each experiment was related to the initial population and expressed as percentage mortality:

$$\text{Mortality (\%)} = \frac{[P_i - P_f] - [P_i - P_c]}{P_i} \times 100$$

where P_i = initial cell abundance, P_f = cell abundance at the end of the crude oil treatment and P_c = cell abundance at the end of the control

experiment. Data generated, generally expressed as mean \pm standard error, were subjected to descriptive statistics, one way analysis of variance (ANOVA) and Pearson correlation analyses using SPSS 10.0 for Windows to establish significant variations and relationships.

RESULTS & DISCUSSION

The physico-chemical parameters of the seawater at the different study locations (Table 1) varied from 1.23 to 2.01 mg/L for NO_3^- , 0.08 to 0.20 mg/L for PO_4^{3-} , 31 to 33‰ for salinity, 6.36 to 7.35 for pH, 49.0 to 61.1 mg/L for COD, 29.0 to 31.5 mg/L for BOD and 3.1 to 5.9 mg/L for DO.

The identified phytoplankton species, whose relative abundance ranged from 2^{10} 516 Cell/mL, consisted of 83.33% diatoms and 16.67% dinoflagellates. They were diatoms *Coscnodiscus centralis*, *Asterionellops glacialis*, *Thalassionema frauenfeldii*, *Odontella mobiliensis*, *Chaetocero mitra*, *Rhizosolenia imbricate*, *Bacellaria paxillifera*, *Cerataulina palegica*, *Navicula spp*, *Pleurosigma spp* and dinoflagellates *Ditylum spp* and *Ceratium trichoceros*.

Phytoplankton mortality due to crude oil (Figs. 3 and 4) increased with concentration of oil irrespective of the exposure period. Generally, crude oil toxicity on the population varied from 0.06 to 36.43% for *C. centralis*, 1.41 to 35.58% for *C. trichoceros*, 1.71 to 46.11% for *T. frauenfeldii* and 0.66 to 44.90% for *O. mobiliensis*. Results revealed that the resistance to crude oil toxicity varied with phytoplankton species. For instance, Fig. 3 showed that at 6 h exposure, the toxicity decreased as follows: *T. frauenfeldii* > *O. mobiliensis* > *C. trichoceros* > *C. centralis*. However, at 12 h exposure, *C. centralis* exchanged position with *C. trichoceros* (*T. frauenfeldii* > *O. mobiliensis* > *C. centralis* > *C. trichoceros*). The same pattern was observed at 18 h contact period. Unfortunately, *C. trichoceros* did not survive beyond this time so became extinct due to natural mortality. Beyond 30 h, *O. mobiliensis* also became extinct and *T. frauenfeldii* could not survive beyond 36 hours so also became naturally extinct (Fig.4). leaving *C. centralis* as the only

Table 1. Physicochemical characteristics of the seawater, phytoplankton species composition and abundance at the study locations

Variable	Study location			
	Awoye	Ayetoro	Ilepete	Bar-Beach
Physicochemical parameters				
pH	7.19 ± 0.2	7.35 ± 0.01	7.35 ± 0.04	6.36± 0.01
Temperature (°C)	33±1	33±1	33±1	30±1
Salinity (‰)	31 ± 1	33± 1	33 ± 1	31 ± 2
Nitrate (mg/L)	1.23± 0.01	1.65±0.02	2.01±0.01	1.77±0.01
Phosphate (mg/L)	0.08 ± 0.01	0.18 0.03	0.10 ±0.01	0.20±
Electrical conductivity (mS/cm)	48.3± 0.2	52.0 0.1	52.0 ±0.3	48.8 ±0.2
Biochemical oxygen demand (mg/L)	31.5±0.4	29.0±0.1	28.3±0.2	29.9±0.3
Chemical oxygen demand (mg/L)	61.1± 0.5	57±0.2	53±0.2	49±0.4
Dissolved oxygen (mg/L)	3.1±0.1	3.7±0.1	3.6±0.2	5.9±0.1
Diatom (Cell/L)				
<i>Coscinodiscus centralis</i>	4.68 ± 1.0	3.14 ± 1.3	3.31 ± 0.1	5.16 ± 1.3
<i>Asterionellopsis glacialis</i>	1.37± 0.3	5.4 ± 0.3	2.2 ± 0.2	2.9 ± 4
<i>Thalassionema frauenfeldii</i>	1.62 ± 1.0	1.13 ± 0.2	9.6± 0.9	1.58 ± 0.6
<i>Odontella mobiliensis</i>	2.13 ±0.7	1.76 ± 1.5	2.0±0.1	1.1 ± 0.3
<i>Chaetoceros mitra</i>	9.6 ± 0.2	2.7 ± 0.2	3.6± 0.1	9.0 ± 0.5
<i>Rhizosolenia imbricate</i>	1.3± 0.1	8.0 ± 0.1	1.9± 0.1	2.9 ± 0.6
<i>Bacillaria paxillifera</i>	2.8 ±0.5	2.0 ± 0.1	1.7 ± 0.2	6.2 ± 0.7
<i>Cerataulina palegica</i>	1.66± 1.3	1.73 ± 1.2	1.6± 0.9	Nil
<i>Navicula species</i>	7.0± 1.0	6.0 ± 0.1	1.3 ± 0.6	1.6 ± 0.4
<i>Pleurosigma species</i>	8.0± 1.0	1.3 ± 0.1	1.6 ± 0.6	1.5 ± 0.3
Dinoflagellate (Cell/L)				
<i>Ditylum species</i>	5.1± 2	4.1 ± 1	4.9 ± 1	4.9 ± 0.5
<i>Ceratium trichceros</i>	5.4 ± 2	3.0±1	7.0 ± 1	1.14 ± 0.2

surviving organism at 42 h - contact with crude oil. Fig. 3 showed that of the four, *T. frauenfeldii* was the most vulnerable to crude oil toxicity within 24 - h contact.

In contrast to the increasing toxicity trend observed with regards to crude oil concentration, toxicity decreased with exposure period for each of the species (Fig.5). The relationships between the crude oil induced toxicity and the two variables (oil concentration and exposure period) showed high and significant correlations, with coefficients in the range of + 0.81 to + 0.97 ($p < 0.01$) for oil concentration and - 0.83 to - 0.90 ($p < 0.01$) for exposure period.

The physico-chemical parameters of the seawater used in this study were generally suitable for supporting lives in the aquatic ecosystem (Chindah, 2003). The dominance by diatoms (Order: Bacillariophyceae) was in line with the report of Wang *et al.* (2004) who stated that

diatoms are a major group of eukaryotic algae and are one of the most common types of phytoplankton, mostly unicellular, green in color, impacting green coloration to water and have a shell of silicon (Kolb, 1986). From previous Nigerian works, notable among phytoplankton found in the aquatic systems were Bacillariophyta, Chlorophyta, cyanophyta, Dinophyceae and Euglenophyta (Opute, 1990; Kadiri, 1993; Kemdirim, 2001; Ezra and Nwankwo, 2001; Ajayi and Akonai, 2003; Akin-Oriola, 2003; Chindah, 2003; Chindah, 2004; Nyananyo *et al.*, 2006; Essien *et al.*, 2006; Onyema, 2007; Akinyemi and Nwankwo, 2007; Essien *et al.*, 2007; Davies *et al.*, 2008).

The adverse impact of low level crude oil concentrations (6 to 50 mg/L) on the phytoplankton population and biodiversity showed toxicological effect of crude oil to these organisms, which are very sensitive ecological indicators, thereby




Fig. 3. Percentage toxicity of crude oil on population of C. centralis, C. trichoceros, T. frauenfeldii and O. mobiliensis as a function of concentration at a given exposure period




Fig. 4. Percentage toxicity of crude oil on population of C. centralis, C. trichoceros, T. frauenfeldii and O. mobiliensis at 30, 36 and 42 hour exposure periods




Fig. 5. Percentage toxicity of crude oil on population of *C.centralis*, *C.trchoceros*, *T.frauenfeldii* and *O.mobiliensis* as a function of at exposure period

demonstrating that crude oil input into the coastal waters is a risk factor to ecological status. A supporting evidence of crude oil toxicity to biota is found in the works of Obire and Anyanwu, (2009) which reported that in soils contaminated with concentrations of crude oil from 1 to 5%, higher concentrations of crude oil had adverse effect on fungal diversity.

As with most aquatic organisms, environmental stress due to the presence of xenobiotic pollutant in the habitat leads to altered physiological characteristics (in response to stress) and a greater degree of susceptibility to infections, disease and even death (Ruggiu *et al.*, 1998) so it probably was with the phytoplankton species in this study which resulted in mortality, though not 100%. The rate or extent to adjust to the crude oil induced-stress differed with the organisms. Results revealed that some phytoplankton species adapted

more easily than the others. In this regard, *C.trichoceros* (dinoflagellate) appeared to resist crude oil stress more than the other three (diatoms). This was attributed to the fact that diatoms (*T.frauenfeldii*, *C.centralis* and *O.mobiliensis*) have lower reproduction and assimilation rates in comparison to dinoflagellate *C.trichoceros* characterized by rapid sexual or asexual reproduction (Klepel, 1992).

The general reduction of phytoplankton population by the added crude oil was attributed to factors such as inhibition of food consumption, decrease in cell size, cell number and bio-volume (Miller *et al.*, 1978; Klepel, 1992). Others were inhibition of photosynthesis, primary production and respiration mechanisms owing to the presence of hydrocarbons (Gordon and Prouse, 1973; Dunsan *et al.*, 1975; Miller *et al.*, 1978; Karydis, 1979; Kusk, 1980; Karydis and Fogg, 1980; Karydis,

1981; Dahl *et al.*, 1983). In a short term bioassay experiment (Miller *et al.*, 1978), oil in water showed photosynthetic inhibition at 5 mg/L crude oil within 24 hours and concentrations as low as 2 mg/L caused up to 50% reduction in photosynthesis (Miller *et al.*, 1978; Dahl *et al.*, 1983). Crude oil consists of up to 80 to 90% hydrocarbons (both aromatic and paraffin fractions), which adversely affect biochemical factors such as protein, and sugar contents and photosynthetic pigments (chlorophyll a and b) of phytoplankton cells (Miller *et al.*, 1978; Karydis and Fogg, 1980). The adverse effect of habitat change on three of the species could be closely related to the transition between the marine environment enriched with nutrients and the controlled systems with limited nutrient supply (Burrige and Shir, 1995; Barron *et al.*, 2002; Folder and Burns, 2003; Chindah, 2004; Carayo *et al.*, 2005; Cermeno, 2008; Nubi *et al.*, 2008). Dinoflagellate *C.trichoceros* which was the most affected could be linked to the fact that it is heterotrophic, very motile and active thus required more food for survival in comparison to diatoms (*T.frauenfeldii*, *C.centralis* and *O.mobiliensis*) which are unicellular and less motile (Miller *et al.*, 1978; Klepel, 1992). Opute (1990) added that the majority of dinoflagellates as well as diatoms found in Nigerian coastal waters manifest stenohaline peculiarities as they could not tolerate a wide salinity range, indicating sensitivity to habitat changes which might explain the natural mortalities observed for the species outside of sea water environment.

The inverse relationship between phytoplankton mortality due to crude oil and exposure period suggest that the negative effect of single dose crude oil in this concentration range may be temporary. This is in concordance with the works of (Miller *et al.*, 1978) in which further prolonged exposure did not evoke further harm of crude oil to phytoplankton species in a bioassay experiment. The observation might be associated with the characteristics of Nigerian crude oil reported to be light (moderately viscous) and characterized with relatively high evaporation loss (Eyong *et al.*, 2004; Igwebuike *et al.*, 2007). Greater adverse impacts with time could likely emanate from chronic oil inputs into the environment. Reports from literature (Kusk, 1980; Karydis and Fogg, 1980; Burrige and Shir, 1995) showed that at non-

toxic levels of crude oil, phytoplankton metabolism could be enhanced due to increased nutrient availability from oil degradation, stimulation of nitrogen-fixing algae and bacteria or increased leaching of nutrients from vascular macrophyte stands, reporting that concentrations less or equal to 100 µg/L may stimulate growth but at up to 1.0 mg/L, growth in some species may be inhibited. The frequent occurrence of oil spills in the Niger Delta region of the country owing to aforementioned factors undermine the chances of the presence of crude oil at non-toxic levels in this area (Awobajo, 1981; Ekekwe, 1981; Ibiebele, 1986).

CONCLUSION

Crude oil concentration in the range of 6.0 to 50mg/L evoked toxicity on four Nigerian indigenous phytoplankton species. In summary, resilience to habitat change from marine environment to laboratory conditions decreased in the order of *C.centralis* > *T.frauenfeldii* > *O.mobiliensis* > *C. trichoceros* but in relation to crude oil toxicity within 24-h contact time, vulnerability decreased in the order *T.frauenfeldii* > *O.mobiliensis* > *C.centralis* > *T.trichoceros* for the four species. This is a potential danger to phytoplankton species biodiversity conservation. Further toxicological research on these and other species, effects mediated through food web interactions as well as a field scale study incorporating natural ecosystem factors are recommended.

REFERENCES

- Adekambi, O.E. (1989). Petroleum hydrocarbon pollution of Nigerian waters and sediments around Lagos and Niger Delta area of Nigeria. Ph.D Thesis, Department of Chemistry, University of Ibadan, Ibadan, Nigeria.
- Adekunle, I.M., Arowolo, T.A., Omoniyi, I.T. and Olubambi, O.T. (2007a). Risk Assessment in the Nile tilapia (*Oreochromis niloticus*) and African mud catfish (*Clarias gariepinus*) exposed to cassava effluent. *Chemistry and Ecology*, **23**(5), 383-392.
- Adekunle, I.M., Adetunji, M.T. Gbadebo, A.M. and Banjoko, O.B. (2007b). Assessment of groundwater quality in a typical rural settlement in southwest Nigeria. *Int. J. Environ. Res. Public Health*, **4**(4), 307-318.
- Adeniyi, O.D., Nwalor, J.U. and Ako, C.T. (2008). A review on water flooding problems in Nigeria crude oil

production. *Journal of Dispersion Science and Technology*, **29(3)**, 362 – 365 .

Ajayi, A.O. and Akonai, K.A. (2003). Physico-chemical properties and microbial ecology of Lagos lagoon, Nigeria. *Biosci. Res. Comm.*, **15(6)**, 453 - 462.

Ajayi, A.O., Balogun, S.A. and Adegbehingbe, K. (2007). Microorganisms in the crude oil-producing areas of Ondo State, Nigeria. *Scientific Research and Essay*, **3(5)**, 174 -179.

Akin-Oriola, G.A. (2003). On the phytoplankton of Awba reservoir, Ibadan, Nigeria. *Rev. Bil. Trop.*, **51(1)**, 99 -106.

Akinyemi, S.A and Nwankwo, D.I. (2007). Effects of seasons on phytoplankton of the Asejire Reservoir (Nigeria). *International Journal on Algae*, **9**, 81 -88.

American Public Health Association (1989). Standard methods for the examination of water and wastewater, 20th edition, AOHA, AWWA, WEF, Washington DC.

Awobajo, S.A. (1981, November). An analysis of oil spill accidents in Nigeria: 1976 – 1980. In: *Proceedings of the Petroleum Industry and the Nigerian Environment*. Nigeria National Petroleum Corporation (NNPC), Warri, 89 – 94 .

Benson, N.U., Essien, J.P., Ebong, G. A. and Williams, A.B. (2008). Petroleum hydrocarbons and limiting nutrients in *Macura repentia*, *Procambius clarkia* & benthic sediment from Qua iboe Esturay, Nigeria. *The Environmentalist*, **28(3)**, 275 -282.

Barron, S., Weber, C., Marid, R., Davidson, E., Tomasky, G. and Howarth., R. (2002). Effects of varying salinity on phytoplankton growth in a low –salinity coastal pond under two nutrient conditions, *Biol. Bull.*, **203**, 260 -261.

Burridge, T.R and Shir, M. (1995). The comparative effects of oil dispersants and oil/dispersant conjugates on germination of the marine macroalga *Phyllospora comosa* (Fucales:Phaeophyta). *Marine Pollution Bulletin*, **3(4-12)**, 446 - 452.

Carayo, C.C., Culango, K.C., Estrera, D., Largo, D.B. and Dy, D.T. (2005). Correlation of inorganic nutrients to the standing crop of green tide algae during the peak growth season in eastern Mactan Island, central Philippine. *Philipp. Scientist*, **42**, 1- 4.

Cermeno, P. (2008). Resource levels, allometric scaling of pollution abundance, and marine phytoplankton diversity. *Limnol. Oceanogr.*, **53(1)**, 312 – 318.

Chindah, A. C. (2004). Response of periphyton community to salinity gradient in tropical estuary, Niger Delta. *Pollution Journal of Ecology*, **52(10)**, 83 -89.

Chindah, A. C. (2003). The physico-chemistry, phytoplankton and periphyton of swan forest stream in the lower Niger, delta. *Scientia Africana*, **1(2)**, 1-15.

Dahl, E., Laake, M., Tjessem, K., Eberlein, K. and Bohle, B. (1983). Effects of Ekofisk crude oil on an enclosed planktonic ecosystem. *Marine Ecology*, **14**, 81 – 91.

Davies, A., Abolude, Z.D.S and Ugwumba, A.A. A. (2008). Phytoplankton of the lower reaches of Okpoka creek, PortHarcourt, Nigeria. *Journal of Fisheries International*, **3(3)**, 83 - 90.

Dunsan, W.M., Atkinson, L.P. and Natok, J. (1975). Simulation and inhibition of phytoplankton growth by low molecular weight hydrocarbons. *Marine Biology*, **31(4)**, 305 -310.

Dzierzbicka-Glowacka, L. (2007). Effect of phytoplankton mortality caused by unpredictable conditions – numerical simulations. *Polish Journal of Ecology*, **35(1)**, 27 -37.

Egberongbe, F.A.O., Nwilo, P.C. and Badejo, O.T. (2006). Oil spill disaster monitoring along Nigeria coastline, 5th FIG Regional Conference, Accra, Ghana, March 8-11, Retrieved January, 23, 2008 from http://www.fig.net/pub/accra/ts16/ts16-06_egberongbe_etal.pdf

Ekekwe, E. (1981, November). The Funiwa-5 oil well blow out. In: *Proceedings of International Seminar on the Petroleum Industry and Environment*. Nigeria National Petroleum Corporation (NNPC), Warri, 19 -22.

Ekom, R.A. (2006). Effect of waste engine oil on phytoplankton of the Calabar River Estuary, Nigeria. *Global Journal of Environmental Sciences*, **292**, 112- 117.

Essien, J.P., Antai, S.P and Benson, N.U. (2007). Microalgae biodiversity and biomass status in Qua Iboe Estuary mangrove swamp, Nigeria. *Aquatic Ecology*, **42(1)**, 71 -81.

Essien, J.P., Ubom, R.M and Antai, S.P. (2006). Productivity and distribution of Epipellic microalgae along salinity gradients in mangrove swamp of the Qua Iboe estuary, Nigeria. *Environmental Monitoring Assessment*, **121(1-3)**, 65-75.

Eyong, E.U., Umoh, I.B., Ebong, P.E., Eteng, M.U. Antai, A.B. and Akpan, A. (2004). Haematotoxic effects following ingestion of Nigerian crude oil and crude oil polluted shellfish by rats. *Nigerian Journal of Physiological Sciences*, **19(1-2)**, 1 - 6.

Ezra, A.G and Nwankwo, D. (2001). Composition of phytoplankton algae in Gubi reservoir, Bauch, Nigeria. *Journal of Aquatic Sciences*, **16(2)**, 115 -118.

Folder, S. and Burns, C. W. (2003). Phytoplankton diversity of shallow tidal lakes; influence of periodic salinity changes on diversity and species number of a natural assemblage. *Journal of Phycology*, **40(1)**, 54 - 61.

- Gordon, D.C. and Prouse, N.J. (1973). The effects of three oils on marine phytoplankton photosynthesis. *Marine Biology*, **22**, 229 -333.
- Ibiebele, D.D. (1986). Point-source inputs of petroleum waste into the Niger Delta, Nigeria. *Sci. Tot. Environ.*, **52**, 233 -238.
- Igwebuike, U.M., Obidike, R.I., Shoyinka, S.V.O., Nwankwo, C.U., Okwechima, I.O. and Aka, L.O. (2007). Effect of Nigeria Qua Iboe Brent crude oil on rat haematological parameters *Vet. Arhiv*, **77**, 247 – 256.
- Isichei, A.O. and Sanford, W.W. (1979). The effects of waste flares on the surrounding vegetation in south – eastern Nigeria. *Journal of Applied Ecology*, **13**, 177 – 187.
- Jinadu, A.K. (1989). Petroleum hydrocarbons of fish and shellfish around Lagos and Niger Delta area of Nigeria. Ph.D Thesis, Department of Chemistry, University of Ibadan, Ibadan, Nigeria.
- Kadiri, M.O. (1993). Seasonal changes in the phytoplankton biomass of shallow tropical reservoir, Nigeria. *Journal of Botany*, **6**, 167 – 175.
- Kakulu, S.E. and Osibanjo, O. (1986). A baseline study of mercury in fish and sediments in the Niger Delta area of Nigeria. *Environ. Pollut. (Series B)*, **11**, 315 – 322.
- Kakulu, S.E. and Osibanjo, O. (1988). Trace heavy metal pollutional status in sediments of the Niger Delta Area of Nigeria. *J. Chem. Soc., Nigeria*, **13**, 9-15.
- Kakulu, S.E., Osibanjo, O. and Ajayi, S.O. (1987). Trace metal content of fish and shellfish around Lagos and Niger Delta area of Nigeria. *Environ. Int.*, **13**, 247 – 251.
- Karydis, M. (1979). Short term effects of hydrocarbons on the photosynthesis and respiration of some phytoplankton species. *Botanica Marina*, **22(5)**, 281 – 285.
- Karydis, M. and Fogg, G.E. (1980). Physiological effects of hydrocarbons on the marine diatom *Cyclotella cryptica*. *Microbial Ecology*, **6(4)**, 281 -290.
- Karydis, M. (1981). The toxicity of crude oil for marine alga *Skeletonema costatum* (Greville) Cleve in relation to nutrient limitation. *Hydrobiologia*, **85(2)**, 137 – 143.
- Kelly, L.D., McGuinness, L.R., Hughes, J.E. and Wainright S.C. (1999). Effects of phenanthrene on primary production of phytoplankton in two New jersey Estuaries. *Bull. Environ. Contam. Toxicology*, **63**, 646 – 653.
- Klepel, G.S. (1992). Environmental regulation of feeding and egg production by *Acartia tonsa* off southern California. *Marine Biology*, **112(1)**, 57 -65.
- Kemdirim, E.C. (2001). Checklist of phytoplankton of shedan reservoir in Plateau State. Nigeria *Jornal of Aquatic Sciences*, **16**, 61 -63.
- Khan, M.A & Ejike, C .1984. Limnology and plankton periodicity of Jos Plateau water reservoir, Nigeria, West Africa. *Hydrobiologia*, **114(3)**, 189 – 199.
- Kolb, J.A. (186). *Marine Biology and Oceanography Grades 9-12*. marine Science project: For sea marine Science center, Poulsbo, WA.
- Kusk, K.O. (1980). Effects of Crude oil and aromatic hydrocarbons on the photosynthesis of the diatom *Nitzschia palea*. *Physiologia Plantarum*, **43(1)**, 1-6.
- Miller, M.C., Alexander, V. R. and Barsadate, J. (1978). The effects of oil spill on phytoplankton in Arctic lake and ponds. *Artic*, **31(3)**, 192 -218.
- Nubi, O.A., Ajao, E. A., Oyewole, E.O and Unyimadu. J.P. (2008). Nutrient levels in Guinea current large marine ecosystem (GC_LME) waters. *Science World Journal*, **3(2)**, 89 – 84.
- Nyananyo, B.I., Okeke, C.U and Mensah, S.I. (2006). Physico-chemical conditions and distribution of phytoplankton in the Brass River, Nigeria. *J. Appl. Sci. Environm. Mgt.*, **10(1)**, 43- 45.
- Obire O and Anyawu, E.C. (2009). Impact of various concentrations of crude oil on fungal populations oof soil, *Int.J. Environ. Sci. Tech.*, **6(2)**, 211-218.
- Onyema, I.C. (2007). The phytoplankton composition, abundance and temporal variation of a polluted estuarine creek in Lagos, Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences*, **7**, 89 -96.
- Opute, F.I. (1990). Phytoplankton flora in the Warri/ Forcados estuary of southern Nigeria, *Hydrobiologia*, **208(1-2)**, 101 – 109.
- Powel, C.B., White, S.A., Baranowska, B., Ibiele, D.D., Isoun, M. and Ofoegbu., F.U. (1985, November). Oshika oil spill environmental impact; effect on aquatic biology in the petroleum industry and the Nigerian environment: Proceedings of an International Seminar, Kaduna. Published by the Petroleum Inspectorate, Nigeria National Petroleum Corporation (NNPC), and the Environmental Planning and Protection Division of the Federal Ministry of Works and Housing.
- Ruggiu, D. Morabito, G., Panzani, P and Pugnetti. A. (1998). Trends and relations among basic phytoplankton characteristics in the course of the long-term oligotrophication of Lake maggiore (Italy). *Hydrobiologia*, 369 & 370: 243 – 257.
- Snowden, R.J. and Ekweozor, I.K.E. (1987). The impact of minor oil spillage in the estuarine Niger Delta. *Marine Pollution Bulletin*, **18(11)**, 595 -599.
- Wang, Y., Lou, Z., Sun, C., Wu, M and Han, S. (2004). Multivariate statistical analysis of water quality and phytoplankton characteristics in Daya Bay, China from 199 to 2002. *Oceanologia*, **48(2)**, 193 -211.