

Estimation of Urban Suspended Particulate Air Pollution Concentration

Halek, F. ^{1*}, Keyanpour, M. ¹, Pirmoradi, A. ² and Kavousi, A. ¹

¹ Environmental group, Dept. of Energy, Materials and Energy Research Center,
P.O.Box 14155-4777, Tehran, Iran

² Graduate Faculty of Environment, University of Tehran & GIS Department, National
Cartographic Center, Tehran, Iran

Received 7 Nov. 2008;

Revised 26 Aug. 2009;

Accepted 15 Sep. 2009

ABSTRACT: A critical eye on the destructive impact of air pollution in Tehran is needed as the basis for urban planning, protection policy and management. This paper is focused on modeling in the GIS (Geospatial Information System) to estimate the concentration of particulate matter (PM) in any point of a typical part of Tehran which extends over 18.2 km² and includes the so-called "Traffic Zone". Many important general hospitals are located in this region, some of which are within the zone. The model is built on the data obtained in 42 stations located within the region. The results strongly indicate that the concentrations of PM₁₀, PM_{2.5} and PM_{1.0} of any points inside the region, including the traffic zone, do not meet the required international standard values. The extracted estimate values for the 22 hospitals reveal that the concentration of PM₁₀ for "Azadi Psychic", "Children" and "Mustafa Khomeini" hospitals are the worst, estimating from the model to be 119.42 µg/m³, 107.09 µg/m³ and 101.14 µg/m³ respectively. The percent ratio of the mean concentrations of PM₁₀/PM_{2.5}/PM_{1.0} in this region is found to be approximately 7: 2: 1.

Key words: GIS, Dynamic modeling, Particulate matter, Air pollution, Traffic

INTRODUCTION

Pollution can be defined as an undesirable change in the physical, chemical or biological characteristics of the air, water or land that can affect health, survival or activities of humans or other organisms (Smith, 1996; WHO, 1997; Patel and Raiyani, 1995; Anu *et al.*, 2002; anthony *et al.*, 2007; Minsi *et al.*, 2007; Marilena & Elias, 2008). Among the major air pollutants released to the atmosphere, suspended particulate air pollution are considered as one of the major health impact and therefore a large number of related studies have been undertaken in developing countries in the last decade (Cautreels & Van, 1978; Zhu *et al.*, 2002; Douglas *et al.*, 2002; Alam *et al.*, 2003; Gramotnev & Ristovski, 2004; Silibello *et al.*, 2008). The most important environmental problem Iran currently faces is air pollution, especially in the capital city of Tehran. The problem is very serious for the city

which considered one of the most polluted cities in the world. Cars are chiefly to blame for Tehran's heavy pollution, because most of the city's more than 2 million cars are at least 20 years old and do not have catalytic converters to reduce pollutants. Unfortunately the city's geographical position is not helping the reduction of pollutants. The city is hammed by the tall Alborz Mountains to the north and therefore traps the pollutants over the city (Halek *et al.*, 2004).

Particulate matter is considered one of the main sources of air pollution problems in Tehran. The role, size distribution of particulate matter in the city's air pollution and also the effect of motor vehicles and trend of air borne particulate, have been the subject of extensive studies (Nabi and Halek, 2007). In air pollution studies, the air quality models are used to predict and estimate concentration of one or more species in space

*Corresponding author E-mail: f-halek@merc.ac.ir

and time as related to the dependent variables. Modeling provides the ability to assess the current and also future air quality in order to enable “informed” policy decisions to be made (Bruckman *et al.*, 1992; Hochadel *et al.*, 2006; Zhou *et al.*, 2006; Gavin *et al.*, 2007; Yuqiong *et al.*, 2008).

One of the systems which have appeared lately is Geospatial Information System (GIS). GIS is not only a system for creating, managing and analyzing graphic and attribute data, but also is a decision supporting system (DSS). In fact, GIS can support managers, planner and decisions maker. Therefore, these days we will face big problems in big cities if we don't use such systems (Pirmoradi, 2008). The rise of GIS technology and its use in a wide range of disciplines provides transportation and air quality modelers with a powerful tool for developing new analysis capability (Goodchild *et al.*, 1996; Burroughs and McDonald, 1998; Appleton and Lovett, 2003; Tolga, 2004; Duanping *et al.*, 2006; Younes *et al.*, 2008). The organization of data by location allows data from a variety of sources to be easily combined in a uniform framework (Wilfred and Gerald, 2005; Mauro and Lorenzo, 2006).

A Comparison and case study conducted to apply the satellite data and GIS for producing maps of amounts of CO, O₃, NO₂ and SO₂ in Tehran's atmosphere (Sohrabinia and Khorshiddoust, 2007). With the help of GIS, concentrations of each of these pollutants were estimated to be much higher than standard values and forecasted that to go still higher. The results of such a study and other air pollution case studies in different countries not only could help the local, but the global environmental pollution experts and decision makers to set environmental politics. So for the application of GIS to estimate the air pollution in Tehran has been limited to find the concentrations of gaseous pollutants and not suspended particles (Sohrabinia and Khorshiddoust, 2007). This paper reports result of a novel study to estimate the concentration and spatial distribution of PM₁₀, PM_{2.5} and PM_{1.0} in an 18.2 km² region selected within the Great Tehran. For this purpose, the study is focused for the first time on modeling in the GIS which used data extracted from 42 stations located in different parts of the region. Special attention is made to estimate the concentrations of particulate matter near 22 hospitals in this

region, some of which are located in the so-called “Tehran's Traffic Zone”. With the help of this modeling, one could estimate the concentrations of PM₁₀, PM_{2.5} and PM_{1.0} in such important areas in this region where setting a station is impossible for sampling.

MATERIALS & METHODS

A region of 18.2 km² was used as our study area because many important general hospitals, including “Tehran heart center”, “Imam Khomeini”, “Shariati”, “Arya”, “Sasan”, “Sajad”, “Pars”, “Mustafa Khomeini”, “Toos” and “Rasoul Akram” are located in this area. Also in this area, a so-called “Traffic Zone” has been set up since 2000, covering the city center during peak traffic hours. Entering and driving inside this zone is only allowed with the special permit. Fig. 1 shows this region in Tehran. 42 sampling sites were set to collect the air samples according to the standard sampling procedures. Daily sampling were done in summer 2007, starting 8.0 AM (the morning rush hour) to 4.0 PM (the afternoon rush hour). The names of the sampling sites and their related geographical coordinates in Tehran are listed in Table 1. The meteorological data were reported by the Iranian Meteorological Organization. The mean temperature in the selected district in Tehran during the study fluctuates between 29-35 °C.

A portable Grimm aerosol spectrometer, model 1.108, which equipped with 15 ports, has been used for sampling. This instrument was able to measure the size distribution of particulate matter in urban and also in the industrial areas. The air flow was set at 1.2 Lit/min. and the particles were collected on PTFE filters. In order to build a “Surface Model” for PM₁₀, PM_{2.5} and PM_{1.0}, different algorithms should be applied to interpolate the data from those obtained for the known sites and extend the results to the “surface”. For this purpose mean concentrations of PM₁₀, PM_{2.5} and PM_{1.0}, in each sampling site for June, July and August, were calculated and interpolated and extended to the surface by “Inverse Distance Weight” or “Spine” algorithms by using ArcGIS9.2. To examine the precision of the surface models, the “Root Mean Square” method was applied. After building the surface models for each month in the summer, the concentration of each particulate matter could be estimated in all the points in the district and




Fig. 1. The border of sampling points

consequently, the related distribution of the particles and their concentration could be evaluated and categorized. Also with the aid of “Mathematics Overlay”, the models for these months could be overlaid and the final results for the particles could be evaluated for summer 2007.

RESULTS & DISCUSSION

The mean concentrations of particulate matter in 42 sampling stations in summer 2007 are shown in Fig. 2. The higher values belong to those crowded places like city squares and street intersections.

As it is shown in Fig. 2. the concentration of PM_{10} is much higher than the rest of particles in Tehran's air pollution. The percent ratios of different categories of particulate matter are found from Fig. 2 to be, 73 % (PM_{10}), 17 % ($PM_{2.5}$) and 10 % ($PM_{1.0}$). The suspended particulate matter concentration, as shown in Fig. 3. is higher for working days, especially on Saturdays which considered as the beginning of the week in Iran, the roads are usually congested with heavy traffic. The concentration of PM_{10} in the selected district along with the locations of the stations is shown in the related distribution map in Fig. 4. The darker the regions in this figure, the higher the concentration of particulate matter. The mean

value of particulate concentration is calculated from the model to be 79.49 for the corresponding summer, having standard deviation of 13.63. It could be deduced from this figure that the highest concentration of the $123.19 \mu\text{g}/\text{m}^3$ for PM_{10} belong to the intersection of “Sheik Fazlollah” and “Jenah” expressways, with and the least belongs to the intersection of “Saeedi” and “Hashemi” expressways, with $63.76 \mu\text{g}/\text{m}^3$.

Similarly, the mean values of concentration of $PM_{2.5}$ and $PM_{1.0}$ are shown in Fig. 5 and Figure 6 respectively. As shown in Fig. 5. the mean value of concentration of $PM_{2.5}$ is recorded to be $19.20 \mu\text{g}/\text{m}^3$, corresponding to the highest value of $45.30 \mu\text{g}/\text{m}^3$ and the lowest of $14.00 \mu\text{g}/\text{m}^3$. The standard deviation of the concentration of $PM_{2.5}$ is calculated to be 3.39. $PM_{1.0}$ concentration, as shown in Fig. 6. has the highest value of $34.80 \mu\text{g}/\text{m}^3$ and the least value of $6.67 \mu\text{g}/\text{m}^3$. The mean value for the concentration of this size of suspended particles is calculated to be 10.80 with the standard deviation of 2.44. As it is deduced from the mean concentration of particulate matter, Tehran, like many big cities, suffers from sever air pollution and therefore the city is often covered by smog making breathing difficult and causing widespread pulmonary illnesses. Unfortunately most of the hospitals in Tehran are located in the

Table 1. The names of the sampling sites and their related geographical coordinates

Site No.	Sampling Site Location:	X	Y
1	Karaj Road	529,208.36	3,950,734.79
2	Saeedi Highway - Dampezhshki intersection	530,759.07	3,949,955.15
3	Fath Square	530,881.52	3,948,624.39
4	Hashemi St. - Jeyhoon intersection	532,909.85	3,949,672.37
5	Dampezhshki St. - Karoon intersection	533,149.77	3,950,075.11
6	Dampezhshki St. - Moeen intersection	531,281.77	3,949,916.59
7	Azadi Square	530,913.31	3,950,790.60
8	Azadi St. - Sharif University	531,750.28	3,950,820.79
9	Azadi St. - Azarbajjan intersection	533,025.93	3,950,866.07
10	Azarbajjan St. - Roudaki intersection	533,948.99	3,950,433.25
11	Islamic Republic Square	534,259.18	3,950,482.14
12	Islamic Republic St. - Jamalzadeh intersection	535,217.38	3,950,436.52
13	Islamic Republic St. - Fakhre-Razi intersection	535,817.68	3,950,402.38
14	Islamic Republic St. - Valii Asr intersection	536,518.47	3,950,368.43
15	Valii Asr Junction	536,585.26	3,950,969.58
16	Enghelab St. - Tehran University	535,801.54	3,950,987.39
17	Enghelab Square	535,382.96	3,950,996.30
18	Azadi St. - Kaveh Parking	534,975.07	3,950,969.12
19	Azadi St. - Eskandari intersection	534,415.68	3,950,940.62
20	Azadi St. - Roudaki intersection	533,899.05	3,950,929.93
21	Khosh St. - Nosrat intersection	533,555.94	3,951,443.21
22	Sattar Khan St. - Behbudi intersection	533,486.26	3,952,164.38
23	Touhid Square	534,109.88	3,951,526.83
24	Chamran Highway - Bagher Khan intersection	534,175.64	3,951,945.33
25	Keshavarz Blv. - Imam Hospital	534,672.10	3,951,727.59
26	Keshavarz Blv. - Kargar intersection	535,307.38	3,951,675.23
27	Keshavarz Blv. - Hejab intersection	535,924.16	3,951,808.26
28	Keshavarz Blv. - Felestin intersection	536,504.66	3,952,005.79
29	Valii Asr St. - Taleghani intersection	536,657.85	3,951,550.26
30	Valii Asr Square	536,790.88	3,952,074.32
31	Valii Asr St. - Zartosht intersection	536,878.83	3,952,766.31
32	Fatemi Square	536,578.55	3,953,064.25
33	Golha Square	536,239.59	3,953,324.69
34	Gomnam Highway - Kurdistan intersection	535,731.10	3,953,601.66
35	Gomnam Highway - Kargar intersection	535,168.89	3,953,436.30
36	Gomnam Highway - Chamran intersection	534,436.39	3,953,698.72
37	Jalal Highway - Sheik Fazlollah intersection	533,348.68	3,954,133.80
38	Sheik Fazlollah Highway - Sattar khan Bridge	532,459.25	3,953,103.95
39	Sheik Fazlollah Highway - Yadekar Bridge	531,830.08	3,952,466.45
40	Sheik Fazlollah Highway - Jenah intersection	530,209.12	3,952,313.70
41	Jenah Highway - Fuel Pump Station	530,452.05	3,951,640.99
42	Azadi Square - Bus Terminal	530,468.40	3,951,129.45




Fig. 2. Mean concentration of suspended particles in 42 site




Fig. 3. Mean of Particulate Matter Concentration in week days

selected district, instead of being placed in the non-polluted areas. Fig. 1 shows the geographical position of the selected district in Tehran. From the aid of GIS database, the concentration of particulate matter in any selected points, especially those for hospitals located in this district could be estimated. The names and places of the 22 hospitals in the selected district with their mean concentrations of PM_{10} are listed in Table 2. As the mean concentrations of PM_{10} indicate, none of the hospitals meet the required international standard air pollution values. The extracted estimated values from GIS modeling revealed that the concentration of PM_{10} for “Azadi Psychic”,

“Children” and “Mustafa Khomeini” hospitals are the worst, estimated from the model to be $119.42 \mu\text{g}/\text{m}^3$, $107.09 \mu\text{g}/\text{m}^3$ and $101.14 \mu\text{g}/\text{m}^3$ respectively.

CONCLUSION

Environmental modeling for an 18.2 km^2 selected district in Tehran is developed to assess the critical state of particulate matter concentration in an important part of Tehran, where the so-called “Traffic Zone” is located and also many hospitals, like “Azadi Psychic” and “Children” are continuing their daily activities. The annual average concentration of PM_{10} and $PM_{2.5}$ in each 42 site




Fig. 4. The distribution of mean concentration of PM_{10} in the selected region




Fig. 5. The distribution of mean concentration of $\text{PM}_{2.5}$ in the selected region

and also in any point in the selected district exceed the National air Quality Standard (NAQS) of $50 \mu\text{g}/\text{m}^3$ and $15 \mu\text{g}/\text{m}^3$ respectively. The mean concentrations of $\text{PM}_{1.0}$, $\text{PM}_{2.5}$ and PM_{10} are found to be $13.14 \mu\text{g}/\text{m}^3$, $22.67 \mu\text{g}/\text{m}^3$ and $95.72 \mu\text{g}/\text{m}^3$ respectively. The highest concentration of PM_{10} , which is found to be $529.24 \mu\text{g}/\text{m}^3$, belongs to “Valii Asr” square, which is also the worst square for $\text{PM}_{2.5}$ ($105.88 \mu\text{g}/\text{m}^3$). The highest

concentration of $\text{PM}_{1.0}$, ($89.87 \mu\text{g}/\text{m}^3$) belongs to “Islamic Republic Square” in Tehran.

REFERENCES

- Alam, A., Shi, J. P. and Harrison, R. M. (2003). Observations of new particle formation in urban air. *J. Geoph. Res.*, **108**, 4093-4107.
- Anthony, L., Barry, S., Anthony, B. and Hugo, Z., (2007). Reducing the healthcare costs of urban air pollution: The South Africa experience. *J. Environ. Manag.*, **84**, 27-37.




Fig. 6. The distribution of mean concentration of PM_{10} in the selected region

Table 2. Names, locations and mean concentration of PM_{10} of the hospitals

	Hospital Name	Location	Mean Concentration of PM_{10} ($\mu g/m^3$)
1	Aria	Vesal St.	99.58
2	Artesh (501)	Fatemi St.	73.01
3	Artesh (503)	Ostad Moeen St.	83.68
4	Azadi	Azadi Sq.	95.76
5	Albourz	Vesal St.	82.31
6	Imam Khomeini	Keshavarz Blvd.	78.06
7	Cancer Institute	Keshavarz Blvd.	74.77
8	Pars	Keshavarz Blvd.	99.60
9	Rasoul Akram	Sattarkhan St.	70.31
10	Ravan Pezeshki	Jenah Highway	119.42
11	Sasun	Keshavarz Blvd.	96.76
12	Central 1	Bagherkhan St.	75.53
13	Sajjad	Fatemi Sq.	93.24
14	Shriati	Kargar St.	62.38
15	Firouzgar	Taleghani St.	97.26
16	Ghalb	Kargar St.	65.83
17	Kudakan	Taleghani St.	107.09
18	UT Health Center	Enghelab St.	90.69
19	Tebbi kudakan	Dr. Ghareeb St.	77.57
20	Mustafa Khomeini	Italia St.	101.14
21	Mehr	Zartosht St.	92.73
22	Meymanat	Azadi Sq.	98.22

- Anu, K., Jaakko, K., Ari, K., Paivi, A. and Tarja, K. (2002). A model for evaluating the population exposure to ambient air pollution in an urban area. *Atmos. Environ.*, **36**, 2109-2119.
- Appleton, K. and Lovett, A. (2003). GIS-based visualization of rural landscapes: defining sufficient realization for environmental decision-making. *Lands. Urban Plan.*, **65**, 117-131.
- Bruckman, L., Dickson, R. J. and Wilkosen, J. G. (1992). Use of GIS software in the development of emissions inventories and emission modeling. *Proceedings of the Air & Waste Management Association*, Pittsburg, PA, USA.
- Burrough, P. A. and McDonald, R. A. (1998). *Principles of geographical information systems*. Oxford University Press, **333**, Oxford.
- Cautreels, W. and Van, Cauwenberghe, K. (1978). Experiments on the distribution of organic pollutants between airborne particulate matter and the corresponding gas phase. *Atmos. Environ.*, **12**, 1133-1141.
- Douglas, W. D., Joel, S. and John, D. S. (2002). Air Pollution and Daily mortality: Associations with particulates and acid aerosols. *Environ. Res.*, **69**, 362-373.
- Duanping, L., Donna, P., Yinkang, D., Eric, W., Jianwei, D., Richard, S., Hung-mo, L., Jiu, C. and Gerardo, H. (2006). GIS approaches for the estimation of residential-level ambient PM concentrations. *Environmental Health Perspectives*, **114**, 1374-1380.
- Gavin, C., Gareth, J., Malkolm, H. and Stephen, D. (2007). Predictive uncertainty in environmental modeling. *Neural Net.*, **20**, 537-549.
- Goodchild, M. F. (1996). *GIS and environmental modeling. Progress and research issues*, Fort Collins, CO; GIS World Book, 486.
- Gramotnev, G. and Ristovski, Z., (2004). Experimental investigation of ultra fine particle size distribution near a busy road. *Atmos. Environ.*, **38**, 1767-1777.
- Halek, F., Kavousi, A. and Montahaie, H. (2004). Role of motor-vehicles and trend of air borne particulate in the Great Tehran area, Iran. *Int. J. Environ. Health Res.*, **14**, 301-307.
- Hochadel, M., Heinrich, J., Gehring, U., Morgenstern, V., Kuhlbusch, T., Link, E., Wichmann, H. E. and Kramer, U. (2006). Predicting long-term average concentrations of traffic-related air pollutants using GIS-based information. *Atmos. Environ.*, **40**, 542-553.
- Marilena, K. and Elias, C. (2008). Human health effects of air pollution. *Environ. Poll.*, **151**, 362-367.
- Mauro, D. D. and Lorenzo, B. (2006). MAPIT: The GIS software for field mapping with tablet pc. *Comp. Geosci.*, **32**, 673-680.
- Minsi, Z., Yu, S. and Xuhui, C. (2007). A health-based assessment of particulate air pollution in urban areas of Beijing in 2000-2004. *Sci. Tot. Environ.*, **376**, 100-108.
- Nabi, Bidhendi, G. R. and Halek, F. (2007). Aerosol Size Segregated of Tehran's atmosphere in Iran. *Int. J. Environ. Res.*, **1**, 58-65.
- Patel, T. S. and Raiyani, C. V. (1995). Indoor air quality: problems and perspectives. In: *Energy Strategies and Green House Gas Mitigation*. Ed. P. R. Shukla, New Delhi, 72.
- Pirmoradi, A. (2008). Creating compatibility matrix of educational use and suitable site selection of new school for urban resource management and supporting decision. 4th GIS Conference along with ISPRS Workshop, 6-7 January 2008, Tehran-Iran.
- Silibello, C., Calori, G., Brusasca, G., Giudici, A., Angelino, E., Fossati, G., Oeroni, E. and Buganza, E. (2008). Modeling of PM₁₀ concentrations over Milano urban area using two aerosol modules. *Environmental Modelling & Software*, **23**, 333-343.
- Smith, K. R. (1996). Indoor air pollution in developing countries: Growing evidence of its role in the global disease burden. In: *Indoor Air 96, proceedings of 7th international conference on Indoor Air and Climate*. Ed. K. Ikeda and I Iwata. Institute of Public Health, Tokyo, 33.
- Sohrabinia, M. and Khorshiddoust, A. M. (2007). Application of satellite data and GIS in studying air pollutants in Tehran. *Habitat Int.*, **31**, 268-275.
- Tolga, E. (2004). A GIS based decision support system for estimation, visualization and analysis of air pollution for large Turkish cities. *Atmos. Environ.*, **38**, 4509-4517.
- Wilfred, W. and Gerald, S. (2005). Environmental software systems for emission inventories. *Environmental Modelling & Software*, **20**, 1469-1477.
- World Health Organization European, (1997). *Regional office for Europe, Copenhagen. Air Quality Guidelines for Europe, Series no 23*.
- Younes, N., Ryuichi, I., Hikari, F. and Toshiaka, T. (2008). GIS integration model for geothermal exploration and well siting. *Geothermics*, **37**, 107-131.
- Yuqiong, L., Hoshin, G., Everett, S. and Thorsten, W. (2008). Linking science with environmental decision making: Experiences from an integrated modeling approach to supporting sustainable water resources management. *Environmental Modelling & Software*, **23**, 846-858.
- Zhou, P., Ang, B. W. and Poh, K. L. (2006). Decision analysis in energy and environmental modeling: An update. *Energy*, **31**, 2604-2622.
- Zhu, Y., Hinds, W. C., kim, S. and Sioutas, S. (2002). Concentration and size distribution of ultrafine particles near a major highway. *J. Air & Waste Manag. Assoc.*, **52**, 1032-1042.