

## Water Quality Assessment in Lakes of Vojvodina

Pamer, E. <sup>1\*</sup>, Vujovic, G. <sup>3</sup>, Knezevic, P. <sup>1</sup>, Kojic, D. <sup>1</sup>, Prvulovic, D. <sup>2</sup>,  
Miljanovic, B. <sup>1</sup>, and Grubor-Lajsic, G. <sup>1</sup>

<sup>1</sup> Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovica 2, 21 000 Novi Sad, Serbia

<sup>2</sup> Department of Field and Vegetable Crops, Faculty of Agriculture, Trg Dositeja Obradovica 8, 21000, Novi Sad, Serbia

<sup>3</sup> ACIMSI, University of Novi Sad, 21000 Novi Sad, Serbia

Received 10 Dec. 2010;

Revised 12 May 2011;

Accepted 19 May 2011

**ABSTRACT:** This study focused on use of bacterial extracellular enzyme activities as biochemical parameters, along with the microbiological and physicochemical characteristics, in a comprehensive assessment of water quality of four lake ecosystems in the Province of Vojvodina (northern Serbia): Provala, Ludas, Zobnatica and Palic. Water samples were collected in June and October, 2008. For assessment of microbiological water quality, heterotrophic plate count, total and fecal coliform count, as well as total bacterial count was determined. Based on microbiological parameters, the water ecosystems were appropriately classified. The following extracellular enzyme activities were determined: alkaline phosphatase,  $\beta$ -D-glucosidase and acetate esterase using fluorogenic 4-methylumbelliferone labeled substrates. The results showed that Provala Lake had the best water quality taking in consideration all of the parameters, followed by Zobnatica, while Ludas and Palic had a significant level of organic water pollution both in June and October. The results indicate that further similar studies should include both microbiological and biochemical analyses, in order to obtain more relevant data on water quality.

**Key words:** Lake water, Alkaline phosphatase activity,  $\beta$ -D-glucosidase activity, Acetate esterase activity, aerobic heterotrophic bacteria, Total coliform count, Total bacterial count

### INTRODUCTION

The growth of heterotrophic bacteria is affected by organic matter. More than 95% of organic matter in subsurface aquatic habitats consists of high molecular weight compounds which cannot be directly exploited by bacteria. In order to provide low molecular weight organic compounds that could be uptaken into the cells, heterotrophic microorganisms produce extracellular enzymes (Münster and Chróst, 1990). Microbial enzymes are therefore, directly related to the concentrations and ratios of limiting nutrients, and may be sensitive indicators of organic pollution in aquatic ecosystems. According to many authors (Boetius, 1995, Jackson *et al.*, 1995, Mallet and Debroas, 1999), enzyme assays can provide powerful tools for studying organic matter presence and degradation in aquatic ecosystems. Activities of many hydrolytic extracellular enzymes can be measured with high sensitivity, using fluorogenic model substrates, which are available for the most of natural compounds (Hoppe, 1993). We studied the extracellular enzymatic activities, physicochemical and

microbiological parameters in surface water of the lake ecosystems Provala, Ludas, Zobnatica and Palic of the Province of Vojvodina (northern Serbia) in June and October, 2008 in order to assess water quality with special attention paid to determine applicability of both microbiological and biochemical parameters for water quality assessment.

### MATERIALS & METHODS

Water samples were taken from four lakes of the Province of Vojvodina (northern Serbia): Provala Lake, Ludas Lake (two sampling sites), Zobnatica and Palic lakes on June 1, and October 1, 2008. The location and coordinates of sampling sites are shown in Fig. 1. The geographic coordinates are: 45°24' N, 019°05' E for Provala, 46°06' N, 019°49' E for Ludas 1, 46°03' N, 019°49' E for Ludas 2, 46°05' N, 019°45' E for Palic and 45°50' N, 019°38' E for Zobnatica Lake. The North latitudes and East longitudes were the same in June and October. Palic and Ludas lakes are protected areas. Palic Lake is proclaimed as a protected area and a Park

\*Corresponding author E-mail: elvira.pamer@dbe.uns.ac.rs

## Archive of SID

of Nature, while Ludas Lake is a special natural preserve and, since 1977, designated as a swamp area of international significance by the Ramsar Convention. Lake water samples were collected at depth of 1.5 m. The samples were kept and transported to the laboratory in cooled containers and analysed within 24h. For determination of total bacterial count (TBC) water samples were fixed with formaldehyde (2% final concentration) after the sampling. The basic parameters of water quality oxygen, temperature, electrical conductivity and pH were determined using oxymeter and conductivity meter (Hanna HI-9142 and HI-933000, Hanna Instruments, USA), as well as pH meter (WTW InoLab-4, Germany). The total suspended solids (TSS), chemical oxygen demand (COD), biochemical oxygen demand (BOD), total organic carbon (TOC) and surfactant (SUR) were measured at sampling site after the sampling using multiparameter PASTEL-UV Portable Analyzer "SECOMAM" (Nova Analytics Corporation, USA). Total nitrogen (TN), nitrate ( $\text{NO}_3^-$ ), total and inorganic phosphorus ( $\text{P}_{\text{tot}}$ ,  $\text{PO}_4^{3-}$ ) were measured in laboratory within 24h. TN was determined according to the method of Kjeldahl in unfiltered samples (CNR-IRSA, 1994).  $\text{NO}_3^-$  was determined spectrophotometrically in filtered samples according to Cataldo (Cataldo *et al.*, 1975). Ammonia was determined by treating the lakewater samples in an alkaline citrate medium with sodium hypochlorite and phenol in the presence of sodium nitroprusside that acted as a catalyser (APHA, 1992). The blue indophenol colour formed with ammonia was measured spectrophotometrically at 640nm.  $\text{P}_{\text{tot}}$  was determined by antimony method in unfiltered samples (Genchi, 1990) as well as  $\text{PO}_4^{3-}$  in filtered samples, using the same method. Organic phosphorus ( $\text{P}_{\text{org}}$ ) was calculated by  $\text{P}_{\text{org}} = \text{P}_{\text{tot}} - \text{PO}_4^{3-}$ . For this analysis the spectrophotometer Jenway 6505 UV/Vis, UK was used. In all samples, counts of aerobic heterotrophic bacteria, fecal and total coliforms were determined on solid media using the spread plate method. The aerobic heterotrophic bacteria (HPC) were grown on Nutrient agar (Torlak, Serbia) using R2A medium (Merck, Germany). The media were incubated at 26°C for 5-7 days. Total and fecal coliforms were counted on Chromocult Coliform Agar (Merck, Germany) after 48 hours of incubation at 37°C. The waters were classified according to Kohl (1975) on the basis of HPC while coliforms were used for classifying water classification was according to Kavka *et al.* (2006). Bacterial counts in samples were determined by epifluorescence microscopy according to Raymond *et al.* (1994). Sample aliquots were filtered under vacuum onto Sudan Black prestained 0.2  $\mu\text{m}$  pore size Nuclepore polycarbonate membrane filter with cellulose-acetate supporter filter. TBC was enumerated by direct counting of acridine

orange stained cells (0.01% w:v, 5 min) using Olympus CX41 epifluorescent microscope with corresponding filter. The bacterial cells were counted on at least 10 fields. The cell count was averaged and the results were expressed per milliliter of water samples. Extracellular enzyme activities were determined in triplicate samples using fluorogenic model substrate 4-methylumbelliferone (MUF) according to Hoppe (Hoppe, 1983). Appropriate MUF-substrates were added to final concentrations: 500  $\mu\text{mol/L}$  MUF-glucose, 250  $\mu\text{mol/L}$  for MUF-phosphate and 100  $\mu\text{mol/L}$  MUF-acetat, according to determined enzyme activities:  $\beta$ -D-glucosidase, alkaline phosphatase and acetate esterase. The reaction mixtures were incubated for 8 hours in dark at room temperature. The reaction was interrupted by boiling for 5 min. The pH of mixtures was adjusted to 10.5 by adding 10:1 of ammonium glycine buffer (0.1 M) before fluorescence measurement. The fluorescence of 4-MUF, hydrolysed from the model substrates was measured, using 96-well microtiter plate and automated microtiter plate fluorometer (Fluoroscan Ascent FL, USA) with excitation and emission filters of 355 and 460 nm. Its quantity determined fluorometrically indicated the level of extracellular enzyme activity in the sample. Standard curve was constructed using an appropriate range of known concentrations. The enzyme activities were expressed as total activity ( $\text{nmol/h/dm}^3$ ) and as specific enzyme activities, calculated per bacterial cell ( $\text{amol/h/cell}$ ).

The correlation coefficients ( $r$ ) and the statistical significance of the relationships ( $p$ -values) were calculated using Statistica 9.1 (StatSoft, USA). The data were analysed using two-way analyses of variance (ANOVA) followed by Tukey's HSD test (Hinkle, 1994).

## RESULTS & DISCUSSION

Water samples were analysed for the physicochemical characteristics at each sampling site and the results are shown in Table 1. The total phosphorous ( $\text{P}_{\text{tot}}$ ) values in the water samples during 2008 varied between 0.00-2625.5  $\mu\text{g P dm}^{-3}$  while organic phosphorous varied between 0.00-1748.8  $\mu\text{g P/dm}^3$  (Table 1). In June, all water samples showed higher organic than inorganic phosphorus fraction, with the exception of Ludas 1 and Palic, where inorganic fraction was 3.57 and 2 fold higher than  $\text{P}_{\text{org}}$ , respectively. In October, higher inorganic than organic phosphorus level was detected in Palic (1.3 fold higher) and in Ludas 2 (1.5 fold higher). The results of microbiological parameters are presented in Table 3 and Fig. 4. Using R2A medium, bacterial count varied from 3.11±0.18 CFU/mL (Zobnatica, June) to 4.76±0.14 CFU/mL (Palic, October). Total coliforms were detected in the majority of the samples. In June, the highest count was detected

Archive of SID

1- Zobnatica

2- Provala

3- Ludas

4- Palic




Fig. 1. Location of the sampling sites of Provala, Ludas 1, Ludas 2, Zobnatica and Palic lakes in the Province of Vojvodina (northern Serbia)

Table 1. Main physicochemical parameters of water quality at sampling sites: Provala, Ludas 1, Ludas 2, Zobnatica and Palic in June and October, 2008

Parameter	Sampling site				
	Provala	Ludas 1	Ludas 2	Zobnatica	Palic
	June - October	June - October	June - October	June - October	June - October
NH <sub>4</sub> <sup>+</sup> [mg/dm <sup>3</sup> ]	0.3 - 0.1	1.8 - 1.2	1.1 - 0.7	0.9 - 0.5	2.0 - 1.1
NO <sub>3</sub> <sup>-</sup> [mg/dm <sup>3</sup> ]	0.7 - 1.2	1.7 - 3.7	1.1 - 1.8	1.0 - 3.3	1.8 - 3.9
NO <sub>2</sub> <sup>-</sup> [µg/dm <sup>3</sup> ]	30.0 - 17.2	107.2 - 80.1	87.2 - 62.9	25.7 - 31.5	58.6 - 115.8
DO [mg/dm <sup>3</sup> ]	4.8 - 3.3	6.1 - 6.4	0.2 - 3.1	3.9 - 3.4	1.88 - 4.5
O <sub>2</sub> [%]	62.3 - 57.8	76.0 - 91.7	2.8 - 46.5	47.5 - 52.2	24.3 - 71.5
Temp. [°C]	28.0 - 17.1	26.3 - 15.4	23.2 - 14.9	25.2 - 15.2	27.5 - 15.0
pH	9.2 - 8.9	9.5 - 9.8	8.1 - 8.5	8.6 - 9.1	8.3 - 9.6
EC <sub>w</sub> [µS/cm]	380.0 - 156.8	896.0 - 1078.0	1300.0 - 1368.0	1210.0 - 1190.0	1111.0 - 939.0
P <sub>tot</sub> [µg P/dm <sup>3</sup> ]	ND - 88.2	1175.0 - 148.4	270.8 - 43.0	32.1 - 32.3	2625.5 - 144.1
PO <sub>4</sub> <sup>3-</sup> [µg P/dm <sup>3</sup> ]	ND - 17.2	918.0 - 53.8	110.2/25.8	13.8/10.8	1748.8/81.7
P <sub>org</sub> [µg P/dm <sup>3</sup> ]	ND - 71.0	257.0 - 94.6	160.6 - 17.2	18.3 - 21.5	876.7 - 62.4
TOC [mg/dm <sup>3</sup> ]	5.1 - 3.3	16.2 - 15.6	-/-	8.4 - 7.3	13.9 - 10.0
TSS [mg/dm <sup>3</sup> ]	<2.5 - 5.7	54.5 - 29.4	-/-	<2.5 - <2.5	<2.5 - 15.4
SUR [mg/dm <sup>3</sup> ]	3.3 - 0.6	6.8 - 5.6	-/-	3.7 - 1.0	8.1 - 2.8
COD [mg/dm <sup>3</sup> ]	11.5 - 9.9	42.5 - 42.0	-/-	21.4 - 22.0	33.0 - 28.0
BOD [mg/dm <sup>3</sup> ]	7.0 - 4.8	22.8 - 22.2	-/-	11.9 - 10.8	17.0 - 14.4

All the values are expressed as mean, number of examinations n=3

## Archive of SID

in Ludas 2 ( $6.43 \pm 0.39$  CFU in 100 mL) and in October in Ludas 1 ( $4.48 \pm 0.48$  CFU in 100 mL). No fecal coliforms were detected in one milliliter of the samples. Total bacterial count in June was on the order of  $10^7$ . In October it decreased to the order of  $10^6$  (Table 3). The highest bacterial density in June was detected in Ludas 1 and Zobnatica ( $3.98 \times 10^7$  and  $3.91 \times 10^7$ , respectively), and the lowest in Palic and Provala ( $1.32 \times 10^7$  and  $1.41 \times 10^7$ , respectively). In October, the bacterial density showed variation from  $1.81 \times 10^6$  (Ludas 1) to  $5.33 \times 10^6$  (Ludas 2).


Alkaline phosphatase activity in June significantly differed depending on the locality, with the highest activity in Ludas 2 (Fig. 2A, Table 2). In relation to the season, in localities Provala and Zobnatica a significant increase in phosphatase activity was recorded in October, while in Palic and Ludas 2 there was a decrease. The results of phosphatase activity in Ludas 1 are not shown due to the lack of samples. The results of specific phosphatase activity also varied depending on the locality and season (Fig. 2B, Table 2). Thus, the specific phosphatase activity was significantly lower in Provala than in Zobnatica, while their activities were lower in comparison with Palic and Ludas 2 (no significant difference was observed). In comparison with June, a significant increase of specific phosphatase activities was detected at all of the sites in October. The highest  $\beta$ -D-glucosidase activity in June was recorded in Ludas 1, being significantly higher than in Ludas 2. Both of these localities showed significantly higher levels in relation to Palic and Provala, whose activities were approximate (Fig. 4A, Table 2). The results for glucosidase activity in Zobnatica are not shown due to the lack of samples. In October, compared to June, there was a significant increase of  $\beta$ -D-glucosidase activity in Provala and Palic and a decrease in Ludas 1 and Ludas 2. The specific  $\beta$ -D-glucosidase activity significantly increased in October at all of the sites (Fig. 4B, Table 2). Acetat esterase activity was different at all sites in June, with the highest activity recorded in Provala (Fig. 3A, Table 2). In terms of specific acetate esterase activity, the highest levels were detected in Provala, while significantly lower ones were recorded in Palic. Ludas 1, 2 and Zobnatica showed similar activities, which proved to be significantly lower in comparison with Palic (Fig. 3B, Table 2). In October the total and specific esterase activity increased significantly at all sites (and between them), the exception being Provala for total activity, which appeared to be significantly lower (Fig. 3A and B, Table 2).

The composition and availability of organic matters, which are mainly derived from allochthonous rather than from autochthonous matters (Findlay, 1991), are the major factors which influence the development and

activity of the microbial heterotrophic communities. Heterotrophic bacteria are the key trophic level in natural organic matter decomposition, nutrient cycling, and structure of the food web (Chróst and Siuda, 2002, Münster, 1999). Therefore the studies of the microbial mobilization, transformation and utilization of a variety of natural organic matters are one of the most important areas of the aquatic microbial ecophysiology (Chróst, 1991).

The results of microbiological analyses indicated that all of the examined ecosystems showed a significant variation of water quality. Generally, better water quality was detected during October in all of the examined lakes. According to HPC, in June water of the four lakes belonged to I-II and II-III class, while in October the water was classified from I to II-III class. Taking into consideration total coliform counts, the water quality varied from II to V class. The detected total coliforms, which naturally occur and multiply in waters (De Zuane, 1997), without fecal coliforms, indicate the absence of water fecal pollution. It is interesting to notice a generally higher count of bacteria on R2A medium than on Nutrient agar reported in certain studies of various waters types (Massa *et al.*, 1998), as the R2A medium is a low nutritive and more suitable for autochthonous water bacteria. However, our results showed that the bacterial counts obtained on Nutrient agar and R2A medium did not statistically differ, neither in June nor in October ( $p = 0.601$  and  $p = 0.08$ , respectively). Such results may indicate that most of the bacteria detected on Nutrient agar are autochthonous, and that they dominate in the ecosystems. Generally, the water of Provala had the best quality, followed by Zobnatica and Palic, while Ludas 1 and 2 had the worst quality, considering all microbiological parameters. Physicochemical parameters of the lake water samples also indicate generally better water quality in October (Table 1). Considering the examined ecosystems, Palic and especially Ludas showed worse water quality than Zobnatica and Provala lakes. Similar findings for physicochemical parameters of the same localities were reported earlier by other authors (Simeunovic *et al.*, 2010).

Our results of estimated extracellular enzymatic activities showed that the hydrolysis activity increased in June as well as in October in the following order:  $\beta$ -D-glucosidase < alkaline phosphatase < acetat esterase, which complies with the results of Mudryk and Skórczewski (Mudryk and Skórczewski, 2004) regarding lake water ecosystems. According to Münster *et al.* (Münster and de Haan, 1998), among the most important microbial extracellular enzymes in surface waters are glucosidases and phosphatases.



**Fig. 2. Alkaline phosphatase activity (A) and specific alkaline phosphatase activity (B) in localities Provala, Ludas 1, Ludas 2, Zobnatica and Palic in June and October, 2008. Differences between groups were tested by two-way analysis of variance (two-way ANOVA) with Tukey's HSD post hoc test, with locality (L) and season (S) as factors with locality (L) and season (S) as factors**


the former participating in depolymerisation of polysaccharides, and the latter in dephosphorylation of organic molecules. Esterases, on the other hand, can be used for the determination of general microbial extracellular enzyme activities due to their varying substrate specificity and biological functions.

In general, the lowest  $\beta$ -D-glucosidase activities in both seasons might be attributed to reduced availability of polymeric carbohydrates (cellulose and cellobiose) as a source of energy in these lake ecosystems. In early summer (June), a significantly

higher level of  $\beta$ -D-glucosidase activity ( $p < 0.001$ ) in Ludas Lake (localities Ludas 1 and 2) in relation to Palic and Provala was detected. According to Chróst and Overbeck (Chróst and Overbeck, 1990) it could indicate on phytoplankton bloom breakdown and water blooms in the lakes examined in the present study are documented during early summer (Simeunovic, 2010). A significant decrease in alkaline phosphatase activity recorded in localities Palic and Ludas 2 and increase in localities Provala and Zobnatica recorded in October is in accordance with the content of organic

**Table 2. Results of statistical analyses two-way ANOVA with Tukey's HSD post hoc test, with locality (L) and season (S) as factors, for enzyme and specific enzyme activity of alkaline phosphatase,  $\beta$ -D-glucosidase and acetate esterase**


Enzyme activity		Locality (L)	Season (S)	L x S
Alkaline phosphatase	df	3	1	3
	MS	$1.65 \times 10^6$	$4.36 \times 10^6$	$4.3 \times 10^6$
	F	<b>453 ***</b>	<b>1200***</b>	<b>1178 ***</b>
$\beta$ -D-glucosidase	df	3	1	3
	MS	5665	1870	7824
	F	<b>111 ***</b>	<b>36.7 ***</b>	<b>153 ***</b>
Acetate esterase	df	4	1	4
	MS	$261 \times 10^3$	$7.2 \times 10^6$	$1.4 \times 10^6$
	F	<b>10.5 ***</b>	<b>292 ***</b>	<b>57.9 ***</b>
<b>Specific enzyme activity</b>				
Alkaline phosphatase	df	3	1	3
	MS	$235 \times 10^3$	$1.9 \times 10^6$	$320 \times 10^3$
	F	<b>701 ***</b>	<b>5714 ***</b>	<b>955 ***</b>
$\beta$ -D-glucosidase	df	3	1	3
	MS	1053	7135	1010
	F	<b>211 ***</b>	<b>1429 ***</b>	<b>202 ***</b>
Acetate esterase	df	4	1	4
	MS	$496 \times 10^3$	$5.5 \times 10^6$	$595 \times 10^3$
	F	<b>845 ***</b>	<b>9346 ***</b>	<b>1014 ***</b>



**Fig. 3. Acetate esterase (A) and specific acetate esterase activity (B) in localities Provala, Ludas 1, Ludas 2, Zobnatica and Palic in June and October, 2008. Differences between groups were tested by two-way analysis of variance (two-way ANOVA) with Tukey's HSD post hoc test, with locality (L) and season (S) as factors with locality (L) and season (S) as factors**

**Table 3. Total bacterial counts, total coliform counts and aerobic heterotrophic bacteria on R2A medium**

Parameter	Samples	Samples				
		Provala	Ludas 1	Ludas 2	Zobnatoca	Palic
Total bacterial count (x10 <sup>6</sup> /mL)	June	14.10±1.90	40.0±2.90	33.90±3.50	40.00±2.50	13.20±1.20
	October	4.45±0.63	3.20±0.63	1.81±0.58	5.33±0.65	3.40±0.32
Heterotrophic bacteria on R2A (CFU/mL)	June	3.38±0.37	4.53±0.32	4.55±0.36	3.11±0.18	4.33±0.20
	October	3.96±0.82	4.62±0.22	4.17±0.13	4.32±0.78	4.76±0.14
Total coliforms (CFU/100 mL) (water class)	June	3.00±0.15	4.48±0.86	6.43±0.39	3.91±2.5	4.39±0.39
	October	N.D.*	4.48±0.48	4.00±0.00	4.00±0.00	N.D.
		-	III	II	II	-



**Fig. 4.  $\beta$ -D-glucosidase activity (A) and specific  $\beta$ -D-glucosidase activity (B) in localities Provala, Ludas 1, Ludas 2, Zobnatoca and Palic in June and October, 2008. Differences between groups were tested by two-way analysis of variance (two-way ANOVA) with Tukey's HSD post hoc test, with locality (L) and season (S) as factors with locality (L) and season (S) as factors**




Fig. 5. Water categorization based on aerobic heterotrophic bacteria

phosphorus fraction in examined lake ecosystems (Fig. 2A, Table 1). The highest hydrolysis activity of acetate esterase from measured extracellular enzyme activities (Fig. 2, 3 and 4A) might be in correlation with the presence of broad substrate spectrum of acetate esterase in the lake ecosystems. The reason for significantly higher acetate esterase activities in October than in June at all localities, except for Provala ( $p < 0.001$ ) (Fig. 3A), could be the presence of high level of surfactants in October (Table 1). SUR matters could affect microbial growth by increasing bioavailability of hydrophobic compounds and accelerating their degradation (Bardi *et al.*, 2000). The highest hydrolytic activity of acetate esterase in Palic and Ludas 1 in October was in correlation with TOC levels in water ecosystems. Significantly higher specific enzyme activities ( $p < 0.001$ ) in October than in June (Table 2), among the other factors, could be related to the effect of temperature on enzyme synthesis and activity. This complies with the well known fact that low water temperature could induce greater enzyme production by certain bacteria (Quist and Stokes, 1969).

## CONCLUSION

The correlation between aerobic heterotrophic bacteria, aerobic heterotrophic bacteria on R2A medium and total coliforms, on one hand, and enzyme activities

of waters of the examined lakes on the other hand, did not prove to be statistically significant. For instance, linear correlation between  $\beta$ -glucosidase activity and bacterial count was established during filtration process at various sites in a groundwater recharge plant at the Ruhr River (Hendel *et al.*, 2001), but not in Danube water (Kavka *et al.*, 1996). The absence of the correlation in the present study could be the result of various factors (physical conditions, nutrient availability, presence of various compounds such as humic acids, biocides and toxins etc.) that influence on heterotrophic bacteria count, enzymatic production and/or activity, and thus on ecophysiology of the ecosystems (Boavida and Wetzel, 1998, Barcina *et al.*, 1997). In addition, some heterotrophic bacteria do not produce phosphatase, lipase and/or glycosidase, while other organisms, including metabolically active but not cultivable microorganisms can produce these enzymes (Barcina *et al.*, 1997). The complexity of the relationships among microbiological and bacteriological parameters indicates the necessity of performing both microbiological and biochemical analyses in order to obtain more relevant data on water quality (Fig. 5). Such analysis provides information on sanitary quality of water in one hand, and gives insight into the processes that take part in examined ecosystems. Taking in consideration all examined



## Archive of SID

parameters, it is obvious that among four lakes of the Province of Vojvodina, Provala and Zobnatica lakes had the best water quality, while Ludas and Palic were overloaded with organic matter both in June and October, 2008.

### ACKNOWLEDGEMENT

This work was supported by the grant from the Ministry of Science and Technological Development of the Republic of Serbia.

### REFERENCES

- APHA, (1992). American Public Health Association, Standard Methods for the Examination of Water and Wastewater. 18th ed., Washington, DC, pp. 42–340.
- Barcina, I., Lebaron, P. and Vives-Rego, J. (1997). Survival of allochthonous bacteria in aquatic systems: a biological approach. *FEMS Microbiology Ecology*, **23**, 1-9.
- Bardi, L., Mattei, A., Steffan, S. and Marzona, M. (2000). Hydrocarbon degradation by a soil microbial population with  $\beta$ -cyclodextrin as surfactant to enhance bioavailability. *Enzyme and Microbial Technology*, **27**, 709–713.
- Boavida, M.-J. and Wetzel, R. G. (1998). Inhibition of phosphatase activity by dissolved humic substances and hydrolytic reactivation by natural ultraviolet light. *Freshwater Biology*, **40**, 285-293.
- Boetius, A. (1995). Microbial hydrolytic enzyme activities in deep-sea sediments. *Helgoländ Meeresuntr.* Springer Berlin / Heidelberg, **49**, 177–187.
- Cataldo, D. A., Haroon, M., Schrader, L. E. and Youngs, V. L. (1975). Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Commun. Soil Sci. Plant Anal.*, **6**, 71-80.
- Chróst, R. J. and Overbeck, J. (1990). Substrate-ectoenzyme interaction: significance of  $\beta$ -glucosidase activity for glucose metabolism by aquatic bacteria, *Arch. Hydrobiol. Beih.*, **34**, 93-98.
- Chróst, R. J. (1991). Ectoenzyme in aquatic environments: Microbial strategy for substrate supply, *Verh. Int. Verein. Limnol.*, **24**, 2597-2600.
- Chróst, R.J. and Siuda, W. (2002). Ecology of microbial enzymes in lake ecosystems, (In: G.R. Burns and P.R. Dick (Eds.), *Enzymes in the environment: Activity, ecology, and applications*, Merceel Dekker, New York, pp. 35-72).
- CNR-IRSA, (1994). *Metodi analitici per le acque*, Istituto Poligrafico e Zecca dello Stato, Rome, Quaderno 100, **2**, 342, (in Italian).
- De Zuane, J. (1997). *Handbook of Drinking Water Quality*, 2nd ed., Van Nostrand Reinhold, New York.
- Findlay, S., Pace, L. M., Lints, D., Cole, J. J., Caraco, F. N. and Peierls, B. (1991). Weak coupling of bacterial and algal production in a heterotrophic ecosystem: the Hudson River Estuary, *Limnol. Oceanogr.*, **36**, 268-278.
- Genchi, G. (1990). Azoto e fosfori totali, *Nova Thalassia* **11**, pp. 171-179, (in Italian).
- Hinkle, E.D., Wiersma W. and Jurs, G.S. (1994). *Applied Statistics for Behavioral Sciences*, 3rd ed. Houghton Mifflin, Boston, pp. 706.
- Hoppe, H. G. (1983). Significance of exoenzymatic activities in the ecology of brackish water: measurements by means of methylumbelliferyl-substrates, *Mar. Biol. Prog. Ser.* **11**, pp. 299-308.
- Hoppe, H. G. (1993). Use of fluorogenic model substrates for extracellular enzyme activity (EEA) measurement of bacteria, (In: F.P. Kemp, F.B. Sherr, B.E. Sherr, and J.J. Cole (Eds.), *Handbook of Methods in Aquatic Microbial Ecology*, Lewis Publishers, London, pp. 423–431).
- Jackson, C.R., Foreman, C.M. and Sinsabaugh, R.L. (1995). Microbial enzyme activities as indicators of organic matter processing rates in a Lake Erie coastal wetland, *Freshwater Biology*, **34**, 329–342.
- Kavka, G. G., Kasimir, G. D. and Farnleitner, A. H. (2006). Microbiological water quality of the River Danube (km 2581-km 15): Longitudinal variation of pollution as determined by standard parameters, (In: Proceedings 36th International Conference of IAD, Austrian Committee Danube Research / IAD, Vienna, pp. 415-421).
- Kohl, W. (1975). *Über die Bedeutung Bakteriologischer Untersuchungen für die Beispiel der Osterreich Donau*, *Arch. Hydrobiol.* **44**, pp. 392-461, (in German).
- Mallet, C. and Debroa, D. (1999). Relations between organic matter and bacterial proteolytic activity in sediment surface layers of a eutrophic lake (Lake Aydat, Puy de Dôme, France), *Arch. Hydrobiol.* **145**, pp. 39–56.
- Massa, S., Caruso, M., Trovatielli, F. and Tosques, M. (1998). Comparison of plate count agar and R2A medium for enumeration of heterotrophic bacteria in natural mineral water. *World Journal of Microbiology and Biotechnology*, **14**, 727-730.
- Mudryk, Z. J. and Skórczewski, P. (2004). Extracellular enzyme activity at the air-water interface of an estuarine lake, *Estuarine. Coastal and Shelf Science*, **59**, 59–67 (in Russian).
- Münster, U. and Chróst, R. J., (1990). Origin, composition and microbial utilization of dissolved organic matter, (In: J. Overbeck, and J.R. Chróst (Eds.), *Aquatic microbial ecology: Biochemical and molecular approaches*. Springer-Verlag, New York, pp. 8-46).
- Münster, U. and de Haan, H. (1998). The role of microbial extracellular enzymes in the transformation of dissolved organic matter in humic waters, (In: O.D. Hessen and J.L. Tranvik (Eds.), *Aquatic Humic Substances: Ecology and Biogeochemistry*, Springer-Verlag, Berlin, Heidelberg, Germany, pp. 199–257).
- Münster, U. (1999). Bioavailability of nutrients, (In: J. Keskitalo and P. Eloranta (Eds.), *Limnology of Humic Waters*, Backhuys Publishers, Leiden, pp. 77–94).

*Archive of SID*

Quist, R. G. and Stokes, J. L. (1969). Temperature range for formic hydrogenlyase induction and activity in psychrophilic and mesophilic bacteria, *Antonie van Leeuwenhoek, J. Microbiol. Serol.*, **35**, 1-8.

Raymond, L., Kepner, J. L. and Pratt, R. J. (1994). Use of fluorochromes for direct enumeration of total bacteria in environmental samples: past and present, *Microbial. Rev.*, **58**, 603–615.

Simeunovic, J., Svircev, Z., Karaman, M., Knezevic, P. and Melar, M. (2010). Cyanobacterial blooms and first observation of microcystin occurrence in freshwater ecosystems in Vojvodina (Serbia). *Fresenius Environmental Biology*, **19**, 198-207.