Int. J. Environ. Res., 6(1):209-218, Winter 2012 ISSN: 1735-6865

Bioaccumulation of Heavy Metals in Fish Species Collected From Former Tin Mining Catchment

Ashraf, M. A.^{1*}, Maah, M. J.¹ and Yusoff, I.²

¹Department of Chemistry, University of Malaya, Kuala Lumpur 50603 Malaysia

²Department of Geology, University of Malaya, Kuala Lumpur 50603 Malaysia

Received 4 June 2011; Re	evised 4 Aug. 2011;	Accepted 14 Aug. 2011
--------------------------	---------------------	-----------------------

ABSTRACT: This study has been carried out to determine the accumulation of heavy metals (arsenic, copper, lead, tin and zinc) in fish samples collected from former tin mining catchment. Total eight types of fish family having fifteen different species are identified. Cyperinidae is the most abundant family with eight different species found in the catchment. Fish samples were analysed by using inductively coupled plasma optical emission spectrophotometer ICP-OES. The accuracy of method is validated by certified reference material. The highest level of arsenic was observed in Hampala macrolepidota while Osteochilus hasselti shows the least. For Copper concentrations, the maximum was obtained in Mastacembelus armatus and the minimum in Oxyeleotris marmorata. High levels of lead, tin and zinc were found in the Rasbora elegans, Trichogaster trichopterus, Oxyeleotris marmorata respectively while Macrobrachium resenbergii, Mastacembelus armatus, Rasbora elegans had the least concentrations. The sequence of order of the heavy metals measured was $S_{n} > 0$ Pb > Zn > Cu > As, respectively. The results showed elevated levels of tin, led and zinc in all the fish samples although copper and arsenic were available in relatively low concentration in the most samples. Sn, Pb, and Zn concentration in the samples were greater than Malaysia food act permissible levels. Hazard index < 1 suggests unlikely adverse health effects whereas HI > 1 suggests the probability of adverse health effects. Although the heavy metals analysed in the catchment did not pose any immediate health risk to humans but due to the bioaccumulation and magnification of these heavy metals in humans, it is essential to safeguard levels of the metals in the environment.

Key words: Species, Classification, Heavy metals, Permissible level, Health effects, Hazard index

INTRODUCTION

During recent years the concentrations of toxic metals in many ecosystems are reaching unprecedented levels. The increasing use of metals in industry and mining activities have lead to serious environmental pollution through effluents and emanations (Goldberg *et al.*, 1978; Phillips, 1980; Sericano *et al.*, 1995). Under certain environmental conditions, heavy metals may accumulate to a toxic concentration (Güven *et al.*, 1999), and cause ecological damage (Harms, 1975; Jefferies & Freestone, 1984; Freedman, 1989).

Specifically aquatic systems are more sensitive to heavy metal pollutants and the gradual increase in the levels of such metals in aquatic environment, mainly due to anthropogenic sources, became a problem of primary concern (Meybeck *et al.*, 1989; Allen *et al.*, 1993). This is due to their persistence as they are not usually eliminated either by biodegradation or by chemical means, in contrast to most organic pollutants. <u>Moreover, the decay of organic materials in aquatic</u> systems together with detritus formed by natural weathering processes and uncontrolled mining activities provides a rich source of nutrients in both the bottom sediments and overlying water body (wade *et al.*, 2008; Jafari & Gunale, 2006). Microorganisms, microflora and algae are capable of incorporating and accumulating metal species into their living cells from various supply sources (Jaffar *et al.*, 1988).

Consequently, small fish become enriched with the accumulated substances. Predatory fish again, general display higher levels than their prey. Eventually man, consuming the fish, inevitably suffers from the results of an enrichment taken place at each trophic level, where less is extracted than ingested (Forstner & Wittmann, 1983). Fish occupies the highest tropic level in aquatic system (APHA, 1981). Besides that, it has high economical values, thus fish is suitable for water quality symbol and easy to be interpreted by public. Fish can response to

^{*}Corresponding author E-mail: chemaqeel@gmail.com

environmental changes that can be used for pollution indicator study. Fish is a good bio-indicator because it is easy to be obtained in large quantity, potential to accumulate metals, long lifespan, optimum size for analysis and easy to be sampled (Batvari *et al.*, 2007). Heavy metal intakes by fish in polluted aquatic environment are different depends on ecological requirements, metabolisms and other factors such as salinity, water pollution level, food and sediment. Fish accumulates metals in its tissues through absorption and human can be exposed to metals via food web. This will cause acute and chronic effect to human (Yilmaz & Dogan, 2007; Fidan *et al.*, 2007). The use of fish as bioindicator can determine the actual situation of pollution level before and during monitoring.

In general, studies on heavy metals by fish analysis can be important in two main aspects. First, from the public health point of view, where the attention has been drawn to the necessity of measuring the accumulation of heavy metals; particularly these metals which pose serious health hazards to humans (e.g. As, Pb, Hg). Second, from the aquatic environment view point, the main problem has been to prevent biological deterioration and to identify the sources which threaten ecological equilibrium. In this regard, the more abundant metals such as copper, zinc and manganese may sometimes represent greater hazard than lead, mercury and cadmium (Kinne, 1984).

Several techniques have been used for determination of metal concentrations in fish species such as flame atomic absorption spectrometry (Abedin, 1986; El-Mehdi, 1987; Ahmet, 1992; Petisleam, *et al.*, 2003; Petisleam, *et al.*, 2007; Bermejo-Barrera, *et al.*, 2001) graphite furnace atomic absorption spectrometry (Sperling, 1988; Mustafa, 2003; Botson *et al.*, 2004), electro-thermal atomic absorption spectrometry (Perez, *et al.*, 2001; Mendez, *et al.*, 2002), inductive coupled plasma (Chirila, *et al.*, 1999; Sanchez, *et al.*, 2003; Milne, *et al.*, 2010) and mass spectrometry (Sanchez, *et al.*, 2003; Petisleam, *et al.*, 2005). Different digestion methods were used as sample preparation methods for determination of heavy metals in fish samples (Olaifa *et al.*, 2004; Huijuan *et al.*, 2005).

The present study was undertaken to study the concentrations of Sn, Cu, Pb, Zn and As in different tissues (bone, skin, flesh and tail) of fish species at former tin mining catchment Bestari Jaya using ICP-MS. concentration levels of selected trace metals in commercially important fish species and correlate the concentration of metals with respect to their weight and length.

Bestari Jaya catchment is located at 3⁰, 24' 40.41" N and 101^o 24' 56.23" E. It is a part of Kuala Selangor district, located in Selangor, biggest state of the country. District Kuala Selangor has three main towns namely, Mukim Batang Berjuntai, Mukim Ulu Tinggi, Mukim Tg.karang. Bestari Jaya is located in Mukim Batang Berjuntai. Tin mining activities has ceased from last ten years, now sand mining. The catchment has total of 442 small and big mining lakes and ponds. Bestari Jaya has a tropical, humid climate, with very little variations in temperature throughout the year. The



Fig. 1. Map of Bestari Jaya Catchment

average temperature of the area is 32°C during day and 23 °C at night. (Ashraf *et al.*, 2010).

MATERIALS & METHODS

Fish samples were collected from Bestari Jaya catchment with the help of local fishermen between March and April of 2011. After collection, the samples were cleaned with deionized-distilled water, weighed, measured, stored in pre-cleaned plastic bags, and kept frozen in an ice box. Fifteen (15) fish samples were double bagged in separate new plastic bags, sealed and labelled accordingly.

The physical characteristics of the sampled fish species were assessed. A measuring board made of perspex and a fixed meter scale was used to measure the total length (cm) of each individual samples. At the zero end of the scale, a suitable stop is placed at 90 angle. The anterior end of the fish is placed against this stop. The total length is recorded for comparison to the known maximum size of the species. An electronic weight scale is used to weigh the individual mass of the fish. In addition, the total number and weight (g) for each batch and species was also determined to represent the overall abundance in the specific river. The weight of fish, together with its size distribution would indicate the fish life stages. The weight is measured in gram (g).

The ratio between the fish whole body weight to body length (head to tail) was examined to identify fishes with anomalously high or low ratios. The overall health of fish species in the catchment were assessed using the method developed by (Williams, 2000). A condition factor index (K) was calculated using the equation:

$K = 100W/L^{3}$

where, W = body weight in grams; L = body length (fork length) in cm

The frozen fish samples were partially thawed on cleaned plastic sheets using scalpels with steel blades and plastic forceps. Whole taxa of a designated length range were cut into small pieces with Teflon-tipped scissors. The samples were pooled and freeze-dried for 10 days to a constant weigh for the determination of metal content, 0.5 g dry muscle samples of the fish. (Moeller *et al.*, 2001).

Two types of digestion procedures were applied. Optimum digestion conditions are given below. The samples were dried to constant weight at 110 °C.

For dry ashing analysis, sample (1 g) was placed in a high form porcelain crucible. The furnace temperature was slowly increased from room temperature to $450 \,^{\circ}$ C

in 1 h. The samples were ashed for about 4 h until a white or grey ash residue was obtained. The residue was dissolved in 5 ml of HNO_3 (25% v/v) and the mixture, where necessary, was heated slowly to dissolve the residue. The solution was transferred to a 25 ml volumetric flask and made up to volume (Vaidya & Rantala, 1996). A blank digest was carried out in the same way. All metals were determined against aqueous standards.

For wet ashing analysis, the samples were solubilized using high-pressure decomposition vessels, commonly known as a digestion bomb. A sample (1 g) was placed into Teflon container and 5 ml of concentrated HNO3 was added. The system was heated to 130 °C for 90 min and finally diluted to 25 ml with deionized water. The sample solution was clear. A blank digest was carried out in the same way. All metals were determined against aqueous standards.

Inductively coupled plasma optical emission spectrometry ICP-OES (Perkin Elmer AA Analyst) was used to analyse the concentration of heavy metals in fish samples. All reagents used were of analytical reagent grade (Merck, Germany). Acid washed glassware analytical grade reagents and double distilled deionised water was used throughout the experiments. Standard stock solutions of arsenic, copper, tin, lead, and zinc were prepared from Titrasol (1000 mg/L). The working solution was freshly prepared by diluting an appropriate aliquot of the stock solutions. In order to check on the purity of the chemical used, a number of chemical blanks were run. There was no evidence of any contamination in these blanks. In order to validate the method for accuracy and precision, certified reference materials (fish: DORM-2, National Research Council, Canada) were analysed for each element. Accurate results are obtained as shown in (Table 2). The detection limit is defined as the concentration corresponding to 3 times the standard deviation of 10 blanks.

RESULTS & DISCUSSION

During sampling March-April 2011, the total number of fishes caught was 66 with 4064 grams. The main family observed in this survey, with 81 % (Fig. 2) of overall caught species was still the family of Cyprinidae, followed by the families of Palamonidae and Nandidae. The cyprinids were mainly represented by Terbul(68), Kawan(42), Rong(16) and Seluang dua titik (16). The condition factor is an estimation of the general well being of fish and crustaceans (Jones *et al.*, 1999). It is based on the hypothesis that heavier individuals of a given length are in better condition than less weightier ones (Bagenal & Tesch, 1978). Condition factor has been used as an index of growth and feeding intensity. Most notably, fish measured with an exceptionally low weight-to-length ratio may indicate the organism is stressed, either by physical injury, disease, dietary limitations, or poor water quality. The sampled fish species recorded an average K factor of 1.7 (Table 1). Age factor or maturity of fish may influence the accumulation of heavy metals (Mohsin & Ambak, 1991). Growth rate is important to stabilize the accumulation of metals. Fish can accumulates high level of pollutants if it has a constant growth and live in polluted ecosystem. This study indicates that mature fish accumulated higher metals compared to juvenile and premature fish. Labiobarbus cuvieri, Chela anommalura and Osteochilus hasseltii (mature omnivour fish) accumulated higher concentrations of Zn rather than Osteochilus vittatus and Channa striatus (premature). Although, comparison between same species is more appropriate, however, the comparison made was based on similar feeding behaviour. This study was unable to obtain sufficient number of fish to make comparison between age for the same species.

Fishes collected from catchment are comprises of three different types of feeding behaviors; herbivor, omnivor and carnivor. Hampala macrolepidota, Rasbora sumatrana, Rasbora elegans, Cyclocheilichthys apogon are carnivores (consume small fishes); Osteochilus vittatus, Trichogaster trichopterus and Pristoplepis fasciatus, Acanthopsis choirorhyhchos, Oxyeleotris marmorata are herbivores (consume plants) while Labiobarbus cuvieri, Chela anommalura, Osteochilus hasseltii, Mastacembelus armatus, Macrobrachium resenbergii and, Channa striatus are omnivores (Mohsin & Ambak,

1991). Feeding behaviour is one of the decisive factors of heavy metals accumulation in fish. Although, many studies showed carnivores accumulate higher metal concentrations (Mohsin & Ambak, 1991; Amundsen et al., 1997) result of this study demonstrates that metal concentrations in three different feeding behavior are not significantly differ. Low metal concentrations in the ecosystem could limits the absorption of these metals in the organisms. As regards to (Mohsin & Ambak, 1991) classification, this study found that most of the fish samples were considered as an adult. The mean metal concentrations of Cu, Zn, Pb Sn and As (mg/kg) in 15 fish species are shown in (Fig. 3). Tin mean concentrations were the highest as compared to other metals and ranged from 56.34 to 153.45 mg/kg. Metals accumulation in fish was found in the order of Sn>Pb>As>Zn>Cu. Metals concentration in fish tissues were detected at small variations.

Most of the locations chosen for the fish sampling for the determination of heavy metals have abandoned mining activities for more than 10 years and have been converted to secondary land uses such as fish breeding and other aqua-culture activities. The mean elemental concentrations measured in the fishes during the study period are also presented in (Fig. 3). Reported values of the various elements are the results of the total concentrations (dry weight) measured in each individual whole fish. Almost all the fish samples collected from the catchment contained detectable amounts of the elements studied (Cu, Pb, Zn, As, Sn). These elements were present in all the fish samples and at varying concentrations. It must be noted that, varying concentrations of the heavy metals were measured in the sampled fishes with some fishes



Fig. 2. Abundance of fish species in the catchment

÷
R.
ă
PL
<u>ਹ</u>
a
<u> </u>
he
Ŧ
.Е
Ś
ii.
ĕ
ġ.
ĩ
S
Ę
of
Ē
<u>ē</u> .
Ę
3
Ë
S
ā
J
<u>е</u> .
E
5
g
X
E
j.
~
č
Ë
÷
ē
5
- E
a
-R
Ĩ
3
Si.
Ň
-
5
le
ab
Ë

S No.	Fanily	Species	Local Name	Length	*Maturity Size/Adult (cm)	Condition of Fish (K)	Status
1	Cyperinidae	Osteochilus hasselti	Terbul	14.5-17.0	6-20	1.8	Mature
2		Osteochilus vittatus	Rong	8-22.3	50.0	1.7	Mature
б		Chela anommalura	Lalang	7-15.8	7.0-20.0	1.5	Premature
4		Labiobarbus cuvieri	Kawan	9.4-18.2	10.0-20.0	1.7	Mature
S		Hampala macrolepidota	Sebarau	7.5-18.5	25.0	1.9	Mature
9		Cyclocheilichthys apogon	Temperas mata merah	18.5-19.7	28.0-100.0	1.5	Premature
Ζ		Rasbora sumatrana	Seluang sumatra	8.8-10.7	48.0	1.7	Mature
8		Rasbora elegans	Seluang dua titik	14.4-17.4	12.0-18.0	1.8	Mature
6	Nandidae	Pristoplepis fasciatus	Patung	7.7-14.3	33.0	1.8	Mature
10	Cobitidae	Acanthopsis choirorhyhchos	Lali	7.5-12.5	20.0-35.0	1.6	Premature
11	Eleotridae	Oxyeleotris marmonata	Ketutu	13.6-17.6	10.0-12.0	2.0	Mature
12	mastacembelidae	Mastac embelus armatus	Tilan	30-41.2	12.0-19.0	2.0	Mature
13	Channidae	Channa striatus	Haruan	10.8-25	12.0-30.0	1.6	Premature
14	Palaemonida e	Macrobrachium resenbergii	Udang galah	4.5-13.5	10.0-30.0	1.8	Mature
15	Siluridae	Trichogaster trichopterus	Sepat padi	5.6-14.7	15.0-35.0	1.7	Mature
S No.	Family	Species	Local Name	Length	*Maturity Size/Adult (cm)	Condition of Fish (K)	Status
1	Cyperinidae	Osteochilus hasselti	Terbul	14.5-17.0	6-20	1.8	Mature
2		Osteochilus vittatus	Rong	8-22.3	50.0	1.7	Mature
Э		Chela anomnalura	Lalang	7-15.8	7.0-20.0	1.5	Premature
4		Labiobarbus cuvieri	Kawan	9.4-18.2	10.0-20.0	1.7	Mature
S		Hampala macrolepidota	Sebarau	7.5-18.5	25.0	1.9	Mature
9		Cyclocheilichthys apogon	Temperas mata merah	18.5-19.7	28.0-100.0	1.5	Premature
7		Rasbora sumatrana	Seluang sumatra	8.8-10.7	48.0	1.7	Mature
8		Rasbora elegans	Seluang dua titik	14.4-17.4	12.0-18.0	1.8	Mature
6	Nandidae	Pristoplepis fasciatus	Patung	7.7-14.3	33.0	1.8	Mature
10	Cobitidae	Acanthopsis choirorhyhchos	Lali	7.5-12.5	20.0-35.0	1.6	Premature
11	Eleotridae	Oxyeleotris marmorata	Ketutu	13.6-17.6	10.0-12.0	2.0	Mature
12	mastacembelidae	Mastac embelus armatus	Tilan	30-41.2	12.0-19.0	2.0	Mature
13	Channidae	Channa striatus	Haruan	10.8-25	12.0-30.0	1.6	Premature
14	Palaemonidae	Macrobrachium resenbergii	Udang galah	4.5-13.5	10.0-30.0	1.8	Mature
15	Siluridae	Trichogaster trichopterus	Sepat padi	5.6-14.7	15.0-35.0	1.7	Mature

Int. J. Environ. Res., 6(1):209-218, Winter 2012

Ashraf, M. A. et al.



Concentration mg/kg

Fig. 3. Concentration of heavy metals in different fish species

reporting very high concentrations whilst other samples measured relatively lower concentrations and in some cases no concentrations of the elements. Zinc and copper tended to be the least concentrated in the fish as compared to other elements measured. Concentrations of copper varied from <1.51 to 36.31 mg/Kg (dry wt.). The low copper values in the catchment fish may be due to the high chlorides content of the catchment since salt-water organisms are known to be more resistant to copper poisoning than freshwater organisms (Asagba et al., 2008). The greatest sources of zinc in humans are sea foods and meats (Reigart & Roberts, 1999). Concentrations of Zinc in the fish samples were relatively high (16.25 to 104.23 mg/Kg). The maximum permissible chromium levels for fish are 102 mg/kg as reported by (FAO, 1983). Arsenic is one of the most toxic elements to fish. Acute exposures can result in immediate death. Chronic exposures can result in the accumulation of the metalloid to toxic levels. In fish, bizarre morphological alterations, as well as early neoplastic alterations are produced in the live. Total arsenic results are similarly low, with only 5 of the 15 fish tissue samples having a total arsenic concentration above the detection limit of 0.11 mg/kg. The highest total arsenic concentration observed was 0.87 mg/kg for Hampala macrolepidota from the catchment. Total concentration of arsenic in fish samples taken from mining ponds are ranged between a minimum of 0.0025 to a maximum of 0.83 mg/ kg. The review concluded that arsenic concentrations in planktivorous fish are elevated over predatory and omnivorous fish. Arsenic does not biomagnify up the aquatic food chain as do some other chemicals. Concentration of lead ranged from 0.07 to 1.78 mg/kg. The highest concentration of zinc was recorded in the muscles of Rasbora elegans and Trichogaster trichopterus. Similarly concentration of tin were ranged from 56.34 to 153.45 mg/kg with highest concentration of tin were found in the Trichogaster trichopterus. The results showed that the concentrations of Pb and Sn, were higher in Trichogaster trichopterus fish, than in fish from other sampling locations showing that Trichogaster trichopterus is more susceptible to heavy metals accumulation but previously there are not reported any marked effect with high concentration of tin metal.

Health risk estimates in this study were calculated based on the integration of the data from heavy metals analysis and the assumed consumption rate based on US.EPA guidelines. The following assumptions were made based:

• Hypothetical body weights of 10 Kg for children between the ages of zero and one year, 30 Kg for children between the ages of one and eleven years and 70 Kg for adults

• Maximum absorption rate of 100% and a bioavailabilty rate of 100%

• Food consumption rate of fish in Malaysia given as 0.08 kg/day Hence, for each type of contaminat (heavy metal), the estimated daily dose (ED) (mg/Kg/day) was obtained using the equation detailed in the US. EPA handbook , 1989:

ED = Conc. of element of interest x Food consumtion rate / Body weight, kg

To estimate the health effects, the estimated lifetime average daily dose of each chemical was compared to its Reference Dose (RfD). The reference dose represents an estimate of a daily consumption level that is likely to be without deleterious effects in a lifetime. And the hazard index (HI) is estimated as the ratio of the estimated metal dose and the reference dose as depicted in US.EPA, 1996.

Hence, HI = ED/RfD

Where, ED = Estimated Dose and RfD = Reference Dose Estimated health risk associated with consumption of the fish is presented in (Table 3). Hazard index < 1 suggests unlikely adverse health effects whereas HI > 1 suggests the probability of adverse health effects. For adults, health risks estimates for copper, zinc, lead, arsenic and tin were lowest for the general public at the highest ingestion rate (Table 3) whilst the same elements were relatively

Sr. No.	Metals	Permissible Limits [*] mg/kg	Measured Value mean	Certified Value	Accuracy %
1	Lead	2	0.89 ± 0.01	0.85±0.13	78
2	Arsenic	0.026	0.44 ± 0.18	0.45 ± 0.15	98
3	Copper	30	89.30±0.42	106 ± 10.00	89
4	Zinc	100	$183.80{\pm}1.70$	180.00 ± 6.00	98
5	Tin	156	198±2.0	204±10.00	95

Table 2. Permissible levels of metals & validation of extraction and analysis with certified reference material

*MFA, 1983

Metal	RfD	Estimated Dose Time (mg/kg)		Hazard Index			
	-	0-1 Year	1-11 Year	Adult	0-1 Year	1-11 Year	Adult
Copper	3.70E-02	0.082	0.027	0.012	1.7	0.58	0.25
Zinc	4.70E-02	0.07	0.024	1.98	1.5	0.5	0.22
Lead	5.70E-05	0.088	0.056	2.56	1.92	0.71	3.44
Arsenic	7.00E-03	0.0065	0.0022	0.00093	0.9	0.007	0.0031
Tin 8.60E-03	0.4728	0.1576	0.0675	54.98	18.32	7.85	5.98

Table 3. Health risk estimates associated with heavy metals in fish species

higher for children at the highest ingestion rate. Even though the values do not pose any health hazards for the ordinary consumer, the levels are likely to be higher for families of fishmen since their meals are more fishdependent. The use of metal-based fertilizers by farmers in the ex-mining area should be monitored and regulated.

The accumulation of metals by the fish depends on the location, feeding behaviour, trophic level, age, size, duration of exposure to metals and homeostatic regulation activities of fish (Sankar et al., 2006; Kargin, 1996) has listed multiple factors that influence metals accumulation in fish such as season, physical and chemical properties of water. Knowledge of metals concentration in fish is important to management for various purposes such as risk of taking fish as part of diet and metals pollution control strategies. Most of fish are at top in aquatic food chain and have potential to accumulate high metals content even in mild polluted conditions. Therefore, metals concentration in fish could be used as an index to estimate level of pollution especially in aquatic bodies (Karadede-Akin & Unlu, 2007) even in the lake system. Although, fish muscle was reported have lowest metal concentrations compare to bone, gill and liver, this study focus metals in fish muscle since, people eat fish muscle and not others. Various studies on metal concentrations in fish samples has done previously (Findan et al., 2007; Chi et al., 2007; Ahmad and Shuhaimi, 2010; Laar et al., 2011). This study found Cu and Zn concentrations were almost as low as from unpolluted lake likes Eber (Findan et al., 2007). However, Sn and Pb concentrations were higher than fishes from Eber Lake and Taihu Lake (Chi et al., 2007). Both metals were probably contributed to the lake through various sources, such as agriculture activities likes oil palm and rubber plantations nearby catchment besides mining activities. A large scale of oil palm plantation was developed by Sime Darby Company nearby the catchment and the use of chemical fertilization could introduce Pb, Zn and Cu to the catchment. Excessive fertilizer may be transported to the catchment through surface run off. The catchment has several dredged out small and big ponds that could transport these

metals into the catchment. Furthermore, the inflow of water from the catchment into Selangor River can make the conditions more adverse. During rainy season or monsoon season also could bring pollutants from nearby agricultural areas including metals into the catchment. Nonetheless, the concentrations of these metals were still below the permissible limit (Table 2, Fig. 3) suggested by the (MFA, 1983).

This study indicates that although still below hazardous level, non-essential metals namely, Pb and Sn and essential metal namely Zn were quite high and continues input could risk the catchment ecosystem. Continual accumulation in fish muscle could transports metals to human through diet. Local aborigines surround the lake catch fish for their daily diet.

CONCLUSION

The concentrations of heavy metals in the fish samples were quite variable and no patterns of distribution and behaviour were noted. However, high concentrations of some heavy metals measured in the fish tissues inhabiting that the Bestari Jaya catchment were related to a high influx of metals as a result of pollution from the dredged mined out ponds thereby increased bioavailability to the fish. The high levels also suggested that the fish were capable of concentrating the metals in their bodies from the aquatic environment. The ability of the fish samples to incorporate the elements into their tissue is another important factor to consider for further study. The data presented in this report indicates that the species have preference to retain some metals in their tissues than other metals. Although, no immediate health risk was estimated from consumption of the fish species, risk prevention on the consumption of fish should therefore focus on reducing the volume of heavy metals discharged from agriculture area as well as mined out ponds into the catchment.

ACKNOWLEDGEMENT

The work reported in this paper is a basic research work carried out at Analytical Laboratory, Department of Chemistry and partially at Department of Geology, University of Malaya. I take this opportunity to express my gratitude to Ministry of Higher Education, Malaysia to provide me scholarship to complete my higher studies in Malaysia. Thanks also go to Institut Pengurusan Dan Pemantauan Penyelidikan, (IPPP) University of Malaya to provide me enough funding (Project No. PS355/2009C) for the completion of this valuable research.

REFERENCES

Abedin,M.Z. (1986). Atomic absorption spectrophotometric (AAS) analysis of zinc and manganese in Libyan fish and canned Tuna fish., Bull. Marine Biol. Res. Center, Libya, **7** (**2**), 46-59.

Ahmad, A. K. and Shuhaimi-Othman, M. (2010). Heavy metal concentrations in sediments and fishes from lake chini, Pahang, Malaysia. J. Biol. Sci., **10** (2), 93-100.

Ahmet, B. (1992). A case of mercury concentration in fish and human. Bull. Marine Biol. Res. Center, Libya, **9B** (2), 11-29.

Allen, H. E., Perdue, E. M. and Brown, D. S. (1993). Metals in Groundwater, Lewis Publishers, pp: 437.

Amundsen, P. A., Staldvik, F. J., Lukin, A. A., Kashulin, N. A., Popova, O. A. and Reshetnikov, Y. S. (1997). Heavy metals contamination in freshwater fish from the border region between Norway and Russia. Sci. Total Environ., **201** (3), 211-224.

APHA, (1981). Standards methods for the examination of water and wastewater. (14 Ed), Washington, D.C., USA.

Asagba, S. O., Eriyamremu, G. E. and Igberaese, M. E. (2008). Bioaccumulation of cadmium and its biochemical effect on selected tissues of the catfish. Fish physiol. Biochem., **34** (1), 61-69.

Ashraf, M. A., Maah, M. J. and Yusoff, I. (2011) Heavy metals accumulation in plants growing in ex tin mining catchment, Int. J. Environ. Sc. Technol. 8 (2), 401–416.

Bagenal, T. B. and Tesch, F. W. (1978). Age and Growth. in: Bagenal, T.B. (Ed.), Methods for Assessing of Fish Production in Freshwaters. 3rd Edn., No. 3, Blackwell Scientific Publication Ltd., 101-136.

Batvari, B. P. D., Kamala-Kannan, S., Shanthi, K., Krishnamoorthy, R., Lee, K. J. and Jayaprakash, M. (2007). Heavy metals in two fish species (Carangoidel malabaricus and Belone stronglurus) from Pulicat Lake, North of Chennai, Southeast Coast of India. Environ. Monit. Assess., **145** (1-3), 167-175.

Bermejo-Barrera, P., Moreda-Pineriro, A. And Bermejo-Barrera, A. (2001). Sample pretreatment methods for the trace elements determination in seafood products by atomic absorption spectrometry. Talanta, **57** (**5**), 969- 984.

Botson, F., Dassenakis, E., Panou, E. and Scoullos, M. M. (2004). Lead and cadmium transfer from a polluted stream to the marine environment. Rapp. Com. Int. Mer., **37** (1), 176-180.

Chi, Q., Zhu, G. and Langdon, A. (2007). Bioaccumulation of heavy metals in fishes from Taihu Lake, China. J. Environ. Sci., **19** (**12**), 1500-1504.

Chirila, E., Birghila, S. and Capota, P., (1999). Biogeochemical characterization of an ecosystem using ICP-AES. S. Afr. J. Chem., **52**, 154-156.

El-Mehdi, H. T. (1987). Mercury in Bluefin Tuna fish (*Thunnus thynnus*) caught from Jamahiriya Coast, Bull. Marine Biol. Res. Centre, Libya, **7**, 46-59.

FAO, (1983). Food and Agriculture Organization, Compilation of Legal Limits for Hazardous Substances in Fish and Fishery Products. Food and Agriculture Organization of the United Nations, Rome, Italy, 5-100.

Feedman, B. (1989). The impacts of pollution and other stresses on ecosystem structure and function. Environ. Ecology. Academic press, London.

Fidan, A. F., Cigerci, I. H., Konuk, M., Kucukkurt, I., Aslan, R. and Dundar, Y. (2007). Determination of some heavy metal levels and oxidative status in *Carassius carassius* L., 1758 from Eber Lake. Environ. Monit. Assess., **147** (1-3), 35-41.

Forstner, U. and Wittmann, G. T. W. (1983). Metal pollution in the aquatic environment. Springer-Verlag, Berlin, 30-61.

Goldberg, F. D., Bowen, V. T. and Farrington J. W. (1983). The mussel watch. Estuarine, Coastal and Shelf Science, **16** (1), 69-93.

Güven, K., Özbay, C., Ünlü, E. and Satar, A. (1999). Acute lethal toxicity and accumulation of copper in Gammarus pulex (L.) (Amphipoda). Tr. J. Biol., **23** (1), 51-521.

Harms, U. (1975). The levels of heavy metals (Mn, Fe, Co, Ni, Cu, Zn, Cd, Pb, Hg) in fish from on shore and off shore waters of the German bight. Z. Lebensum. Unters. Forsch., **157** (3), 125-130.

Huijuan, J., Huifeng, R., Schuichi, S., Hideaki, E. and Tetsuhito, H. (2005). Comparison of pretreatment condition of cadmium in fish samples and diet by microwave digestion method for ICP-AES, J. Tokyo Uni. Marine Sc. Technol., **1** (**1**), 41-46.

Jafari, N. G. and Gunale, V. R. (2006). Hydobiological study of algae of an urban freshwater river. J. Appl. Sci. Environ. Mgt., **10** (2), 153-158.

Jaffar, M., Ashraf, M. and Rasool, A. (1988). Heavy metal contents in some selected local freshwater fish and relevant waters, Pak. J. Sci. Ind. Res., **31** (3), 189–193.

Jefferies, D. J. and Freestone, P. (1984). Chemical analysis of some coarse fish from a Suffolk River carried out as part of the preparation for the first release of Captive bred otters. J. Otter Trust., **8** (3), 17-22.

Jones, R. E., Petrell, R. J. and Pauly, D. (1999). Using modified length-weight relationship to assess the condition of fish. Aquat. Eng., **20** (**4**), 261-276.

Karadede-Akin, H. and Unlu, E. (2007). Heavy metal concentrations in water, sediment and fish and some benthic

organisms from Tigris River, Turkey. Environ.Monit. Assess., **131** (1-3), 323-337.

Kargin, F. (1996). Seasonal changes in levels of heavy metals in tissues of Mullus barbatus and Sparus aurata collected from ýskenderun gulf (Turkey). Water Air Soil Pollut., **90** (**3-4**), 557-562.

Kinne, O. (1984). Marine Ecology, Ocean management, V5, John Wiley, 618-627.

Laar, C., Fianko, J. R., Akiti, T. T., Osae, S. and Brimah, A. K. (2011). Determination of Heavy Metals in the Black-Chin Tilapia from the Sakumo Lagoon, Ghana Res. J. Environ. Earth Sci. **3** (1), 8-13.

Mendez, H., Alava, F., Lavilla, I. and Bendicho, C. (2002). Ultrasonic extraction combined with fast furnace analysis as an improved methodology for total selenium determination in seafood by electrothermalatomic absorption spectrometry, Analtica chemical Acta, **452** (2), 217-222.

Meybeck, M., Chapman, D. and Helmer, R. (1996). Global Fresh Water Quality: A second Assessment. Blackwell Reference. Oxford, 306-310.

MFA, (1983). Malaysian Food Act, Malaysian Food and Drug. MDC Publishers Printer Sdn. Bhd., Kuala Lumpur.

Milne, A, Landing, W., Bizimis, M. and Morton, P. (2010). Determination of Mn, Fe, Co, Ni, Cu, Zn, Cd and Pb in seawater using high resolution magnetic sector inductively coupled mass spectrometry (HR-ICP-MS) Analytica Chimica Acta, **665** (2), 200-207.

Moeller, S., Croning, D. R. and Apweiler, R., (2001). Evaluation of methods for prediction of membrane spanning regions. Bioinformatics, **17** (7), 646-653.

Mohsin, A. K. M. and Ambak, M. A. (1991). Ikan Air Tawar Di Semenanjung Malaysia. Dewan Bahasa Dan Pustaka, Kuala Lumpur, ISBN: 9836219560.

Mustafa, T. (2003). Determination of heavy metals in fish samples of the middle Black Sea (Turkey) by graphite furnace atomic adsorption spectrometry, Food Chem., **80** (1), 119-123.

Petisleam, T., Popescu, I. V., Ciupina, V. and Belc, M. (2003). Trace elements from marine environmental samples of Black Sea determined by atomic absorption spectrometry, Proceedings of 4 th International Balkan Workshop on Applied Physics, Oviaius University Press, Constantza Romania, 103-108.

Olaifa, F. E., Olaifa, A. K., Adelaja, A. A. and owolabia, A. G. (2004). Heavy metal contamination of Clarias gariepinus from a lack and fish farm in Ibadan, Nigeria, African J. Biomed. Res., **7** (1), 145-148.

Perez, Cid, B., Boia, C., Pombo, L. and Rebelo, E. (2001). Determination of trace metals in fish species of the Ria de Aveiro (Portugal) by electro-thermal atomic absorption spectrometry, Food chem., **75** (1), 93-100.

Petisleam, I. V., Popescu, I. V, Ciupina, V. and Belc, M. (2007). Considerations regarding Cu and Ni determination

from marine environmental samples of Black Sea Using FAAS, Rom. J. Phys., **52** (3-4), 441-444.

Phillips D. J. H. (1980). Quantitative aquatic biological indicators: Their use to monitor trace metal and organochlorine pollution. Applied Science Publ., London, 136.

Reigart, R. J. and Roberts J. R. (1999). Recognition and management of pesticide poisonings. 5th Edn., U.S. Environmental Protection Agency, Office of Pesticide Programs, Washington, D.C.

Sanchez Lopez, F. J., Gil Garcia, M. D., Sanchez Morito, N. P. and Martinez Vidal, J. L. (2003). Determination of heavy metals in Cray fish by ICP-MS with a microwave-assisted digestion treatment, Ecotoxicol. Environ. Safe., **54** (2), 223-228.

Sankar, T. V., Zynudheen, A. A., Anandan, R. and Nair, P. G. C. (2006). Distribution of organochlorine pesticides and heavy metal residues in fish and shellfish from Calicut region, Kerala, India. Chemosphere, **65** (**4**), 583-590.

Sericano, J. L., Wade, T. L. and Jackson, T. J. (1995). Trace organic contamination in the Americas: An overview of the US national status and trends and the international mussel watch progammes. Mar. Pollut. Bull., **31** (**4-12**), 214-225.

Sperling, K. R. (1988). Determination of heavy metals in sea water and in marine organisms by graphite furnace AAS, J. Anal. Chem., **332** (6), 565-567.

Vaidya, O. C. and Rantala, R. T. T. (1996). A comparative study of analytical methods: determination of heavy metals in mussels (*Mytilus edulis*) from Eastern Canada. Int. J. Environ. Anal. Chem., **63 (3)**, 179–185.

Wade, T. L., Sweet, S. T. and Klein, A. G. (2008). Assessment of sediment contamination in Casco Bay, Maine, USA. Environ. Poll., **152** (3), 505-521.

Williams, J. E. (2000). The coefficient of condition of fish. in: Schneider, J.C. (Ed.), The manual of fisheries survey methods II: with periodic updates. Michigan department of natural resources, Fisheries special report 25, Ann Arbor, Michigan.

Yilmaz, A. B. and Dogan, M. (2007). Heavy metals in water and in tissues of himri (*Carasobarbus luteus*) from Orontes (Asi) River Turkey. Environ. Monit. Assess., **144** (1-3), 437-444.