Int. J. Environ. Res., 6(4):893-902, Autumn 2012 ISSN: 1735-6865

Assessment of Water Quality Parameters in Mangrove Ecosystems Along Kerala Coast: A Statistical Approach

Manju, M. N.*, Resmi, P., Gireesh Kumar T.R., Ratheesh Kumar, C.S., Rahul, R., Joseph, M. M. and Chandramohanakumar, N.

Department of Chemical Oceanography, School of Marine Sciences, Cochin University of Science and Technology, Kochi, Kerala, India

Received 25 Aug. 2011;	Revised 2 June 2012;	Accepted 12 June 2012
------------------------	----------------------	-----------------------

ABSTRACT: Water samples were collected from five selected mangrove ecosystems along Kerala coast, North of Cochin, on seasonal basis and analysed for various hydrographic parameters. To explain spatiotemporal variations and the processes controlling their distribution, ANOVA (two factor without replication), correlation and principal component analysis were carried out. Concentration of total nitrogen varied from 10.26 to 188.38 μ M and total phosphorous from 1.53 to 22.88 μ M, with comparatively higher concentration in monsoon season which could be attributed by land run off. Chlorophyll pigments recorded significant seasonal variation ranging from ND to 40.86 μ g/L (chlorophyll a), ND to 6.00 μ g/L (Chlorophyll b) and ND to 13.80 μ g/L (Chlorophyll c). Both monsoon and post monsoon seasons recorded higher concentration of pheophytin compared to Chlorophyll a and the maximum concentration of chlorophyll was observed during pre monsoon. Station 2 (Pappinissery) exhibited elevated concentration of nutrients (especially NH₄⁺) and lower DO content which pointed towards the reducing environment by anthropogenic stress. Factor analysis revealed six components which explained 92.77% of the total variance. It also described the processes like diagenesis, sediment remineralisation, anthropogenic activities, tidal and river influx which make the ecosystem highly complex.

Key words: Mangrove, Northern Kerala coast, Water quality, Nutrients, Major ions, Chlorophyll

INTRODUCTION

Anthropogenic activities have remarkably influenced the quality of ecosystems in aquatic environments (Jiang et al., 2011; Pei et al., 2011; Jing and Zhiyuan, 2011; Hallare et al., 2011; Siddiqui, 2011; Feng et al., 2012; Clemente et al., 2012; Kiteresi et al., 2012). Mangroves dominate majority of the world's tropical and subtropical coastline, forming 15 million hectares of forests worldwide, which provide habitat for rich biodiversity, ranging from bacteria, fungi and algae through to invertebrates, birds and mammals (FAO, 2004). The high productivity in mangrove ecosystems is often attributed to greater litter degradation rates and efficient recycling of nutrients, which are supplied by both autochthonous litter and allochthonous inputs from natural and anthropogenic sources (Lee, 1990; Kazungu et al., 1993; Bouillon et al., 2002). The stability of the mangrove is influenced by salinity, soil type and chemistry, nutrient content and dynamics, physiological tolerance, predation and

competition at local level (Smith *et al.*, 2003). Besides the environmental stress factors, habitat destruction through human encroachment has been the primary cause of mangrove loss.

Despite the fact that nutrients in the tropical marine ecosystems are generally low (Qasim & Wafar, 1990), mangrove ecosystems are considered to be the most productive and complex ecosystems (Naskar & Mandal, 1999) with the ambience being rich in organic production and nutrients (Manson *et al.*, 2005). Primary production by microalgae can also contribute to the complexity of the temporal and spatial variation of dissolved compounds (Harrison *et al.*, 1997; Trott & Alongi, 1999). The nutrients such as inorganic phosphorus, nitrogen, potassium, and organic carbon are provided to adjacent coastal and marine, as well as terrestrial ecosystems through active and passive transport (Odum, 2000; Gonneea *et al.*, 2004). Changes in proportions of dissolved nutrients in mangrove

^{*}Corresponding author E-mail: manjubhat2009@gmail.com

water are primarily caused by supply from agricultural seepage, aquaculture effluents, human settlements, etc., and leads to changes in nutrient stoichiometric ratios (Si:N, N:P and Si:P), significantly impacting the coastal food web dynamics (Moncheva *et al.*, 2001; Zhao *et al.*, 2005).

Total area of mangroves in India is about 4461 km², constituting 0.41% of countries geographical area (Jagtap et al., 2002). Mangrove vegetation in Kerala is now confined largely to river mouths and tidal creeks and there has been no significant mangrove cover south of Cochin in Kerala coast (Kurien, et al., 1994; FSI, 2003). Inland and coastal wetlands around Cochin have been lost over years due to reclamation, conversion to industrial use, dumping of solid waste, discharge of untreated sewage from municipal waste, effluents from industries and encroachment for construction. A recent study by Radhakrishnan et al., 2006 pointed out mangrove vegetation in four northern districts of Kerala - Kasargod, Kannur, Kozhikode and Malappuram - is approximately 3,500 hectare, which represents about 83 percent of mangrove cover in the State.

The conservation, management and sustainable development of the mangroves depend on the maintenance of hydrogeochemical characteristics of the system. Most of the research works on hydrochemical aspects of mangrove ecosystems are limited to central part of Kerala and studies on northern Kerala coast has not been reported so far. The present study is a preliminary attempt to reveal ecological health and status of mangrove ecosystems along northern Kerala Coast for management strategy and conservation. Objective of the study is to evaluate the spatial and temporal variation of general hydrographical parameters along with chlorophyll pigments for water samples collected from the selected mangrove ecosystems, employing statistical methods.

MATERIALS & METHODS

Water samples were taken from five mangrove ecosystems located at Kannur and Calicut districts of northern Kerala on west coast of India. The sampling locations are depicted in Fig. 1. Station 1 (Thalassery), is the mangrove ecosystem found in the vicinity of Arabian Sea, is fringed with dense growth of *Rhizophora spp.* and *Avicennia officinalis, Acanthus illcifolius, Excoecaria agallocha, Aegiceros corniculatum* and *Thespesia populnea* with isolated column of old trees of *Sonneratia sp.* and *Kandelia candel.* Station 2 (Pappinissery) is formed on the banks of Valapattanam estuary (area of about 20 hectare) by covering a distance of 4-5 km from the coastline. Major mangrove species found at this site includes: Avicennia, Rhizophora, Kandelia and Acanthus with isolated growths of Aegiceras corniculatum and creeper Derris trifoliata. Station 3 (Pazhayangadi) is situated at a distance about 3-4 km from coastline was found to be almost free from human activities. Major species found at this station include: Avicennia marina, Avicennia officinalis, Aegiceras corniculatium and Rhizophora mucronata. Station 4 (Kunjimangalam) is situated in the estuarine environment formed by Pullamcode puzha and Kunjimangalam river which is located at 2 km away from coastline with an area of around 18 hectares. The major species occurred in this region include: Avicennia, Acanthus ilicifolius, Rhizophora mucronata, Rhizophora apiculata, Kandelia candel, Clerodendron inerme, Aegiceros corniculatium and Excoecaria agallocha. Station 5 (Kadalundi) is situated on the banks of Kadalundi river which joins the Arabian Sea through a permanent bar mouth. Mangroves cover an area of 10 hectare which includes Rhizophora mucronata, Excoecaria agallocha, Aegiceros corniculatum, and Acanthus ilicifolius Avicennia officinalis etc. (CMFRI, 2002).

There are three seasonal conditions prevailing in Kerala viz. monsoon (June-September), post monsoon (October-January) and pre monsoon (February-May). Sampling was carried out during October 2009, May 2010 and August 2010 representing three seasons. Water samples were collected using a clean plastic bucket, transferred to clean plastic bottles and transported to the laboratory on ice and stored in a deep freezer (-20°C) till analysis. Samples were collected in triplicate from each station and average value for each parameter was reported. pH of water samples was measured in situ and salinity was estimated by Mohr-Knudsen method (Muller, 1999). Modified Winkler method was used for the estimation of dissolved Oxygen (Hansen, 1999). Alkalinity of the water samples were estimated by the method of Koroleff (Anderson et al., 1999). Nutrients (nitrite, nitrate, phosphate, silicate,) were estimated spectophotometrically using standard methods (Grasshoff, 1999). The total nitrogen and total phosphorous content were measured after alkaline persulphate oxidation (Hansen & Koroleff, 1999). The major elements like sodium, potassium, lithium were analyzed using flame photometer, calcium and magnesium were determined using EDTA titration, iron was analyzed using phenanthroline method and nephelometric method was employed for sulphate determination (APHA, 1995). Chlorophyll pigments and pheophytin in water samples were filtered through 0.45 µm glass fiber filter paper, extracted using 90% acetone and measured spectrophotometrically (APHA, 1999).

894



Fig. 1. Map showing location of sampling stations

Analysis of variance (ANOVA) was employed to evaluate the temporal and spatial variation in concentration of the analysed parameters. The effect of natural and anthropogenic flux was evaluated using factor analysis by applying Varimax rotation.

RESULTS & DISCUSSION

The seasonal and spatial variations of the analysed parameters are depicted in Fig. 2. pH displayed significant seasonal variation (P< 0.01) and comparatively lower values were detected during post monsoon with its minimum was recorded at station 2. In post monsoon, pH varied from 7.10 and 7.40 (avg.7.24 \pm 0.11), in pre monsoon 7.20 to 8 (avg.7.82 \pm 0.35), and in monsoon it varied from 7.81 to 8.05 (avg.7.92 \pm 0.11). A profound seasonal variation (P<0.01) was noticed for salinity ranging from 4.26 to 9.25 ppt (avg.6.73 \pm 2.32 ppt; post monsoon), 29.31 to 35.97 psu (avg.7.89 \pm 11.44 ppt; monsoon). Dissolved Oxygen (DO) showed variation in concentration from 2.86 to 6.41 mg/L (avg.4.66 ± 1.37 mg/L), 0.51 to 9.34 mg/ L (avg.4.52±3.34 mg/L) and 3.76 to 7.35 mg/L (avg.5.82±1.42 mg/L) during post monsoon, pre monsoon and monsoon seasons respectively. Alkalinity displayed significant seasonal variation (P <0.01), exhibiting higher values during pre monsoon and lower values during monsoon. It varied from 44.55 to 79.20 mgCaCO₂/L during post monsoon (avg.55.44±13.44 mgCaCO₂/L), 121.25 to 167.33 mg CaCO₂/L (avg.135.8±18.78 mg CaCO₂/L) and 22.31 to 83.42 mgCaCO₃/L (avg.42.49±24.51 mgCaCO₃/L), pre monsoon and monsoon seasons respectively. Nitrite concentration ranged from 0.15 to 0.55 µM (avg.0.28±0.16 µM; post- monsoon), 0.29 to 0.99 µM (avg.0.63 \pm 0.25 μ M; pre monsoon) and 0.31 to 0.99 μ M (avg. $0.50\pm0.28\,\mu$ M; monsoon) with a minimum of 0.15 µM from station 4 during post monsoon. Nitrate displayed a significant seasonal variation (P<0.01) and it recorded variation in concentration from 1.37 to 8.43µM (avg.3.63±2.82 µM; post monsoon), 0.29 to 4.33 μ M (avg.1.82 \pm 1.59 μ M; pre monsoon) and 3.14 to 20.79 Water Quality Parameters in Mangrove Ecosystems



Fig. 2. Variation of water quality parameters in mangrove ecosystems

μM (avg.13.62±6.70 μM; monsoon). Ammonia content varied from 4.73 to 27.95 µM during post monsoon $(avg.18.20 \pm 10.13 \ \mu M)$, ND to 98.09 μM (avg.22.38±42.45 µM) and 5.52 to 80.36 µM (avg.42.15±34.67 µM), pre monsoon and monsoon seasons respectively. Inorganic phosphate showed significant seasonal variations and its concentration varied from 0.59 to 1.37µM (avg.0.86±0.32µM) during post monsoon season, 1.32 to 6.56µM (avg.3.49±1.91µM) during pre monsoon and 9.98 to $15.07 \,\mu$ M (avg. $13.09 \pm 2.37 \mu$ M) during monsoon season. Significant seasonal variation was exhibited by silicate (P<0.05) and its content ranged from 7.33 to 47.79 μ M $(avg.22.72 \pm 16.10 \mu M; post monsoon), 15.55 to$ 106.60µM (avg.41.08±38.48 µM; pre monsoon) and 44.98 to 80.76 µM (avg.70.9±15.02 µM; monsoon). Concentration of total nitrogen (TN) was found to vary from 34.55 to 63.69 μ M (avg.46.17 \pm 12.20 μ M; post monsoon), 10.26 to 102.78 µM (avg.55.19±40.1 µM; pre monsoon) and 124.78 to 188.38 μM (avg.163.68±29.67µM; monsoon). A significant seasonal variation was observed for TN (P<0.01), with its minimum content reported during pre monsoon and maximum during monsoon (station 5). Total phosphorous (TP) content ranged from 2.96 to 8.61 μ M (avg.5.87 \pm 2.30 μ M), 1.53 to 6.65 μ M (avg.3.67 \pm 1.87 μM), and 10.04 to 22.88 μM (avg.19.29±5.77 μM) during post monsoon, pre monsoon and monsoon seasons respectively. TP also displayed highly significant seasonal variation (P<0.01) with its maximum concentration recorded during monsoon and the minimum during pre monsoon.

Significant seasonal variation for chlorophyll pigments were observed from ANOVA (P < 0.05). Chlorophyll a (chl a) recorded variation in concentration from 0.77 to 17.29 µg/L (avg. 5.82 $\pm 6.75 \mu g/L$; post monsoon), 2.18 to $40.86 \mu g/L$ $(avg.16.07\pm15.66; \mu g/L; pre monsoon)$ ND to $3.09 \mu g/L$ (avg.1.47±1.41 µg/L; monsoon). The samples collected during pre monsoon recorded comparatively higher chl a content (station 4) whereas lower concentration was reported during monsoon. Concentration of chlorophyll b (chl b) varied from ND to1.61 µg/L (avg.0.45 $\pm 0.66\,\mu g/L$), 1.12 and 6.00 $\mu g/L$ (avg.3.04 ± 1.89 μ g/L) and ND to 1.71 μ g/L (avg.0.96 \pm 0.89 μ g/L) during post monsoon, pre monsoon and monsoon seasons respectively. Comparatively higher chl b content was recorded at station 5 (pre monsoon). Chlorophyll c (chl c) ranged from ND to $4.34 \,\mu\text{g/L}$ (avg. $1.58 \pm 1.73 \,\mu\text{g/L}$), from 1.70 to 13.80 $\mu g/L$ (avg.6.50 \pm 4.67 $\mu g/L$) and ND to $2.44 \,\mu\text{g/L}$ (avg.0.67 \pm 1.06 $\mu\text{g/L}$) during post monsoon season, pre monsoon and monsoon seasons respectively. All stations recorded higher pheophytin content during post monsoon season with a maximum at station 4 (31.01 µg/L). It varied from 4.01 to 31.01 µg/L (avg.17. 96±12.84µg/L; post monsoon), 1.01 to 22.27µg/L (avg.8.68±8.56µg/L; pre monsoon) and ND to 23.20 µg/L (avg.5.59±9.94 µg/L; monsoon).

Among the major elements studied, calcium content found to vary from 21.78 to 172.51 mg/L (avg. 85.99±58.74 mg/L) during post monsoon, 148.37 to 453.13 mg/L (avg. $266.38 \pm 126.81 \text{ mg/L}$) during pre monsoon and 8.02 to 108.27 mg/L (avg. 30.45 ±43.31 mg/L) during monsoon. Concentration of magnesium ranged from 65.44 to 540.23 mg/L (avg.260.23±175.99 mg/L; post monsoon), 616.65 to 1362.8 mg/L (avg.1017.77±284.73 mg/L; pre monsoon) and 8.63 to 621.09 mg/L (avg.141.39 ±268.36 mg/L; monsoon season). The sodium content in water samples varied from 434 to 2523 mg/L (avg.1249±777mg/L) during post monsoon, 4000 to 7400 mg/L (avg. 5700±1245mg/L) during pre monsoon and 50 to 2600 mg/L (avg.626±1105.23 mg/L) during monsoon seasons. Concentration of potassium ranged from 18 to 105 mg/ L (avg.56.22±31.9 mg/L), 160 to 320 mg/L (avg. 228.57±68.78 mg/L) and 3.3 to 130 mg/L (avg.31.34±55.21 mg/L) during post- monsoon, pre monsoon and monsoon seasons respectively. The sample from station 3 recorded higher concentration for sodium and potassium during pre monsoon. Lithium content varied from 0.2 to 1.0 mg/L during post monsoon season, 1.6 to 3.2 mg/L during pre monsoon and ND to 1.4 mg/L during monsoon.

The sulphate content varied from 235.24 to1086.5 mg/L (avg.594.93 \pm 359.75 mg/L; post monsoon), 1700 to4166.67 mg/L (avg.3104.76 \pm 1057.09 mg/L; pre monsoon) and 12 to 966.67 (avg.224.4 \pm 415.41 mg/L; monsoon). The maximum concentration of sulphate was observed at station 4 during pre monsoon and the minimum was recorded at station 2 during monsoon. The iron content ranged from 0.089 to 0.85 mg/L (avg.0.31 \pm 0.31 mg/L), 0.078 to 0.624 mg/L (avg.0.31 \pm 0.25 mg/L) and 0.04 to 1.43 mg/L (avg.0.43 \pm 0.58 mg/L) during post monsoon, pre monsoon and monsoon season respectively.

Salinity exhibited minimum value during monsoon season due to fresh water runoff. The maximum was reported from station 4 which might be due to the land locked nature of the sampling site. Compared to other stations, stations 1 and 5 are situated in the close proximity of Arabian Sea and therefore prominent tidal activity can alter salinity of these systems. In aquatic systems, oxygenation is the result of an imbalance between the process of photosynthesis, degradation of organic matter, re-aeration (Granier *et al.*, 2000), and physicochemical properties of water (Aston, 1980).The organic pollution by domestic sewage at station 2 resulted in the depletion of dissolved oxygen. The dissolved oxygen super saturation (132 %) was observed at station 4 (pre monsoon) could be resulted by the photosynthesis by phytoplankton, confirmed by the higher chlorophyll content. Limited tidal rhythm and flushing also contributed to the dissolved oxygen super saturation. Dissolved gas super saturation (DGS) can be produced in rivers and lakes which have high densities of plankton, aquatic plants, and algae (White *et al.*, 1991).

Nitrate exhibited higher concentration during monsoon (station 5) and lower content during pre monsoon (station 4). According to Solanki *et al.*, 2010, the proportion of different forms of nitrogen in any water body is determined by the balance between assimilation, mineralization, nitrification, denitrification and nitrogen fixation. Station 2 received large quantity of domestic garbage, poultry waste resulting to the reducing environment and recorded elevated concentration of nutrients (especially NH₄⁺) and low DO. At stations 2 and 3, burrowing crabs produce sediment micro pores which are reducing in nature resulting in the build up of ammonium, by the process of anaerobic ammonification (Smith *et al.*, 1991).

Comparatively higher content of phosphate was recorded during monsoon and lower content during post monsoon season. During the monsoon, nitrate and phosphate level in water increases considerably due to land drainage and anthropogenic input. Faecal contamination and excreta of birds enhanced the higher level of phosphate in these mangrove ecosystems. The lower content observed during post monsoon and summer seasons might be due to decreased runoff, adsorption to sediments, and utilization by phytoplankton (Ramakrishnan et al., 1999). Phosphate was found to be positively correlated with pH (Table 1), which may be due to the release of phosphorous from iron and aluminium oxides by ligand exchange reactions (Boers, 1991) under favourable conditions. Water samples from station 1 during post monsoon recorded a minimum silicate content of 7.33 µM and sample from station 2 during pre monsoon showed higher value of 106.60 µM. Weathering process and land run-off mainly contribute to silicate concentration of these mangrove ecosystems.

The ratio of dissolved inorganic nitrogen to dissolved inorganic phosphate (DIN: DIP) ranged from 10.63 to 42.60 (avg.25.70 \pm 12.71; post monsoon), 0.62 to 15.25 (avg.5.75 \pm 5.90; pre monsoon) and 0.82 to 8.73 (avg.4.43 \pm 3.29; monsoon). Maximum value for this ratio was estimated at station 2 during post monsoon and the minimum at station 3 during pre monsoon. All stations were found to be nitrogen depleted during pre monsoon and monsoon which might be due to the slow regeneration of NO₃⁻ compared to PO₄³⁻.

Total nitrogen to total phosphorus (TN: TP) ratio in the study region varied from 4.27 to 12.39 (avg.8.79±3.33; post monsoon), 6.70 to 25.06 $(14.40\pm7.01; \text{ pre monsoon})$ and 6.33 to 12.42 (8.84 $\pm2.22;$ monsoon). Samples collected during pre monsoon exhibited maximum value for TN: TP ratio and the minimum recorded during monsoon. At stations 3 and 4, major portion of the total nitrogen was in organic form, because naturally occurring organic nitrogen from decaying plant remains are slowly converted to NH_{4}^{+} state by biological degradation (Valentina & Joseph, 2005). Sewage from excreta being richer in nutrients seemed to enhance the eutrophication problem in the study region. Uptake by trees and benthic microorganisms and sorption into the sediment matrix can considerably reduce nutrient concentrations in pore-water which can make the mangrove forest to an effective sink for nutrients (Alongi, 1996). From the interrelationships between nutrients (Table.1), it could be inferred that they were originated from the same source.

Chlorophyll displayed highly significant seasonal variation (P<0.01) with higher concentration during pre monsoon. Chlorophyll is considered as the most reliable index of phytoplankton biomass. Also, chl a to pheophytin ratio provides the first hand information on the physiological status of phytoplankton. This ratio ranged from 0.15 to 0.56 (average 0.28±0.16; post monsoon), 1.63 to 2.14 (average 1.88±0.19; pre monsoon) and ND to 1.54 (average $0.77\pm0.0.76$; monsoon). Higher concentration of pheophytin compared to chl a during post monsoon and monsoon seasons indicated the presence of more detritus matter in these environments, which could be attributed to decomposition of organic matter, from the sediment and community structure, harbouring in the surrounding water (Tripathy et al., 2005). In pre monsoon season, the reverse trend was observed due to the growth of phytoplankton in the high light intensity and low turbulent waters of mangrove ecosystems. Absence of correlation between chlorophyll pigments and nutrients indicated that nutrients have no significant role on primary production. The presence chl b revealed the contribution of green algae to the productivity of mangrove ecosystems. Chl b to chl a and chl c to chl a ratios are <1 suggesting the possibility of healthy phytoplankton populations in lower light intensity and low turbulent areas (Takahashi & Nakamoti, 1972).

The high correlations obtained between salinity and cation concentrations pointed towards the same origin of investigated elements by intrusion of Sea water. At stations 2, 3 and 4 seasonal patterns of measured parameters were modulated by precipitation, evaporation and temperature and at stations 1 and 5 along with these factors, tidal influx of Sea water also

Mumano. Mumano. </th <th></th> <th>Hu</th> <th>salinity</th> <th></th> <th>Alkalinity</th> <th>-ON</th> <th>- ON</th> <th>Ammonia</th> <th>PO.³⁻</th> <th>SiO.²⁻</th> <th>Ę</th> <th>đL</th> <th>ť</th> <th>Ċ</th> <th>Ċ</th> <th>theon</th> <th>DIN</th> <th>Ma</th> <th>Ğ</th> <th>Š</th> <th>Х</th> <th>:1</th> <th>S</th>		Hu	salinity		Alkalinity	-ON	- ON	Ammonia	PO. ³⁻	SiO. ²⁻	Ę	đL	ť	Ċ	Ċ	theon	DIN	Ma	Ğ	Š	Х	:1	S
Mit I Mit I <th>Parameters</th> <th>hid</th> <th>заштыу</th> <th>3</th> <th>(mmaarc</th> <th>1402</th> <th>504</th> <th></th> <th>ф Т</th> <th>4010</th> <th>, i</th> <th>=</th> <th>Ça</th> <th>5</th> <th>ç</th> <th>hytin</th> <th>NICT</th> <th>SIM</th> <th>Ç</th> <th>110</th> <th>4</th> <th>1</th> <th></th>	Parameters	hid	заштыу	3	(mmaarc	1402	504		ф Т	4010	, i	=	Ça	5	ç	hytin	NICT	SIM	Ç	110	4	1	
Motione in the sector of	Hq	1																					
Monorizatione di constructione di	Salinity	0.35	1																				
Main Main <th< td=""><td>DU</td><td>0.36</td><td>0.01</td><td>-</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></th<>	DU	0.36	0.01	-																			
With with with with with with with with w	NO. ⁻	0.09	0.79**	-0.29	1 051	-																	
Momenta 10. 40. 10.	0N	1 1 0	14:0	01.0	100	1	-																
POL 433 0.0 0.33 0.34 0.4 0.35 0.34 0.4 0.35 0.34 0.4 0.35 0.34 0.4 0.35 0.34 <th0.34< th=""> <th0.34< th=""> <th0.34< th=""></th0.34<></th0.34<></th0.34<>	Ammonia	/ 6.0	97 N-	0.42		c1.0	-																
BU U2 U2<	PO_4^{3-}	-0.10	-0.28	-0.24	0.06	0.23	0.36	-															
Integrity 01 02 03 <	SiO_4^{2-}	0.56*	-0.24	0.17	-0.40	0.06	0.77**	* 0.44	-														
IP 0.1 0.2 0.4 0.4 0.7 <th0.7< th=""> 0.7</th0.7<>	NL	0.14	-0.12	-0.15	-0.05	0.16	0.60**	* 0.73**	0.76**	1													
Understand Underst	ТР	0.42	-0.33	0.27	-0.45	-0.04	0.75**	* 0.48	**960	0.74^{**}	1												
	Ca	0.34	-0.50	0.17	-0.67**	-0.20	0.70**	* 0.29	0.91^{**}	0.60^{**}	0.91**	-											
	ç	0.21	0.54*	0.29	0.48	0.07	-0.49	-0.40	-0.35	-0.40	-0.29	-047	Т										
	ට	0.44	0.53*	0.11	0.55*	0.53*	-0.33	-0.14	-0.12	-0.12	-0.20	-026	0.56^{*}	1									
$ \begin{array}{{ \ \ \ \ \ \ \ \ \ \ \ \ \$	Phe ophytin	0.28	0.64 **	0.25	0.62**	0.22	-0.48	-0.28	-0.31	-0.29	-0.28	-048	0.96**	0.72**	-								
	DIN	-0.25	-0.18	0.34	-0.09	-0.33	-0.30	-0.31	-0.35	-0.34	-0.21	-029	0.49	0.06	0.41	1							
	Mg	-0.02	-0.30	-0.14	-0.04	0.25	0.52*	0.98**	0.55*	0.79**	0.58*	040	-0.46	-0.19	-0.35	-034	1						
	Ca	0.06	0.79**	-0.29	0.92**	0.42	-0.48	0.11	-0.36	-0.03	-0.41	-0.64**	0.39	0.33	050	-023	0.01	1					
$ \begin{array}{ ccccccccccccccccccccccccccccccccccc$	Na	-0.02	0.64**	0.54*	0.78**	0.14	-0.53*	-0.03	-0.41	-0.03	-0.49	-0.58*	0.35	0.37	0.43	-018	-0.13	0.74**	-				
$ \begin{array}{r[r]lllllllllllllllllllllllllllllllllll$	K	0.15	0.84**	-0.32	0.92**	0.34	0.52*	-0.04	-0.38	-0.09	-0.47	-0.65**	0.48	0.44	0.57*	-017	-0.14	0.94^{**}	0.88**	-			
$ \sum_{0,4}^{2} \begin{array}{cccccccccccccccccccccccccccccccccccc$	Li	0.09	0.77**	-0.43	0.90**	0.33	-0.49	0.03	-0.37	-0.01	-0.46	-0.63**	0.37	0.38	0.47	-023	-0.06	0.91^{**}	0.93**	**860	-		
$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\mathrm{So_4}^{2\text{-}}$	0.09	0.77**	-0.45	0.90**	0.40	-0.43	0.12	-0.31	0.08	-0.41	-058*	0.32	0.44	0.45	-030	0.03	0.91**	0.91^{**}	**960	0.98**	1	
0.27 -0.25 0.06 -0.18 -0.14 0.04 0.00 0.20 0.07 0.17 0.19 -0.09 0.23 0.02 0.40 0.01 -0.33 -0.07 -0.21 -0.18 -0.21	Fe	0.35	0.87**	0.08	0.81^{**}	0.53*	-0.49	-0.32	-0.44	-0.38	-0.52*	-0.68**	0.68**	0.69 **	0.78**	000	-0.38	0.78**	0.57*	0.81^{**}	0.72**	0.71^{**}	
		0.27	-0.25	0.06	-0.18	-0.14	0.04	0.00	020	0.07	0.17	0.19	-0.09	0.23	0.02	0.40	0.01	-0.33	-0.07	-021	-0.18	-0.21	· ' I

Table 1. Correlation matrix for the estimated parameters (n=15)

	Component						
-	1	2	3	4	5	6	
pН	0.21	0.71	-0.18	0.04	0.51	0.24	
Salinity	0.83	0.09	-0.28	0.15	0.28	-0.16	
DO	-0.42	0.27	-0.16	0.55	0.51	-0.18	
Alkalinity	0.90	-0.24	0.06	0.15	0.20	0.02	
Nitrite	0.33	-0.02	0.21	-0.08	0.86	-0.01	
Nitrate	-0.45	0.57	0.38	-0.19	0.31	-0.18	
Ammonia	0.04	0.05	0.96	-0.15	0.02	0.01	
Phosphate	-0.22	0.87	0.40	-0.15	0.06	0.07	
Silicate	0.08	0.52	0.76	-0.18	-0.08	0.02	
TN	-0.32	0.82	0.46	0.01	-0.02	0.00	
ТР	-0.50	0.78	0.24	-0.23	-0.12	0.07	
Ca	0.43	0.00	-0.34	0.80	0.05	-0.01	
C _b	0.47	0.06	-0.18	0.29	0.47	0.47	
C _c	0.55	0.00	-0.24	0.75	0.16	0.12	
Pheophytin	-0.28	-0.31	-0.12	0.78	-0.19	0.28	
DIN	-0.04	0.15	0.95	-0.17	0.09	-0.03	
DON	-0.36	0.90	-0.09	0.13	-0.09	0.01	
DOP	-0.73	0.34	-0.11	-0.26	-0.34	0.04	
Mg	0.90	-0.21	0.08	0.07	0.13	-0.20	
Ca	0.90	-0.20	-0.06	-0.09	-0.19	0.14	
Na	0.97	-0.16	-0.07	0.07	0.07	-0.05	
K	0.97	-0.19	0.01	-0.04	0.01	-0.01	
Li	0.97	-0.16	0.09	-0.08	0.06	-0.01	
Sulphate	0.74	-0.15	-0.34	0.29	0.45	-0.01	
Iron	-0.19	0.11	0.04	0.10	-0.02	0.91	

Table 2. Factor loadings for the analysed parameters

Rotated Component Matrix, Extraction Method: Principal Component Analysis, Rotation Method: Varimax with Kaiser Normalization

had a profound influence. The observed negative correlations of calcium with DO suggest the uptake of calcium by primary producers (Cohen *et al.*, 1999). Sodium, potassium, lithium, calcium and sulphate contents varied seasonally while concentration of magnesium displayed both seasonal and spatial variation and iron exhibited a significant spatial variation only.

The effect of natural and anthropogenic flux was evaluated using factor analysis with Varimax rotation. A total of 92.77% of variance was explained by all the six factors. The first factor contributed 35.63% of the total variance (Table 2), and depicted strong positive loadings on all major elements, alkalinity, salinity and moderate loading on sulphate. It suggested the fact that intrusion of Sea water was the major parameter controlling the distribution of major elements. This factor showed negative loadings on nitrate TP and DOP with positive loadings on chlorophyll a, b and c. The variables in the second factor were characterised by strong positive loadings on phosphate, silicate, total nitrogen and DON, moderate loading on total phosphate and pH, accounting for 18.15% of the total variance. It also displayed a positive loading on nitrate. Release of phosphate from sediment during alkaline condition can be confirmed with this factor loading. This factor pointed towards the sediment remineralisation and digenetic processes occurring in the mangrove ecosystem.

Third factor contributed 14.24 % of the total variance exhibiting high positive loadings on ammonia and DIN, moderate loading on silicate, phosphate and total nitrogen. Faecal materials along with other anthropogenic and sewage wastes entering the mangrove ecosystem by land run off contributed towards this factor loading.

Fourth factor reflected 10.53 % of the total variance which had a loading on DO and strong loadings on chl a, chl c and pheophytin. This factor pointed towards the fact that primary production had a control over the DO concentration in the study area. Fifth and sixth factors accounted for 8.73% and 5.50% of the total variance respectively. Variables in component 5 consist of nitrite (strong), pH, DO, chlorophyll b and sulphate while that in component 6 consisted of iron (strong) and chl b. The effect of dissolved oxygen on nitrification

and sulphate reduction can be predicted in the ecosystem. This factor can also provide information about effect of pH on nitrification rates.

CONCLUSION

All the mangrove ecosystems in the study region were found to be nitrogen depleted during pre monsoon and monsoon which may be due to the slow regeneration of NO₃ compared to PO₄³⁻. Internal loading of phosphorous was evident from the correlation and factor analysis; hence phosphorous was not a limiting nutrient in the present investigation. Compared to other seasons, pre monsoon was characterised by higher chlorophyll content. From factor and correlation analysis it was evident that nutrients has no control over primary production. High turbulent water restricted the light attenuation resulting in lower primary production in post monsoon and monsoon season. The dissolved oxygen super saturation (132 %) was noticed at Kunjimangalam during pre monsoon which might be due to higher rate of photosynthesis and limited flushing. All major elements showed positive correlation with salinity indicating saline water influx from adjacent estuarine system. Factor analysis can also explain the processes like diagenesis, river influx and anthropogenic stress in the study region.

ACKNOWLEDGEMENT

The authors gratefully acknowledge the facilities and the support provided by the Dean and Director, School of Marine Sciences, Cochin University of Science and Technology. We also acknowledge the financial support rendered by Ministry of Earth Sciences (MoES) and Department of Science and Technology (DST), Government of India. The authors also wish to thank the reviewers for their comments and suggestions.

REFERENCE

Alongi, D. M. (1996). The dynamics of benthic nutrient pools and fluxes in tropical mangrove forests. Journal of Marine Research, **54**, 123-148.

Anderson, L. G., Turner, D. R., Wedborg, M. and Dyrssen, D. (1999). Determination of alkalinity, In: Methods of Sea Water Analyses, Grasshoff, K., Ehrhardt, M., Kremling K. (Eds.), Verlag Chemie, Weinheim, 127-147.

APHA, (1995). Standard methods for the examination of water and wastewater, Am. Public health association. Washington DC, **5**, 47-48.

APHA, (1999). Standard methods for examination of water and waste water. Am. Public health association. Washington DC.

Aston, S. R. (1980). Nutrients dissolved gasses and general biochemistry in estuaries. In: Chemistry and biogeochemistry of estuaries, E. Olausson & I. Cato (Eds.), New York: Wiley, 233-262.

Billen, G., Lancelot, C. and Meybeck, M. (1991). N, P, and Si Retention along the Aquatic Continuum from Land to Ocean, In: Ocean Margin Processes in Global Change, Mantoura, R. F. C, Martin, J.M. and Wollast, R. (Eds), New York, Wiley, 19- 44.

Boers, P. C. M. (1991). The influence of pH on phosphate release from lake sediments. Water Research, **25** (**3**), 309-311.

Bouillon, S., Koedam, N., Raman, A. V. and Dehairs, F. (2002). Primary producers sustaining macro-invertebrate communities in intertidal mangrove forests. Oecologia., **130**, 441–448.

Clemente, Z., Castro, V. L., Jonsson, C. M. and Fraceto, L. F. (2012). Ecotoxicology of Nano-TiO2 – An Evaluation of its Toxicity to Organisms of Aquatic Ecosystems. Int. J. Environ. Res., 6 (1), 33-50.

CMFRI, (2002). Central Marine Fisheries Research Institute, Technical extension series, Mar. Fish. Infor. Serv. T&E Ser., No. 172.

Cohen, M. C. L., Lara, R. J., Ramos, J. F., da F. and Dittmar T. (1999). Factors influencing the variability of Mg, Ca and K in waters of a mangrove creek in Braganc, a, North Brazil. Mangroves and Salt Marshes, **3**, 9-15.

FAO, (2004). Status and trends in mangrove area extent worldwide. Forest Resources Assessment Working Paper, **063**, 287.

Feng, X. Y., Luo, G. P., Li, C. F., Dai, L. and Lu, L. (2012). Dynamics of Ecosystem Service Value Caused by Land use Changes in Manas River of Xinjiang, China. Int. J. Environ. Res., **6** (2), 499-508.

FSI, (2003), Forest Survey of India, State of Forest Report, Dehra Dun, India.

Gonneea, M. E., Payton, A. and Herrera-Silveira, J. A. (2004). Tracing organic matter sources and carbon burial in mangrove sediments over the past 160 years. Estuarine, Coastal and Shelf Science, **61**, 211-217.

Granier, J., Billen, G. and Palfner, L. (2000). Understanding the oxygen budget and related ecological processes in the river Mosel: The Riverstrahler approach. Hydrobiologia, **410**, 151-166.

Grasshoff, K., Ehrhardt, M. and Kremling, K. (1999). Methods of Seawater Analysis. Verlag Chemie, Weinheim.

Hallare, A. V., Lasafin, K. J. A. and Magallanes, J. R. (2011). Shift in Phytoplankton Community Structure in a Tropical Marine Reserve Before and After a Major oil Spill Event. Int. J. Environ. Res., **5** (3), 651-660.

Hansen, H. P. and Koroleff, F. (1999). Determination of nutrients, In: Methods of Seawater Analyses, Grasshoff, K., Ehrhardt, M., Kremling, K. (Eds.), Verlag Chemie,Weinheim, 159-202.

Hansen, H. P. (1999). Determination of oxygen, In: Methods of Seawater Analyses, Grasshoff, K., Ehrhardt, M., Kremling, K. (Eds.), Verlag Chemie, Weinheim, 75-90.

Harrison, P. J., Khan, N., Yin, K., Saleem, M., Bano, N., Nisa, M., Ahmed, S. I., Rizvi, N. and Azam, F. (1997).

Nutrient and phytoplankton dynamics in two mangrove tidal creeks of the Indus River delta, Pakistan. Mar. Ecol. Prog. Ser., **157**, 13–19.

Jagtap, T. G., Murthy, P and Komerpant, D. (2002). Mangrove ecosystems of India: major biotic constituents conservation and management, In: Wetland conservation and management, Hosetti. B.B (Ed.), Pointer publishers, Jaipur, India, 34-64.

Jiang, D. X., Li, Y., Li, J. and Wang, G. X. (2011). Prediction of the Aquatic Toxicity of Phenols to Tetrahymena Pyriformis from Molecular Descriptors. Int. J. Environ. Res., **5** (4), 923-938.

Jing, L. and Zhiyuan, R. (2011). Variations in Ecosystem Service Value in Response to Land use Changes in the Loess Plateau in Northern Shaanxi Province, China. Int. J. Environ. Res., **5** (1), 109-118.

Kazungu, J., Dehairs, F. and Goeyens, L. (1993). Distribution of particulate and organic material in Gazi Bay. In: Dynamics and Assessment of Kenyan Mangrove Ecosystems, Commission of the European Communities, Woitchik, A.N. (Ed.), Project TS2-0240-C (GDF), Final Report, 175–193.

Kiteresi, L. I., Okuku, E. O., Mwangi, S. N., Ohowa, B., Wanjeri, V.O., Okumu, S. and Mkono, M. (2012). The Influence of Land Based Activities on the Phytoplankton Communities of Shimoni-Vanga system, Kenya. Int. J. Environ. Res., **6** (1), 151-162.

Kurian, C. V. (1994). Fauna of the mangrove swamps in Cochin estuary. Proceedings of the Asian Symposium on the Mangrove Environment. University of Malaya, Kuala Lumpar, Malaysia, 226–230.

Lee, S. Y. (1990). Primary productivity and particulate organic matter flow in an estuarine mangrove-wetland in Hong Kong. Marine Biology, **106**, 453–463.

Manson, F. J., Loneragan, N. R., Harch, B. D., Skilleter, G. A. and Williams, L. (2005). A broad-scale analysis of links between coastal fisheries production and mangrove extent: a case-study for northeastern Australia. Fish. Res. **74**, 69–85.

Moncheva, S., Gotsis-Skretas, O., Pagou, K. and Kraster A. (2001). Phytoplankton blooms in Black Sea and Mediterranean coastal ecosystems subjected to anthropogenic eutrophication: similarities and differences. Estuarine, Coastal and Shelf Science, **53**, 281-295.

Muller, T. J. (1999). Determination of salinity, In: Methods of SeaWater Analyses, Grasshoff, K., Ehrhardt, M., Kremling, K. (Eds.), Verlag Chemie, Weinheim, 41-74.

Naskar, K. and Mandal. R. (1999). Ecology and biodiversity of Indian mangroves Part I Global status. Daya publishing House, New Delhi.

Odum, E. P. (2000). Tidal marshes as outwelling/pulsing systems. In: Concepts and controversies in tidal marsh ecology. Weinstein, M. P, Kreeger, D. A (eds). Kluwer, Dordrecht, The Netherlands.

Pei, H., Liu, Q., Hu, W. and Xie, J. (2011). Phytoplankton Community and the Relationship with the Environment in Nansi Lake, China. Int. J. Environ. Res., **5** (1), 167-176. Qasim, S. Z., Wafar, M. V. M. (1990). Marine resources in the tropics. Resource Management and Optimization, **7**, 141-169.

Radhakrishnan, C., Gopi, K. C. and Palot, M. J. (2006).Mangroves and their faunal associates in Kerala. Occasional paper No 246. Records of Zoological Survey of India. Zoological Survey of India, Calicut, Kerala, 81pp.

Ramakrishnan, R., Perumal, P. and Santhanam P. (1999). Spatio-temporal variations of hydrographical features in the Pichavaram mangroves and Mohi aqua farm, Southeast coast of India. In: Proc. Intl. Sem. Appl. Hydrogeochem., Annamalai University, Annamalai Nagar, India, Published by Dept. of Geology, Annamalai University, Chidambaram, Tamil Nadu, 197-203.

Siddiqui, Z. (2011). Holistic Approach to Mitigate the Pollution Impacts in the Coastal Ecosystem of Thailand Using the Remote Sensing Techniques. Int. J. Environ. Res., **5** (2), 297-306.

Smith, K.A., Ball, T., Conen, F., Dobbie, K., Massheder, J. and Rey, A. (2003). Exchange of greenhou se gases between soil and atmosphere: interactions of soil physical factors and biological processes. European Journal of Soil Science, **54**, 779-91.

Smith, T. J., Boto, K. G., Frusher, S. D. and Giddins, R. L. (1991). Keystone species and mangrove forest dynamics: The influence of burrowing by crabs on soil nutrient status and forest productivity. Estuarine, Coastal and Shelf Science, **33**, 419 - 432.

Solanki, V. R., Hussain, M. M. and Raja, S. S. (2010). Water quality assessment of Lake Pandu Bodhan, Andhra Pradesh State, India. Environ. Monit. Assess., **163**, 411-419.

Takahashi, M. K. S. and Nakamoti, N. (1972). Chlorophyll distribution and photosynthetic activity in the North and Equatorial Pacific Ocean along 155°W. J. Oceanogr. Soc. Japan, **28**, 27-34.

Tripathy, S. C., Ray, A. K., Patra, S. and Sarma, V. V, J. (2005). Water quality assessment of Gautami-Godavari mangrove estuarine ecosystem of Andhra Pradesh, India during September 2001. Earth Syst. Sci, **114(2)**, 185-190.

Trott, L. A. and Alongi, D. M. (1999). Variability in surface water chemistry and phytoplankton biomass in two tropical, tidally dominated mangrove creeks. Mar. Freshwat. Res., **50**, 451-457.

Valentina, G. C., and Joseph, N. B. (2005). Spatial patterning of water quality in Biscayne Bay, Florida as a function of land use and water management. Marine Pollution Bulletin, **50**, 1416-1429.

White, P. A., Kalff, J., Rasmussen, J. B. and Gasol, J. M. (1991). The effect of temperature and algal biomass on bacterial production and specific growth rate in freshwater and marine habitats. Microb. Ecol., **21**, 99-118.

Zhao, S. J., Jiao, N. Z., Shen, Z. L. and Wu, Y. L. (2005). Causes and consequences of changes in nutrient structure in the Jiaozhou Bay. J. Integra. Plant Biol., **47** (**4**), 396–410.