

ORIGINAL ARTICLE

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

How to Cite This Article: Tavirani Rezaei M, Okhovatian F, Zamanian Azodi M, Rezaei Tavirani M.
Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping. Iran J Child Neurol. Autumn 2017;11(4):7-14

Mostafa REZAEI TAVIRANI MD¹,
Farshad OkHOVATIAN MD²,
Mona ZAMANIAN AZODI MD¹,
Majid REZAEI TAVIRANI MD³

1. Department of Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Department of Physiotherapy Research Centre, School of Rehabilitation, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

3. Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran.

Corresponding Author:
Tavirani Rezaei M. MD
Department of Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
Email: tavirany@yahoo.com

Revised: 4 -Dec- 2016
Last Revised: 01 -Feb -2017
Accepted: 04- Feb-2017

Abstract**Objective**

Duchenne muscular dystrophy (DMD) is one of the mortal diseases, subjected to study in terms of molecular investigation. In this study, the protein interaction map of this muscle-wasting condition was generated to gain a better knowledge of interactome profile of DMD.

Materials & Methods

Applying Cytoscape and String Database, the protein-protein interaction network was constructed and the gene ontology of the constructed network was analyzed for biological process, molecular function, and cellular component annotations.

Results

Among 100 proteins related to DMD, dystrophin, utrophin, caveolin 3, and myogenic differentiation 1 play key roles in DMD network. In addition, the gene ontology analysis showed that regulation processes, kinase activity, and sarcoplasmic reticulum were the highlighted biological processes, molecular function, and cell component enrichments respectively for the proteins related to DMD.

Conclusion

The central proteins and the enriched ontologies can be suggested as possible prominent agents in DMD; however, the validation studies may be required.

Keywords: Duchenne muscular dystrophy (DMD); Protein-protein interaction network; Gene ontology; Hub-bottlenecks; Biomarker panel

Introduction

Duchenne muscular dystrophy (DMD) is a severe muscle degradation condition that consequent in death of the patient. This inheritable disease is common in males with the prevalence of 1 out of 3500 (1). A functional neural deficit in this disease is reported. However, there are some ethical problems for investigation on the sick boys; EEG abnormality is reported for the number of the patients (2). There is not yet a suitable treatment for DMD (3).

There are some molecular studies for DMD (4, 5). One of the main extensively changed expressed molecules in this disease is known as dystrophin protein that the lack of its gene expression is linked to the disease development (6). This protein

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

can cause vast amount of changes in morphology and function of skeletal muscle tissue (5). Therefore, it accounts as putative biomarker for DMD. Beside important role of dystrophin, there are other introduced proteins for DMD, mostly conducted by proteomic approaches (7, 8).

Understanding other related proteins can help for better understanding and managing the disease. In addition to the role of these proteins, the interaction pattern of them can add more information about their contribution in DMD (9). Consequently, protein interaction networks provide information related to systematic communication scale of different kinds of diseases. Proteins that are dysfunctional in a specific disease condition are in complex interactions with each other. Dysregulation of each of these components can induce modification of the interaction of the whole interactome and changes the functions of other proteins. These disruptions of the whole system are responsible for the abnormal phenotype such as diseases (10). Some modification can be referred to the proteins that are essential for an interacted system (11). These proteins are known as central components that their dysfunctions lead to massive changes in the interaction profile (10). Therefore, detecting topological features can be prominent for disease analysis in terms of molecular origin (9). Here for obtaining a better knowledge about molecular mechanisms of DMD, the protein interactome and gene ontology analysis of disease was conducted using Cytoscape and DAVID software, respectively.

Materials and Methods

Determination of related proteins to a certain disease can be handled through different sources. One of the important sources is Cytoscape 3.4. This common software is free and is compatible with different sources. Many applications are available via Cytoscape and can provide further information for the mapped network. One of the famous interaction sources is String Database (DB). It is accessible through Cytoscape software.

Cytoscape can provide different interaction data from different sources. One of them, as mentioned earlier, is the String DB (<http://string-db.org/>) (12, 13). It has three options for retrieving interaction data including PubMed, disease, and proteins queries.

Protein interaction map was constructed according to String Database criteria. Here, the disease query was used for extracting 100 top related proteins to DMD. The used cut off for interaction evidence was set at 0.5. This score is a combined score from different sources. The disease score shows the relation of the disease and the obtained proteins. String DB provides interaction information from different sources. Furthermore, for network centrality analysis, Network Analyzer can be used. Here, the examined centrality parameters are degree and betweenness centrality. The nodes with highest degree (connections) are known as hub nodes and similarly, the nodes based on a function of larger number of shortest paths passing through them are called bottlenecks (14). These elements are crucial for the disease onset and progress (15).

Gene ontology analysis of the whole components of the network was done by the application of The Database for Annotation, Visualization and Integrated Discovery (DAVID) Bioinformatics 6.8 (<http://david.niaid.nih.gov>). The ontology analysis was based on biological process (BP), cellular component (CC), and molecular function (MF). The associated proteins were clustered for each of these annotations. In each cluster, similar enrichments are included (16). DAVID uses kappa statistics to group the terms.

Results

The protein network was constructed from 100 nodes including 18 isolated nodes and 1 main connected component. The network (main connected component) contains 82 nodes with different disease score relations and 395 edges. The highest disease combined score [4] was assigned for dystrophin protein. This protein also showed the highest value in degree (Figure 1).

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping




Fig 1. The protein-protein interaction mapping of DMD after analysis with network analyzer, as the nodes get bigger, the degree increases. The network (main connected component) consists of 82 nodes and 395 edges.

The centrality analysis of the network components based on selection of top 5% of the hub nodes and cut off 0.05 for betweenness revealed that 4 proteins including dystrophin, utrophin, caveolin-3 and myogenic

differentiation-1 are hub-bottleneck proteins (Table 1). Since connection between the determined crucial nodes provides informative data, these findings are tabulated in Table 2.

Table 1. The list of hub-bottleneck nodes of DMD PPI network based on degree and betweenness centrality analysis

Gene name	Description	Degree	BC	Disease score
DMD	Dystrophin	40	0.4	4
UTRN	Utrophin	28	0.1	3.9
CAV3	Caveolin 3	19	0.05	2
MYOD1	Myogenic differentiation 1	19	0.1	2.3

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

Table 2. Interactions properties between the determined hub-bottleneck nodes. The combined score was set to 0.5.
There is no interaction between CAV3 and MYOD1 (16)

Interactions	Combined Score
CAV3 (pp) DMD	0.992
DMD (pp) UTRN	0.964
MYOD1 (pp) DMD	0.815
CAV3 (pp) UTRN	0.699
MYOD1 (pp)UTRN	0.724

DAVID Bioinformatics Resources 6.8 (16), applied for functional annotation of the all contributed proteins in DMD network including biological process (BP), cell component (CC), and molecular function (MF) (Table 3,4). Each of abbreviations refers to the certain operation

for example “RT” (related term) refers to the terms related to the main introduced term in the same row. The accepted P-value is 0.1 and the terms are sorted base on P-value amounts. Enrichment score corresponds to the important role of the introduced terms in the cluster.

Table 3. Clustering annotation of the contributing elements in DMD network based on BP analysis. *Benjamini-Hochberg* is the correction method for P value

Annotation Cluster 1	Enrichment Score: 3.7	G	Count	P_Value	Benjamini
GOTERM_BP_DIRECT	muscle cell fate commitment	RT	4	1.5E-6	4.3E-4
GOTERM_BP_DIRECT	skeletal muscle tissue development	RT	6	3.1E-6	6.6E-4
GOTERM_BP_DIRECT	positive regulation of skeletal muscle fiber development	RT	4	4.2E-6	5.9E-4
GOTERM_BP_DIRECT	positive regulation of myoblast fusion	RT	4	4.9E-5	4.7E-3
GOTERM_BP_DIRECT	positive regulation of myoblast differentiation	RT	4	1.1E-4	9.3E-3
GOTERM_BP_DIRECT	positive regulation of muscle cell differentiation	RT	4	1.4E-4	1.1E-2
GOTERM_BP_DIRECT	skeletal muscle cell differentiation	RT	4	1.3E-3	7.0E-2
GOTERM_BP_DIRECT	regulation of transcription from RNA polymerase II promoter	RT	6	3.9E-2	6.5E-1
GOTERM_BP_DIRECT	transcription from RNA polymerase II promoter	RT	3	6.5E-1	1.0E0
Annotation Cluster 2	Enrichment Score: 1.51	G	Count	P_Value	Benjamini
GOTERM_BP_DIRECT	response to hypoxia	RT	5	9.8E-3	3.2E-1
GOTERM_BP_DIRECT	positive regulation of transcription, DNA-templated	RT	7	2.3E-2	4.9E-1
GOTERM_BP_DIRECT	aging	RT	4	5.7E-2	7.3E-1
GOTERM_BP_DIRECT	female pregnancy	RT	3	6.9E-2	7.6E-1

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

Table 4. Clustering annotation of the contributing elements in DMD network based on CC analysis. *Benjamini-Hochberg* is the correction method for P-value

Annotation Cluster 1		Enrichment Score: 3.83		G	Count	P_Value	Benjamini
GOTERM_CC_DIRECT	sarcoplasmic reticulum membrane	RT		5	9.9E-6	2.4E-4	
GOTERM_CC_DIRECT	T-tubule	RT		5	2.6E-5	5.6E-4	
GOTERM_CC_DIRECT	sarcoplasmic reticulum	RT		4	4.4E-4	7.4E-3	
GOTERM_CC_DIRECT	I band	RT		3	4.0E-3	4.8E-2	
Annotation Cluster 2		Enrichment Score: 1.54		G	Count	P_Value	Benjamini
GOTERM_CC_DIRECT	costamere	RT		3	2.7E-3	3.5E-2	
GOTERM_CC_DIRECT	focal adhesion	RT		6	2.3E-2	2.0E-1	
GOTERM_CC_DIRECT	cell-cell adherens junction	RT		3	3.9E-1	8.5E-1	

Table 5. Clustering annotation of the contributing elements in DMD network based on MF analysis. *Benjamini-Hochberg* is the correction method for P-value.

Annotation Cluster 1		Enrichment Score: 3.28		G	Count	P_Value	Benjamini
GOTERM_MF_DIRECT	creatine kinase activity	RT		4	1.4E-6	3.0E-4	
GOTERM_MF_DIRECT	kinase activity	RT		7	5.7E-4	1.5E-2	
GOTERM_MF_DIRECT	ATP binding	RT		10	1.8E-1	7.9E-1	
Annotation Cluster 2		Enrichment Score: 2.38		G	Count	P_Value	Benjamini
GOTERM_MF_DIRECT	transcription factor activity, RNA polymerase II distal enhancer sequence-specific binding	RT		5	1.5E-4	6.3E-3	
GOTERM_MF_DIRECT	E-box binding	RT		4	3.9E-4	1.3E-2	
GOTERM_MF_DIRECT	transcriptional activator activity, RNA polymerase II transcription regulatory region sequence-specific binding	RT		4	7.6E-3	1.1E-1	
GOTERM_MF_DIRECT	protein heterodimerization activity	RT		7	1.4E-2	1.5E-1	
GOTERM_MF_DIRECT	RNA polymerase II core promoter proximal region sequence-specific DNA binding	RT		4	1.9E-1	8.0E-1	

Discussion

Protein interaction mapping as a powerful analytical method is attracted a great attention of scientists in the field of molecular analysis of diseases (17). One of the recently introduced disciplines is network medicine that investigates about network topological features of a specific disease (18). The information which introduces numbers of essential proteins in terms of interactions can be crucial in disease progression (19). These proteins can be considered as biomarkers that by validation tests would be introduced for clinical approaches.

The aim of this study was to provide a preliminary knowledge of these key components for more in depth

comprehension of Duchenne muscular dystrophy related mechanisms and further investigation in this field. Consequently, a network of all 100 top reported proteins for this disease was constructed and analyzed based on centrality and gene ontology. The associated proteins for this disease were obtained from String source and this source assigned disease score relation based on differently combined evidence. Only 82 genes included in the network and the numbers of 18 genes remained isolated. Each node is connected to the about 5 neighbor nodes. However, due to heterogeneity of the network, the number of the edges for the nodes is different. The centrality analysis of the illustrated network differentiated

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

the nodes and the important genes were highlighted. The important proteins are visible as the largest nodes in the network. Network Analyzer is the applied plug-in for this analysis, provide quantitative information. We choose the most ranked proteins according to the highest degree and largest betweenness centrality. The proteins that showed both characteristics are listed in Table 1. They are known as hub-bottleneck proteins (18). Hub implies on high degree and equally, bottlenecks denote large BC values (20). Therefore, these four proteins including dystrophin, utrophin, caveolin 3, and myogenic differentiation-1 are the key agents of the DMD network. Dystrophin as the putative biomarker of DMD (21), also express high centrality values in our network. Attributing top score to dystrophin in our research is a potent validation for the constructed and analyzed PPI network related to DMD. This protein is the top rated node for the integrity of the network. About 10% of all edges are linked to this protein connected to the 50% of all nodes of the network. Dystrophin is characterized by top scores of degree; betweenness and disease score (Table 1). Other key proteins are also important and their role needs more evaluation in the disease. In fact, their involvement with different expression changes has been reported extensively for DMD (3, 22, 23).

Utrophin the second key protein in our analysis possess near scores to dystrophin in compare to the other two proteins. Its disease score is equal to Utrophin so it is related to DMD strongly. For more resolution, mutual interaction between the crucial proteins as a useful tool was done. In Table 2, the linkage of theses essential proteins together is tabulated. The interactions scores show that the proteins have high degree of interactions between each other. However, only CAV3 and MYOD1 have no connections. The introduced central proteins construct an integrated network related to DMD. Combined score value for the paired genes is higher if DMD is paired. On the other hand, MYOD1 reduces this score. This finding confirms the order of the sorted genes.

Moreover, to identify the role of these linked proteins to the disease, the gene ontology analysis was used. All the extracted proteins [82] were under consideration of this purpose as they show some association to the disease. Accordingly, DAVID Bioinformatics clustered these

agents with the respect to query annotations. The ontology aspects include BP, CC, and MF. The first implemented ontology is biological process of the DMD relevant proteins. In each enrichment analysis, two important clusters are introduced. The first one is the significant cluster and the related terms in this cluster may play more roles that are important in DMD. The main terms in cluster 1 are the biological processes are most active in regulatory processes. Muscle cell fate commitment is the most significant term with the contribution of 4 of the proteins in the network. Skeletal muscle tissue development, positive regulation of skeletal muscle fiber development, positive regulation of myoblast fusion, positive regulation of myoblast differentiation, positive regulation of muscle cell differentiation, skeletal muscle cell differentiation is the other terms related to the involved proteins in DMD. In cell component analysis, the sarcoplasmic reticulum is the proposed resident for the most of the DMD proteins, in which sarcoplasmic reticulum membrane is the most significant part. T tubule and I band are the other two components determined as related parts to the DMD. DAVID functional annotation for MF in Table 5 showed that kinase activity is the highlighted function of the DMD network and creatine kinase activity is the most highlighted one. ATP binding is the other molecular function attributed to the analyzed proteins of DMD PPI network. Investigation about roles of dystrophin (24), utrophin (25), caveolin-3 (26), and myogenic differentiation-1 (27) indicates that relationship between cluster-1 and DMD is highlighted; terms of cluster-1 play roles that are more important in DMD relative to cluster-2.

In Conclusion, dystrophin, utrophin, caveolin 3, and myogenic differentiation-1 are introduced as biomarker panel for DMD onset and development. However, the first two proteins play crucial roles. More investigation in the field can lead to validation of the elements of this panel for diagnostic and therapeutic proposes.

Acknowledgement

This project was supported by Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Author's Contribution

Dr. Mona Zamanian-Azodi and Dr. Mostafa Rezaei-Tavirani provided manuscript and edited revised version

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

of paper. All authors agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Conflict of Interest

The authors declare that there is no conflict of interests.

References

1. Nguyen-Tran D-H, Hait NC, Sperber H, et al. Molecular mechanism of sphingosine-1-phosphate action in Duchenne muscular dystrophy. *Dis Model Mech* 2014; 7: 41-54.
2. Anderson J, Head S, Rae C, Morley J. Brain function in Duchenne muscular dystrophy. *Brain* 2002; 125: 4-13.
3. Hirst RC, McCullagh KJ, Davies KE. Utrophin upregulation in Duchenne muscular dystrophy. *Acta Myol* 2005; 24: 209-216.
4. Coenen-Stass AM, McClorey G, Manzano R et al. Identification of novel therapy-responsive protein biomarkers for Duchenne muscular dystrophy by aptamer-based serum proteomics. *Sci Rep* 2016; 6:37047.
5. Fröhlich T, Kemter E, Flenkenthaler F, et al. Progressive muscle proteome changes in a clinically relevant pig model of Duchenne muscular dystrophy. *Sci Rep* 2016; 6:33362.
6. Li HL, Fujimoto N, Sasakawa N, et al. Precise correction of the dystrophin gene in duchenne muscular dystrophy patient induced pluripotent stem cells by TALEN and CRISPR-Cas9. *Stem Cell Reports* 2015; 4: 143-154.
7. Rouillon J, Zocevic A, Leger T, et al. Proteomics profiling of urine reveals specific titin fragments as biomarkers of Duchenne muscular dystrophy. *Neuromuscul Disord* 2014; 24: 563-573.
8. Holland A, Dowling P, Zweyer M et al. Proteomic profiling of cardiomyopathic tissue from the aged mdx model of Duchenne muscular dystrophy reveals a drastic decrease in laminin, nidogen and annexin. *Proteomics* 2013; 13: 2312-23.
9. Zamanian-Azodi M, Rezaei-Tavirani M, Rahmati-Rad S, Hasanzadeh H, Tavirani MR, Seyyedi SS. Protein-Protein Interaction Network could reveal the relationship between the breast and colon cancer. *Gastroenterol Hepatol Bed Bench* 2015; 8: 215-24.
10. Rezaei-Tavirani M, Zamanian-Azodi M, Rajabi S, Masoudi-Nejad A, Rostami-Nejad M, Rahmatirad S. Protein Clustering and Interactome Analysis in Parkinson and Alzheimer's Diseases. *Arch Iran Med* 2016: 19.
11. Zamanian-Azodi M, Mortazavi-Tabatabaei SA, Mansouri V, Vafae R. Metabolite-protein interaction (MPI) network analysis of obsessive-compulsive disorder (OCD) from reported metabolites. *Arvand J Health Med Sci* 2016.
12. Lopes CT, Franz M, Kazi F, Donaldson SL, Morris Q, Bader GD. Cytoscape Web: an interactive web-based network browser. *Bioinformatics* 2010; 26: 2347-2348.
13. Szklarczyk D, Morris JH, Cook H, et al. The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res* 2016; gkw937.
14. Safari Alighiarloo N, Taghizadeh M, Tabatabaei SM, et al. Identification of new key genes for type 1 diabetes through construction and analysis of the protein protein interaction networks based on blood and pancreatic islet transcriptomes. *J Diabetes* 2017 9(8):764-777.
15. Safaei A, Tavirani MR, Oskouei AA, Azodi MZ, Mohebbi SR. Evaluation of protein clustering of pancreatic cancer. *Arvand J Health Med Sci* 2016;1(2):68-77.
16. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 2009; 4: 44-57.
17. Safari-Alighiarloo N, Taghizadeh M, Rezaei-Tavirani M, Goliaei B, Peyvandi AA. Protein-protein interaction networks (PPI) and complex diseases. *Gastroenterol Hepatol Bed Bench* 2014;7(1):17-31.
18. Azodi MZ, Peyvandi H, Rostami-Nejad M, et

Duchenne Muscular Dystrophy (DMD) Protein-Protein Interaction Mapping

- al. Protein-protein interaction network of celiac disease. *Gastroenterol Hepatol Bed Bench* 2016; 9: 268-277.
19. Abbaszadeh H-A, Peyvandi AA, Sadeghi Y et al. Er: YAG Laser and Cyclosporin A Effect on Cell Cycle Regulation of Human Gingival Fibroblast Cells. *J Lasers Med Sci* 2017; 8: 143-9.
20. Safaei A, Tavirani MR, Oskouei AA, Azodi MZ, Mohebbi SR, Nikzamir AR. Protein-protein interaction network analysis of cirrhosis liver disease. *Gastroenterol Hepatol Bed Bench* 2016; 9: 114.
21. Mendell JR, Campbell K, Rodino-Klapac L et al. Dystrophin immunity in Duchenne's muscular dystrophy. *N Engl J Med* 2010; 363: 1429-1437.
22. Deconinck AE, Rafael JA, Skinner JA, et al. Utrophin-dystrophin-deficient mice as a model for Duchenne muscular dystrophy. *Cell* 1997; 90: 717-727.
23. Galbiati F, Razani B, Lisanti MP. Caveolae and caveolin-3 in muscular dystrophy. *Trends Mol Med* 2001; 7: 435-441.
24. Blake DJ, Weir A, Newey SE, Davies KE. Function and genetics of dystrophin and dystrophin-related proteins in muscle. *Physiol Rev* 2002; 82: 291-329.
25. Haenggi T, Fritschy J-M. Role of dystrophin and utrophin for assembly and function of the dystrophin glycoprotein complex in non-muscle tissue. *Cell Mol Life Sci* 2006; 63: 1614-1631.
26. Matsuda C, Hayashi YK, Ogawa M, et al. The sarcolemmal proteins dysferlin and caveolin-3 interact in skeletal muscle. *Hum Molecular Gen* 2001; 10: 1761-1766.
27. Blais A, Tsikitis M, Acosta-Alvear D, Sharan R, Kluger Y, Dynlacht BD. An initial blueprint for myogenic differentiation. *Genes Dev* 2005; 19: 553-569.