

ORIGINAL ARTICLE

Incidence and Etiology of Stroke among Hospitalized Children: A Case-Series Study

How to Cite This Article: Keihani Doust ZT¹, Noori Sari F¹, Akbari Asbagh P¹, Farahani Z¹, Tehrani F¹, Shariat M¹. Incidence and Etiology of Stroke among Hospitalized Children: A Case-Series Study. *Iran J Child Neurol*. Winter 2019; 13(1): 65-70

Zarin Taj KEIHANIDOUST MD¹,
Fateme NOORI SARI MD¹,
Parvin AKBARI ASBAGH MD¹,
Zahra FARAHANI MD¹,
Fateme TEHRANI MD¹,
Mamak SHARIAT MD¹

1. Maternal & Child Health Specialist,
Maternal-Fetal Neonatal Research
Center, Tehran University of Medical
Sciences, Tehran, Iran

Corresponding Author

Shariat M. MD

Maternal & Child Health Specialist,
Maternal-Fetal Neonatal Research
Center, Tehran University of Medical
Sciences, Tehran, Iran

Email: shariat_soufi@yahoo.com

Received: 08-Oct-2017

Last Revised: 30-Dec-2017

Accepted: 15-Jan-2018

Abstract

Objectives

Stroke is a sudden blockage or rupture of brain vessels resulting neural defect or impairment. We aimed to investigate the incidence and causes of stroke in hospitalized children (Tehran-Iran, 2008-2013).

Materials & Methods

This case series study was carried out in pediatric ward of tertiary care Vali-Asr Hospital, Imam Khomeini Hospitals Complex (Tehran-Iran) from 2008 to 2013. One month to 15 yr old admitted children due to stroke were enrolled into this case series study. Diagnosis was confirmed with brain imaging. Participants' demographic data, potential risk factors and neuroimaging findings were obtained from Hospital Reporting System. Recorded data were studied and considered regarding the incidence of stroke and its causes. Indeed we investigated cardiological causes as well as different items related to hematological disorders.

Results

Of 20000 admitted subjects in Imam Hospital during 5 yr, stroke was diagnosed in 15 cases. The incidence among the population study was 0.75 per 100000 children. Stroke was more frequent in males than females (). The most common age of stroke was 4-6 yr and the mean age of stroke was 58.8 months equal to 4.9 year. The most frequent stroke was hemorrhagic stroke (26%), followed by vascular (20%) and coagulopathy disorders (20%).

Conclusion

The incidence of stroke in children was 0.75 per 100000. Hemorrhagic stroke due to major trauma, coagulopathy and vasculopathy were observed as most frequent causes that necessitate implementing some strategies for prevention, earlier diagnosis, and treatment.

Keywords: Stroke; Infant; Adult children; Cerebral blood flow

Introduction

Stroke is rapidly declined of cerebral blood flow resulting impairment of brain function (1). The incidence of childhood stroke is about 13-16 in 100000 (2). In the United States, stroke with mortality rate of 0.34 deaths per 100000 is one of top 10 causes responsible for death among children (3).

The causes of stroke in children are different from adults. Thrombophilia, sickle cell anemia, infections, acquired or congenital emboligenic heart diseases were reported as important causes. Stroke in children may lead to death, cognitive-motor impairment and long-lasting seizure (24 h or longer) (2).

Congenital heart disease is reported as a profound risk factor for ischemic and hemorrhagic stroke (4). The rate of ischemic stroke in children with CHD is about 10-fold higher than age and sex-matched children in the control group (5). Moreover, the frequency of vaso-occlusive stroke may increase during cardiac surgery based on child's age, duration of bypass or reoperation (6). Coagulopathy and thrombophilia in combination with other factors are reported a potential risk factor for arterial ischemic stroke. Genetic factors like Moyamoya Syndrome and Sickle cell disease are also responsible for 6% and approximately 1% of overt stroke, respectively (7). Metabolic diseases and trauma resulting infarction are uncommon; however, these conditions are important for the pediatric population (1).

Stroke is known as an important cause of mortality (18%) or lifelong disability in two-thirds of the children (1-3). We aimed to evaluate the incidence and main underlying causes of stroke which would be of value in implementing strategies to decrease morbidity and mortality rate as well as economic burden on families and society.

Materials & Methods

This case series study was carried out in Pediatric Ward of tertiary care Vali-Asr Hospital, Imam Khomeini Hospitals Complex (Tehran- Iran) from 2008 to 2013. All admitted patients due to stroke; aged one month to 15 yr old entered the study. Inclusion criteria were presence focal acute neural impairment confirmed by brain imaging. Neonatal strokes were excluded due to variable etiologic patterns. Participants' demographic data and potential risk factors including gender, age, clinical presentation and neuroimaging findings were obtained from Hospital Reporting System. Underlying disease and causes (prenatal problems, congenital heart disease, hematologic disorders, infection of CNS and trauma) were also collected and studied. Recorded data were considered regarding frequency of stroke and the causes. Frequency was presented as mean \pm SD.

This manuscript was taken from a thesis with ID 22664. Our study was approved by the institutional Review Board of Tehran University of Medical Sciences according to Helsinki declaration. The gathered data were confidential.

Results

Of 20000 admitted children in Vali-Asr Hospital during 5 yr led study, stroke was diagnosed in 15 cases. The incidence rate among the population study was 0.75 per 100000 children. Stroke was more frequent in males than females; male/female ratio was 1.14/1. The incidence of stroke in males and females were 0.40 and 0.35 per 100000, respectively. The most common age of stroke was 4-6 yr and the mean age of stroke was 58.8 months equal to 4.9 yr. The most frequent cause of stroke was hemorrhagic stroke due to trauma, followed by vascular and coagulopathy disorders. Of 4

cases with hemorrhagic stroke, stroke in 3 of them was due to child sexual abuse and a case due to severe car accident. Demographic data and factors

associated with stroke in 15 patients are shown in Table 1.

Table 1. Demographic data and factors associated with stroke in 15 patients

| Variables | n (%) |
|--|----------|
| Sex | |
| Male | 8 (53) |
| female | 7 (47) |
| Age(yr) | |
| 1-3 | 6 (40) |
| 4-6 | 7 (46) |
| 7-9 | 0 |
| 10-12 | 1 (6.66) |
| 13-15 | 1 (6.66) |
| Causes | |
| - Congenital Cardiovascular | 1 (6.66) |
| -Vascular disease; (Artery-Vein malformation and hemangioma) | 3 (20) |
| -Hemorrhagic stroke | |
| -Coagulopathy; (Protein C,S deficiencies and thromboembolism) | 4 (26.6) |
| -Post surgery | 3 (20) |
| -Congenital metabolic disease; (Mitochondrial disease and lactic acidosis) | 1 (6.66) |
| -PROM* | 1 (6.66) |
| -Idiopathic | 1 (6.66) |
| | 1 (6.66) |
| | 1 (6.66) |

*premature rupture of the membranes

Table 2. Comparison of male/female ratio in different studies

| Variables | Present study | Report by Tavassoli et al. (14) | Report by Deng et al. (12) | Report by Shi et al. (13) | Report by Chung et al. (3) |
|-------------------|---------------|---------------------------------|----------------------------|---------------------------|----------------------------|
| Male | 53% | 60% | 60% | 57% | 56% |
| Female | 47% | 40% | 40% | 43% | 44% |
| Male/Female ratio | 1.14/1 | 1.5/1 | 1.5/1 | 1.32/1 | 1.27/1 |

Table 3. Comparison of causes in different studies

| Causes | Present study | Report by Tavassoli et al. (14) | Report by Shi et al. (13) | Report by Chung et al. (3) | Report by Simma et al. (11) |
|---------------------|---------------|---------------------------------|---------------------------|----------------------------|-----------------------------|
| Trauma | 26.6% | 5% | 11% | - | 0 |
| Vascular disease | 20% | 2.5% | 0 | 26% | 85% |
| Coagulopathy | 20% | 7.5% | 0 | 28% | 50% |
| Heart disease | 6.66% | 17.5% | 0 | 30% | 0 |
| Hb disorders | 0 | 0 | 0 | - | 0 |
| Metabolic disorders | 6.66% | 0 | 0 | - | 0 |
| Post surgery | 6.66% | 0 | 0 | - | 0 |
| Idiopathic | 6.66% | 40% | 32.5% | 12% | 13.6% |

Discussion

Although pediatric stroke is not common, it significantly effects on morbidity- mortality rate and quality of life (8, 9). This retrospective population-based study was conducted to determine the incidence and risk factors for pediatric stroke. Based on our results, the incidence of stroke was 0.75 per 100000 among the population study. Former study from Iran has shown a higher rate; the annual incidence of children stroke (<15 yr) in 2006 was reported 1.83/100000 (10). The reported incidence of pediatric stroke (<15 yr of age) in Hong Kong between 1998 and 2001 was higher than in our population (2.1 cases per 100000) (3).

The incidence of pediatric stroke (ischemic and hemorrhagic stroke) was reported 2.70/100000 child-yr in Austria, from 1984-2005 (11). The incidence of stroke varies in different countries based on some factors like the ethnic background, age-dependent references and the number of population study (8).

Our results also showed that stroke was more frequent among males than females. Inconsistent with our results, more than 60% of stroke was observed in male gender (12). In a study, of 157 children with arterial ischemic stroke; male/female ratio was 1.4:1 (13). This ratio was compared among different studies and is shown in Table 2.

The most common age of stroke in the present

study was 4 to 6 year, consistent with other a mean age of 5.5 yr among 17 cases with stroke (10). The most common age of stroke was 2 to 5 yr old among 40 Iranian children aged 3 months to 14 yr old (14). Of the 27 children aged between 6months and 15 yr with stroke, the average age was 6.91 (± 2.08) year (2). On the other hand, stroke was more common in children less than 1 yr (51.35%) (8).

Traumatic hemorrhagic stroke was the most frequent cause of stroke in our pediatric population (26.6%). This rate is inconsistent with another study where head trauma and arterial dissection reported in 25.7% of pediatric brain infarction (15). On the other hand, spontaneous intraparenchymal hemorrhage, non-traumatic subarachnoid hemorrhages, and arterial ischemic stroke are the most frequent causes of stroke with incidence 1.5–2.9 per 100.000 children (16).

Of 94 pediatric strokes, 14 strokes (28%) were hemorrhagic (3). Moreover, hemorrhagic strokes were reported in 20% of patients (14). Different etiologies from different studies are compared and shown in Table 3.

Another common etiology observed in our population was vascular disease by 20%. Vasculopathy was seen in 23.5% of their population study (10).

Coagulopathy disorders as frequent causes were responsible for 20% of strokes in our study. This frequency was higher in comparison with another study from Iran; prothrombotic disorders were showed in 7.5% of cases with stroke (14). However, there is a higher incidence regarding prothrombotic abnormality (50%) (11).

Although congenital heart disease is reported the main causes of childhood stroke (responsible for 15%-30%), this rate in our study was not notable (6.66%).

In conclusion, the incidence of stroke in children was 0.75 per 100000. Hemorrhagic stroke due to major trauma, coagulopathy and vasculopathy were the most frequent causes that necessitate implementing strategies for prevention, earlier diagnosis, and treatment.

Acknowledgement

This study was supported by Tehran University of medical sciences (TUMS). We acknowledge the parent for their participation in this study

Authors' contributions

Keihani Doust: Conception- Revising and Final approval

Noori Sari: Data gathering and Drafting the work

Akbari Asbagh: Revising and consultant

Farahani: Drafting the work

Tehrani: Interpretation of Data

Shariat: Designing and Data analysis

All authors agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Conflict of interest

The authors had no potential conflicts of interest during study.

References

1. Rosa M, De Lucia S, Rinaldi VE, Le Gal J, Desmarest M, Veropalumbo C, Romanello S, Titomanlio. Pediatric arterial ischemic stroke: acute management, recent advances and remaining issues. *Ital J Pediatr* 2015; 41:95.
2. Mallick AA, O'Callaghan FJ. The epidemiology of childhood stroke. *Eur J Paediatr Neurol* 2010; 14(3):197-205.
3. Chung B, Wong V. Pediatric stroke among

- Hong Kong Chinese subjects. *Pediatrics* 2004; 114(2):e206-12.
4. Fox CK, Sidney S, Fullerton HJ. Community-based case-control study of childhood stroke risk associated with congenital heart disease. *Stroke* 2015; 46(2):336-40.
 5. Mandalenakis Z, Rosengren A, Lappas G, Eriksson P, Hansson P, Dellborg M. Ischemic stroke in children and young adults with congenital heart disease. *J Am Heart Assoc* 2016; 5:e003071.
 6. Domi T, Edgell DS, McCrindle BW, Williams WG, Chan AK, MacGregor DL, Kirton A, deVeber GA. Frequency, predictors, and neurologic outcomes of vaso-occlusive strokes associated with cardiac surgery in children. *Pediatrics* 2008; 122(6):1292-8.
 7. DeVeber G. Arterial ischemic strokes in infants and children: an overview of current approaches. *Semin Thromb Hemost* 2003; 29:567-573.
 8. Eltayeb AA, Askar GA, Abu Faddan NH, Kamal TM. Prothrombotic risk factors and antithrombotic therapy in children with ischemic stroke. *Ther Adv Neurol Disord* 2015; 8(2):71-81.
 9. Inaloo S, Ghofrani M. Stroke in an infant; its association with antiphospholipid antibody and acquired protein C and S deficiencies. *Iran J Allergy Asthma Immunol* 2004; 3(2):93-5.
 10. Ghandehari K, Izadi-Mood Z. Incidence and Etiology of Pediatric Stroke in Southern Khorasan. *Advanced Research Yields Across Atherosclerosis (ARYA) Journal* 2007;3 (1): 29-33.
 11. Simma B, Martin G, Müller T, Huemer M. Risk factors for pediatric stroke: consequences for therapy and quality of life. *Pediatr Neurol* 2007;37(2):121-6.
 12. Deng Y, Wang Y, Yang W, Yu Y, Xu J, Wang Y, Gao B. Risk factors and imaging characteristics of childhood stroke in China. *J Child Neurol* 2015;30(3):339-43.
 13. Shi KL, Wang JJ, Li JW, Jiang LQ, Mix E, Fang F, Wu HS, Jin X, Jing H, Zou LP. Arterial ischemic stroke: experience in Chinese children. *Pediatr Neurol* 2008;38(3):186-90.
 14. Tavassoli A, Ghofrani M. Clinical Survey of Cerebrovascular Disease in Children. *Iran J Pediatr* 2008; 18(Suppl 1):53-58.
 15. Gulati S, Karla V. Stroke in children. *Indian J Pediatrics* 2003; 70:639-648.
 16. Ciccone S, Cappella M, Borgna-Pignatti C. Ischemic Stroke in infants and children: practical management in emergency. *Stroke Res Treat* 2011; 2011:736965.