Percutaneous Revascularization of Patients with History of Coronary Artery Bypass Grafting

Special thanks are due to Dr. Fatemeh Behboudi et al. for their invaluable work on percutaneous intervention on grafted veins as well as native coronary arteries in patients with previous history of coronary artery bypass graft (CABG) surgery.¹ They report 71 patients with a history of CABG, in whom percutaneous coronary intervention (PCI) was performed on native vessels in 60%, on grafted vessels in 32%, and on both in the remaining 8%.

In the modern era of cardiovascular medicine, CABG and PCI are not rivals but could be complementary to each other. With the increasing age of patients with a history of CABG, atherosclerotic changes progress in their native as well as grafted vessels. Within 10 years after CABG, nearly half of saphenous vein grafts (SVGs) fail or demonstrate significant atherosclerotic disease and the patients become symptomatic.² Atherosclerotic plaques in SVGs are always complex and friable and may be associated with thrombus.³

The first approach to the symptomatic stable post-CABG patient is optimizing medical treatment as well as assessing the amount of myocardium in jeopardy and localizing the ischemia by non-invasive tests such as imaging modalities. In case of medical treatment failure or high-risk non-invasive test results, or if the presentation of the patient is acute coronary syndrome, coronary angiography may be indicated and revascularization is on the table. Redo CABG could be an option, but there are some obstacles. Higher mortality and morbidity has been reported compared with first CABG, especially in subjects with advanced age and with comorbid states.⁴ Sternotomy could be a potential hazard for the grafted internal mammary artery. The second approach is PCI on either native coronary arteries or grafted vessels or both whenever feasible and is indicated based on the area of the ischemia. PCI is often the preferred treatment option in this population since reoperation imposes substantial risk on these subjects.

The study conducted by Dr. Behboudi and her colleagues is a report on in-hospital and mid-term outcome of PCI on patients with a previous history of CABG. The favorable outcome of the subjects in this study encourages the cardiologists to perform PCI on this group of patients. Whether the native vessel or the SVG is preferable for PCI is not answered in this survey. The target vessel for PCI in post-CABG patients is sometimes a matter of debate, and the selection of the native vessel or SVG with significant stenosis for intervention is not always a simple decision. A comparison of the major adverse cardiac events (MACE) rate, procedural complications, and outcome between the two groups can help solve this problem.

The percutaneous treatment of the SVG has been a matter of interest for many years, and there is a large body of data on this procedure. The SVG could be a target because of lesser tortuosity, less calcification, and larger diameter compared with native vessels. The mechanisms of stenosis in the SVG are somehow different from those in native coronary arteries.³ Native coronary artery and SVG atherosclerosis should be considered different diseases. Atherosclerotic plaques in the graft are more diffuse, friable, soft, and lipid-rich. These characteristics render SVG lesions prone to fragmentation and distal embolization during PCI.5 The incidence of no reflow and rate of periprocedural myocardial infarction are higher in SVG angioplasty, owing in large part to the embolization of the abundant and friable atherosclerotic debris in diseased SVGs.^{3, 6} The use of embolic protection devices (EPDs) has been demonstrated to reduce the major adverse cardiac events rate as well as no reflow in SVG angioplasty⁷ and these devices are recommended in guidelines whenever technically feasible.8 Despite these supporting data, EPDs are used only in 22% of patients in the United States.9

In this study, Dr. Behboudi et al. mention that PCI on post-CABG patients is feasible and safe. A challenge is that which vessel is better for intervention ,the native vessel or the SVG? Was there any difference in the MACE rate between the group who underwent PCI on their native coronaries and those with intervention on their SVGs? It seems that the sample size was not sufficient for such an analysis. Provision of information on the no-reflow rate in the SVG group and utilization rate of EPDs would be helpful. There is room for investigation about the procedural time, radiation dose, and amount of contrast injection in SVG intervention and a comparison with PCI on native vessels in post-CABG patients.

We will encounter more symptomatic post-CABG patients in the future. More PCI procedures will be carried out on these patients either on their native or their grafted vessels. More data are needed to help select the best target vessel to obtain maximal efficacy and minimal risk. Sometimes we have to open as much vessel as we can, especially when the patient remains symptomatic despite opening one target vessel.

The important role of optimal medical treatment should



Fatemeh Behboudi et al.

not be forgotten. Preventive efforts to halt the progression of atherosclerotic changes in grafted veins and native coronary arteries perhaps are the better way ;be that as it may ,the impact of risk factor modification and changing life style on this issue needs to be confirmed with larger scale clinical trials.

1411713138. Tel: + 98 21 88029702. Fax: + 98 21 88029731. E-mail: kianhoseini@tums.ac.ir.

Seyed Kianoosh Hoseini,

Assistant Professor of Cardiology, Tehran University of Medical Sciences, Tehran Heart Center, Karegar Ave, Tehran, Iran.



References

- Acholoider, Oakfill, Hashemi SR. Immediate results and sixmonth clinical outcome after percutaneous coronary intervention in patients with prior coronary artery bypass surgery. J Teh Univ Heart Ctr 2011;6:31-36.
- Goldman S, Zadian K, Moritz T, Ovitt T, Sethi G, Copeland JG. Long term patency of saphenous vein and internal mamary artery grafts after coronary artery bypass surgery; results from a department of veterans affairs cooperative study. J Am Coll Cardilo 2004;44:2149-2156.
- Kereiakes DJ, Turco MA, Breall J. A novel filter-based distal embolic protection device for percutaneous intervention of saphenous vein graft lesions. J Am Coll Cardiol 2008;1:248-257.
- Fitzgibbon GM, Kafka HP, Leach AJ. Coronary bypass graft fate and patient outcome: angiographic follow-up of 5,065 grafts related to survival and reoperation in 1,388. J Am Coll Cardiol 1996;28:616-626.
- Roffi M, Mukherjee D. Current role of emboli protection devices in percutaneous coronary and vascular interventions. Am Heart J 2009;157:263-270.
- Kirtane AJ, Heyman ER. Correlates of adverse events during saphenous vein graft intervention with distal embolic protection, a PRIDE substudy. J Am Coll Cardiol 2008;1:186-191.
- Baim DS, Wahr D, George B, Leon MB, Greenberg J, Cutlip DE, Kaya U, Popma JJ, Ho KK, Kuntz RE; Saphenous vein graft Angioplasty Free of Emboli Randomized (SAFER) Trial Investigators. Randomized trial of a distal embolic protection device during percutaneous intervention of saphenous vein aortocoronary bypass grafts. Circulation 2002;105:1285-1290.
- 8. Smith SC, Jr, Feldman TE, Hirshfeld JW, Jr, Jacobs AK, Kern MJ, King SB, 3rd, Morrison DA, O'Neil WW, Schaff HV, Whitlow PL, Williams DO, Antman EM, Adams CD, Anderson JL, Faxon DP, Fuster V, Halperin JL, Hiratzka LF, Hunt SA, Nishimura R, Ornato JP, Page RL, Riegel B; American College of Cardiology/ American Heart Association Task Force on Practice Guidelines; ACC/AHA/SCAI Writing Committee to Update 2001 Guidelines for Percutaneous Coronary Intervention. ACC/AHA/SCAI 2005 guideline update for percutaneous coronary intervention: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Circulation 2006;113:e166-286.
- Mehta SK, Frutkin AD, Milford-Beland S, Klein LW, Shaw RE, Weintraub WS, Krone RJ, Anderson HV, Kutcher MA, Marso SP; American College of Cardiology-National Cardiovascular Data Registry. Utilization of distal embolic protection in saphenous vein graft interventions. Am J Cardiol 2007;100:1114-1118.