



Anatomy of Permanent Mandibular First Molars in a Selected Iranian Population Using Cone-beam Computed Tomography

Aria Choupani Dastgerdi^a, Manije Navabi^{b*}, Ladan Hafezi^c, Zohre Khalilak^d, Vahid Rakhshan^a

^a Private Practice, Tehran, Iran; ^b Department of Removable Prosthodontics, Dental Branch, Islamic Azad University, Tehran, Iran; ^c Department of Oral Radiology, Dental Branch, Islamic Azad University, Tehran, Iran; ^d Department of Endodontics, Dental Branch, Islamic Azad University, Tehran, Iran

ARTICLE INFO

Article Type:

Original Article

Received: 14 Nov 2017

Revised: 27 Feb 2018

Accepted: 13 Mar 2018

Doi: 10.22037/iej.v13i2.19035

*Corresponding author: Manijeh Navabi, No. 4, 10th Neyestan St., Pasdaran Ave., Tehran, Iran.

Tel: +98-910 980 8568

E-mail: dr.ma.navabi@gmail.com

ABSTRACT

Introduction: Knowledge of radicular anatomy has a crucial impact on endodontic practices. Since some anatomic features such as modifications of Vertucci are not evaluated adequately, this study was conducted. **Methods and Materials:** In this *in vivo* study, cone-beam computed tomography (CBCT) images of 312 intact bilateral first molars from 156 patients (79 males and 77 females with an average age of 35.58±11.17 years) were investigated by a trained dentist in terms of number of roots, number of canals in each root and in each tooth, and shapes of canals according to Vertucci's classification and its modifications. Groups were compared using the *Chi-square* test. The level of significance was set at 0.05. **Results:** Of all teeth, 5.2% had 3 roots. Mesial roots had mostly 2 canals while distal roots had a similar frequency of 1 and 2 canals. Of all teeth, 39.7% had 3 canals, 45.2% had 4 canals, 13.8% had 5 canals, and 1.3% had 6 canals. There were no significant differences between males and females, in terms of number of roots ($P=0.137$), number of canals in mesial ($P=0.453$) or distal roots ($P=0.328$), and total number of canals ($P=0.138$). The most frequent Vertucci classes in mesial and distal roots were IV followed by II and I, respectively. There were no significant differences between males and females in terms of Vertucci classes of mesial ($P=0.211$) or distal ($P=0.205$) roots. **Conclusion:** In this population, there were 3 to 6 canals per tooth (mostly 4 and 3 canals). Males and female's might be similar regarding the number of roots, or number of canals in each root, number of canals in each tooth, or the predominant canal shape in each root.

Keywords: Anatomy; Cone-beam Computed Tomography; Endodontics; Root Anatomy

Introduction

Proper knowledge of the internal anatomy of root canals is crucial for a successful endodontic treatment which relies on appropriate cleaning and shaping [1, 2]. Since mandibular first molars emerge very early in the mouth and have complex surfaces, they are one of the teeth most requiring root canal therapy; and therefore the knowledge of their root canal anatomy is important [2, 3]. These teeth usually have three canals within two roots although they can change due to ethnicity or normal variations [4-7].

Since ethnic background can affect root anatomy of mandibular first molars, it is important to document properties of

these teeth in various populations. However, studies on Iranians are few and controversial [2, 3, 8-11]. Many aspects of anatomic features are not usually covered by them (such as modifications of Vertucci classification which is not covered in many studies worldwide). Many of older studies have used conventional or 2D radiography techniques that can be less accurate than 3D radiography techniques [12]. Nevertheless, recent studies have mainly used cone-beam computed tomography (CBCT) due to its numerous advantages [8-11].

This study evaluated number and shape of roots and canals of permanent mandibular first molars using CBCT images of an Iranian sample population.




Figure 1. Examples of transverse CBCT sections




Figure 2. Schematic diagrams representing canals within a root [black columns drawn together from top (coronal) to bottom (apical)], arranged as all Vertucci classes (top row) and as certain Vertucci modifications that were observed in this sample of mandibular first molars (bottom row)

Materials and Methods

This *in vivo* study was performed on CBCT images of patients aged 20 to 60 years who had attended two centers in Tehran. All CBCTs had been retrospectively taken solely for clinical purposes. No x-ray was emitted to patients for this study, and study ethics were approved by the research committee of Azad University of Medical Sciences, Dental branch, Tehran, Iran (#14859). All CBCTs had been taken with the same unit (NewTom, GiANO, Verona, Italy); with similar field of view (8×5 cm), focal size (0.3 mm), current (12 mA), peak kilovoltage (85 kVp), and time (0.4 sec). Inclusion criteria were availability of both mandibular first molars and in each patient, full patient information was obtained. Exclusion criteria were resorption, previous endodontic treatment, open apex, agenesis, fractures, or pathologies. A total of 312 images of first molars from 156 patients were included.

All measurements were done by the same dentist trained by a maxillofacial radiologist, using NTT Viewer software program (NTT Software Corporation, Yokohama, Japan). CBCT images were examined in coronal, sagittal, and mainly axial dimensions (Figure 1). Evaluated parameters were number of roots, number of canals in each root and in each tooth, and shapes of canals according

to the Vertucci classification [13] and its modifications [14-17]. Vertucci classes can be summarized based on the number of canals from coronal to apical portions of the root: type I (coronal canals: 1, apical canals: 1 [1-1]), type II (2-1), type III (1-2-1), type IV (2-2), type V (1-2), type VI (2-1-2), type VII (1-2-1-2), type VIII (3-3), type IX (3-1), type X (3-1-2-1), type XI (4-2), type XII (3-2), type XIII (2-3), type XIV (4-4), type XV (5-4), type XVI (1-3), type XVII (1-2-3-2), type XVIII (1-2-3), type XIX (3-1-2), type XX (2-3-1), type XXI (2-3-2), type XXII (3-2-1), and type XXIII (3-2-3). Schematic diagrams of Vertucci classes and certain modifications observed are presented in Figure 2.

Descriptive statistics were calculated. Groups were compared using the *chi-square* of Statistical Package for Social Science (SPSS, version 24.0, SPSS, Chicago, IL, USA). The level of significance was predetermined as 0.05.

Results

The observer was calibrated through repeating the diagnosis of cases (especially more difficult cases) under the supervision of a dental anatomist and an endodontist. Of 156 enrolled patients, 79 were males and 77 were females. Patients' average age was 35.58 ± 11.17 years. Of patients, 101 (64.7%) were aged between 20 and 39 years old, while 55 (35.3%) were between 40 and 60 years old.

Number of roots and canals

Table 1 summarizes the number of canals. Among 312 assessed teeth, 16 (5.2%) bilateral teeth in 8 patients had 3 roots; in all these cases, the third root was distolingual. All other teeth had 2 roots. Mesial roots had mostly 2 canals and in few cases 3 canals (Table 1). Distal roots showed a rather similar distribution of 1 and 2 canals, in addition to very few 3 canals (Table 1). All distolingual roots contained only 1 canal. Overall, number of canals ranged between 3 and 6. Of 312 teeth, 39.7% had 3 canals, 45.2% had 4 canals, 13.8% had 5 canals, and 1.3% had 6 canals. All 6-canaled teeth were 3-rooted. There were no significant differences between males

and females, in terms of number of roots ($P=0.137$), number of canals in mesial ($P=0.453$) or distal roots ($P=0.328$), and total number of canals ($P=0.138$).

Vertucci classifications

Tables 3 and 4 present Vertucci classes and Vertucci modifications. In mesial roots, the most common classes were type IV followed by II. In distal roots, the most common class was type I (Tables 2 and 3, Figure 2). The Chi-square did not show significant differences between males and females in terms of Vertucci classes in the mesial root ($P=0.211$) or distal root ($P=0.205$).

Discussion

The findings of the present study showed that only 5.2% of first molars had three roots. Our results were similar to other Iranian studies reporting prevalence of third roots ranging between zero and 3.9% [2, 8-11]. Also our findings were consistent with studies on Jordan [14], India [18], Turkey [19, 20] and Sudan [21]. However, they were not in line with results from South Korea [22] and Burma [7]. It seems that presence of distolingual roots might depend on ethnicity, and results in Asians indicate a rather high prevalence of the third root (Table 4) [4-7, 22, 23].

In this study, prevalence of three and four canals in each tooth were rather similar. This finding was comparable to results of Kim *et al.* [22], Zhang *et al.* [23], Al-Qudah and Awadeh [14] who reported similar prevalence of three and four canals, but was in contrast to observations of Demirbuga *et al.* [20], Chourasia *et al.* [18], Ballullaya *et al.* [6], Nur [19] and Masoudi [9] who reported a considerably greater prevalence of 3-canaled roots compared to 4-canaled roots. Three-canaled roots can be problematic in endodontic treatments. This research showed prevalence of 3-canaled roots about 13% in mesial roots and 3.5% in distal roots; this was consistent with findings of Al-Qudah and Awawdeh [14], De Pablo *et al.* [5] regarding mesial roots and Ahmed *et al.* [21] regarding distal root. However, our result was not in line with other reports, which might be due to differences in ethnicity or methodologies. In this study, most of mesial roots had 2 canals but about 13% of them had 3 canals. This was similar to other studies stating that 2-canaled roots are much more prevalent (summarized in Table 5). In distal roots, similar prevalence of one or two canals were observed in this study (with very few cases of three canals). Such a high frequency of two canals in this root was similar to studies of Ahmed *et al.* [21], Arjmand *et al.* [10], and Al-Qudah and Awadeh [14]. All distolingual roots were single-rooted as what was reported by Akhlaghi *et al.* [2].

Table 1. Distribution (%) of canal number in roots of mandibular first molars across mesial, distal, and distolingual roots

Root	Canal number		
	1	2	3
Mesial	-	271 (86.9%)	41 (13.1%)
Distal	136 (43.6%)	165 (52.9%)	11 (3.5%)
Distolingual	16 (100%)	-	-

Table 2. Distribution of canal types according to Vertucci classes (I to VIII) in mesial and distal roots of mandibular first molars

		I	II	III	IV	V	VI	VII	VIII
		Mesial root	N	-	66	20	125	13	27
	%	-	21.1	6.4	40.0	4.1	8.6	3.2	-
Distal root	N	136	15	55	-	47	-	31	-
	%	43.6	4.8	17.6	-	15.0	-	9.9	-

Table 3. Distribution of canal types according to Vertucci modifications (IX to XXIII) in mesial and distal roots of mandibular first molars.

Modification with all-empty cells are removed from the table

		X	XII	XIII	XVI	XVII	XX	XXI	XXII
		Mesial root	N	10	9	4	-	7	13
	%	3.2	2.9	1.3	-	2.2	4.2	2.5	-
Distal root	N	17	-	3	4	-	-	-	4
	%	5.4	-	0.9	1.3	-	-	-	1.3

Table 4. Summary of reports on number of roots

Author /Country	Method	Sample size	Number of roots		
			1	2	3
Kim et al. /South Korea [22]	CBCT (<i>in vivo</i>)	1952	0.67%	73%	25%
Abella et al. [4]	review of literature of 45 articles	19056	-	-	14% (Range 0* to 29%)
Zhang et al./China [23]	CBCT (<i>in vivo</i>)	232	0.4%	70%	29%
DePablo et al./Spain [5]	review of literature of 41 articles	18787			13% (Range 0 to 32%)
Demirbuga et al./Turkey [20]	CBCT (<i>in vitro</i>)	1748	0.85%	95.8%	2%
Chourasia et al./India [18]	<i>in vitro</i> (clearing technique)	150		94.6%	5.3%
Garg et al./India [24]	<i>in vitro</i> (periapical radiography)	1054			6%
Al-Qudah, Awawdeh/Jordan [14]	<i>in vitro</i> (clearing technique)	330		96%	4%
Ahmed et al./Sudan [21]	<i>in vitro</i> (clearing technique)	100		94%	3%
Mirzaie et al./Iran [11]	CBCT (<i>in vivo</i>)	66		100%	
Ballullaya et al. [6]	review of literature of 97 articles				Range: 3-35%
Gulabivala et al./Myanmar [7]	<i>in vitro</i> (clearing technique)	139		90%	10%
Nur et al./Turkey [19]	CBCT (<i>in vivo</i>)	966	0.3%	99.2%	0.5%
Zafar et al./Saudi Arabia [25]	CBCT (<i>in vitro</i>)	100		100%	
Arjmand et al./Iran [10]	CBCT (<i>in vivo</i>)	121		97.5%	2.5%
Masoudi et al./Iran [9]	CBCT (<i>in vivo</i>)	129		96.1%	3.9%
Akhlaghi et al./Iran [2]	<i>in vitro</i> (clearing technique)	150		69.7%	3.3%

*In studies from Iran, Uganda, Pakistan, Turkey, Kuwait and Spain, no three-root teeth had been observed

Table 5. Summary of reports on number of canals

Author /Country	Method	Sample size	Number of canals		
			Overall	Mesial root	Distal root
Kim et al. /South Korea [22]	CBCT (<i>in vivo</i>)	1952	48.6% three, 49.2% four	-	-
Zhang et al. /China [23]	CBCT (<i>in vivo</i>)	232	53% three-, 43% four-	95% two-canal	
De Pablo et al. /Spain [5]	review of literature of 41 articles	18787	61% three-, 35.7% four-	94.4% two, 2.3% three	63%: single-canal
Demirbuga et al. /Turkey [20]	<i>in vitro</i> (CBCT)	1748	79.9% three-, 15.4% four-	-	-
Chourasia et al. /India [18]	<i>in vitro</i> (clearing technique)	150	64% three, 36% four		
Al-Qudah, Awawdeh/Jordan [14]	<i>in vitro</i> (clearing technique)	330	49% three-, 46% four-	93% two-, 6% three-	54% single 45% two s
Ahmed et al. /Sudan [21]	<i>in vitro</i> (clearing technique)	100		86% two	59% two-, 38% single, 3% three-
Mirzaie et al. /Iran [11]	CBCT (<i>in vivo</i>)	66	63% three-, 37% four-		
Ballullaya et al. [6]	review of literature of 97 articles		35% four-	Three canals: Range 0.95 to 15%	
Nur et al. /Turkey [19]	CBCT (<i>in vivo</i>)	966	63% three-, 37% four-	96.8% two , 0.2% three, 3% single	
De Pablo et al. /Spain [5]	CBCT (<i>in vitro</i>)	53	41.5% three, 29.4% four, 28.3% five	96.8% two , 0.2% three, 3% single	
Zafar et al. /Saudi Arabia [25]	CBCT (<i>in vitro</i>)	100		95.5% two	20% double, 80% single
Arjmand et al. /Iran [10]	CBCT (<i>in vivo</i>)	121		92.5% two	58.6% single, 37.2% double
Masoudi et al. /Iran [9]	CBCT (<i>in vivo</i>)	129	72.1% three, 24.8% four, 1.6% five		74.7% single
Akhlaghi et al./Iran [2]	<i>in vitro</i> (clearing technique)	150		100% two	61.3% single, 38.7% double

Table 6. Summary of reports on Vertucci types and modifications (all reported values in percent)

Author /Country	Method	Size	Mesial root	Distal root	Vertucci modifications
Kim <i>et al.</i> /South Korea [22]	CBCT (<i>in vivo</i>)	1952	IV 71 II 20	I 66 II 19 IV 12	0.35%
Zhang <i>et al.</i> /China [23]	CBCT (<i>in vivo</i>)	232	IV 81 V 15	-	-
De Pablo <i>et al.</i> /Spain [5]	review of literature of 41 articles	18781	IV 52.3 II 35	I 63 II 14 IV 12.4	
Demirbuga <i>et al.</i> /Turkey [20]	CBCT (<i>in vitro</i>)	1748	IV 68 II 30	I 82 II 6 IV 5.6	
Chourasia <i>et al.</i> /India [18]	<i>in vitro</i> (clearing technique)	150	IV 54 II 36.6	I 65.3 II 20.6 IV 9.3	
Al-Qudah, Awawdeh/Jordan [14]	<i>in vitro</i> (clearing technique)	330	IV 53 II 36	I 54 II 17 V 11 IV 9	Mesial root: 5.7% Distal root: 1.7%
Ahmed <i>et al.</i> /Sudan [21]	<i>in vitro</i> (clearing technique)	100	IV 73 II 14	I 38 II 28 V 22	
Gulabivala <i>et al.</i> /Myanmar [7]	<i>in vitro</i> (clearing technique)	139	IV 38.1 II 28.8 V 6.5	I 92.9	Mesial: 12.9% Distal: 10%
Faraz <i>et al.</i> [26]	<i>in vitro</i> (clearing technique)	123	IV 70.7 II 26.8	I 65.8 II 14.6 V 19.5	
Nur <i>et al.</i> /Turkey [19]	CBCT (<i>in vivo</i>)	966	IV 92 II 5	I 60 II 12 IV 20 V 7	
Zafar <i>et al.</i> /Saudi Arabia [25]	CBCT (<i>in vitro</i>)	100	II 30 IV 27.5 III 20 V 12.5	I 58.6 II 20.3 III 16.9	
Arjmand <i>et al.</i> /Iran [10]	CBCT (<i>in vivo</i>)	121	IV 65.3 II 27.2	I 47.2 III 18.9	
Masoudi <i>et al.</i> /Iran [9]	CBCT (<i>in vivo</i>)	129	II 62.1 IV 29.5	I 74.7 III 18.6 II 17.8	
Akhlaghi <i>et al.</i> /Iran [2]	<i>in vitro</i> (clearing technique)	150	IV 55.3 II 41.3 VIII 2.7	I 61.2 II 26.6 IV 9.4	

The most common types of canal which were found in mesial roots of this study were Vertucci classes IV (40%), II (21.1%) and VI (8.1%), while about 16% of cases were Vertucci modifications. Higher prevalence of type IV followed by type II was seen in most other studies [2, 8] except those reported by Zafar *et al.* [24] and Masoudi *et al.* [9] (Table 7). In distal roots, most common Vertucci types were I (43.6%), III (17.6%), and V (15%); about 9% of cases were Vertucci modifications (Table 6). This was

consistent with studies of Zafar *et al.* [37], Arjmand *et al.* [10], and Masoudi *et al.* [9]. Although Vertucci classes can simplify reports, they are not sufficient to cover all complexities of root canal structures which were observed in this study and few others that have evaluated Vertucci modifications [7, 14]. Sometimes, real canals are much more irregular to be easily categorized into one of Vertucci classes or modifications. Differences might be attributed to ethnicity.

Conclusion

In this population, there were 3 to 6 canals per tooth (mostly 4 and 3 canals). Males and females might be similar regarding number of roots, or number of canals in each root, or number of canals in each tooth. The most frequent Vertucci classes in mesial and distal roots were IV (followed by II) and I, respectively without a significant sex dimorphism.

Conflict of Interest: 'None declared'.

References

- Akhlaghi NM, Abbas FM, Mohammadi M, Shamloo MR, Radmehr O, Kaviani R, Rakhshan V. Radicular anatomy of permanent mandibular second molars in an Iranian population: A preliminary study. *Dent Res J (Isfahan)*. 2016;13(4):362-6.
- Mohammadzadeh Akhlaghi N, Khalilak Z, Vatanpour M, Mohammadi S, Pirmoradi S, Fazlyab M, Safavi K. Root Canal Anatomy and Morphology of Mandibular First Molars in a Selected Iranian Population: An In Vitro Study. *Iran Endod J*. 2017;12(1):87-91.
- Razmi H, Shokouhinejad N, Hooshyar M. An In Vitro Study of the Number of Distal Roots and Canals in mandibular First Molars in Iranian Population. *Iran Endod J*. 2008;2(4):126-30.
- Abella F, Patel S, Durán-Sindreu F, Mercadé M, Roig M. Mandibular first molars with disto-lingual roots: review and clinical management. *Int Endod J*. 2012;45(11):963-78.
- de Pablo ÓV, Estevez R, Péix Sánchez M, Heilborn C, Cohenca N. Root Anatomy and Canal Configuration of the Permanent Mandibular First Molar: A Systematic Review. *J Endod*. 2010;36(12):1919-31.
- Ballullaya S, Vemuri S, Kumar P. Variable permanent mandibular first molar: Review of literature. *J Conserve Dent*. 2013;16(2):99.
- Gulabivala K, Aung TH, Alavi A, Ng YL. Root and canal morphology of Burmese mandibular molars. *Int Endod J*. 2001;34(5):359-70.
- Madani ZS, Mehraban N, Moudi E, Bijani A. Root and Canal Morphology of Mandibular Molars in a Selected Iranian Population Using Cone-Beam Computed Tomography. *Iran Endod J*. 2017;12(2):143-8.
- Masoudi SM, Rouhi N. Evaluation of root canal number and anatomy of first and second mandibular molars with available CBCT in Qazvin in 2013. Qazvin, Iran: Qazvin University of Medical Sciences; 2014.
- Arjmand N, Kolahdouzan A, Rouhi N. Evaluation of Root and Canal Morphology of First and Second Mandibular Molars according to Vertucci and Weine Classification by using Cone - Beam Computed Tomography Archive in Partow Radiology Center in Qazvin in 1391. Qazvin, Iran: Qazvin University of Medical Sciences; 2014.
- Mirzaie M, Tork Zaban P, Mohammadi V. Cone-beam Computed Tomography Study of Root Canals in a Hamadani Population in Iran. *Avicenna J Dent Res*. 2012;4(2):25-31.
- Lee KW, Kim Y, Perinpanayagam H, Lee JK, Yoo YJ, Lim SM, Chang SW, Ha BH, Zhu Q, Kum KY. Comparison of alternative image reformatting techniques in micro-computed tomography and tooth clearing for detailed canal morphology. *J Endod*. 2014;40(3):417-22.
- Ingle J, Bakland L. *Endodontics* (5th edn). BC Decker, Hamilton. 2002.
- Al-Qudah AA, Awawdeh LA. Root and canal morphology of mandibular first and second molar teeth in a Jordanian population. *Int Endod J*. 2009;42(9):775-84.
- Sert S, Bayirli G. Evaluation of the Root Canal Configurations of the Mandibular and Maxillary Permanent Teeth by Gender in the Turkish Population. *J Endod*. 2004;30(6):391-8.
- Peiris R, Takahashi M, Sasaki K, Kanazawa E. Root and canal morphology of permanent mandibular molars in a Sri Lankan population. *Odontology*. 2007;95(1):16-23.
- Faramarzi F, Vossoghi M, Shokri A, Shams B, Vossoghi M, Khoshbin E. Cone Beam Computed Tomography Study of Root and Canal Morphology of Maxillary first Molar in an Iranian Population. *Avicenna J Dent Res*. 2015;7(1):e24038.
- Chourasia HR, Meshram GK, Warhadpande M, Dakshindas D. Root Canal Morphology of Mandibular First Permanent Molars in an Indian Population. *Int J Dent*. 2012;2012:1-6.
- Nur B, Ok E, Colak M, Gungor E, Altunsoy M, Aglarci O. Evaluation of the root and canal morphology of mandibular permanent molars in a south-eastern Turkish population using cone-beam computed tomography. *Eur J Dent*. 2014;8(2):154.
- Demirbuga S, Sekerci AE, Dincer AN, Cayabatmaz M, Zorba YO. Use of cone-beam computed tomography to evaluate root and canal morphology of mandibular first and second molars in Turkish individuals. *Med Oral Patol Oral Cir Bucal*. 2013:e737-e44.
- Ahmed HA, Abu-bakr NH, Yahia NA, Ibrahim YE. Root and canal morphology of permanent mandibular molars in a Sudanese population. *Int Endod J*. 2007;40(10):766-71.
- Kim S-Y, Kim BS, Woo J, Kim Y. Morphology of Mandibular First Molars Analyzed by Cone-beam Computed Tomography in a Korean Population: Variations in the Number of Roots and Canals. *J Endod*. 2013;39(12):1516-21.
- Zhang R, Wang H, Tian YY, Yu X, Hu T, Dummer PMH. Use of cone-beam computed tomography to evaluate root and canal morphology of mandibular molars in Chinese individuals. *Int Endod J*. 2011;44(11):990-9.
- Zafar M, Alrahabi M. Cone Beam Computed Tomography for Exploring Morphology of Mandibular First Molar. *Br J Med Res*. 2015;6(5):514-21.

Please cite this paper as: Choupani Dastgerdi A, Navabi M, Hafezi L, Khalilak Z, Rakhshan V. Anatomy of Permanent Mandibular First Molars in a Selected Iranian Population Using Cone-beam Computed Tomography. *Iran Endod J*. 2018;13(2):251-6. Doi: 10.22037/iej.v13i2.19035.