

# Preanalytical Errors in Hematology Laboratory- an Avoidable Incompetence

Vikram Narang, Harsimran Kaur, Pavneet Kaur Selhi, Neena Sood, Aminder Singh

Dept. of Pathology, Dayanand Medical College & Hospital, Ludhiana, Punjab, India

## KEY WORDS

Quality control  
Preanalytical  
Hematology

## ABSTRACT

**Background:** Quality assurance in the hematology laboratory is a must to ensure laboratory users of reliable test results with high degree of precision and accuracy. Even after so many advances in hematology laboratory practice, pre-analytical errors remain a challenge for practicing pathologists. This study was undertaken with an objective to evaluate the types and frequency of preanalytical errors in hematology laboratory of our center.

**Methods:** All the samples received in the Hematology Laboratory of Dayanand Medical College and Hospital, Ludhiana, India over a period of one year (July 2013-July 2014) were included in the study and preanalytical variables like clotted samples, quantity not sufficient, wrong sample, without label, wrong label were studied.

**Results:** Of 4,71,006 samples received in the laboratory, preanalytical errors, as per the above mentioned categories was found in 1802 samples. The most common error was clotted samples (1332 samples, 0.28% of the total samples) followed by quantity not sufficient (328 sample, 0.06%), wrong sample (96 samples, 0.02%), without label (24 samples, 0.005%) and wrong label (22 samples, 0.005%)

**Conclusion:** Preanalytical errors are frequent in laboratories and can be corrected by regular analysis of the variables involved. Rectification can be done by regular education of the staff.

## ARTICLE INFO

Received 13 Jan 2015;

Accepted 08 Jul 2015;

©Iran J Pathol. All rights reserved.

**Corresponding Information:** Dr. Vikram Narang, Dept. of Pathology, Dayanand Medical College & Hospital, Ludhiana, Punjab, India.  
Email: drvikramnarang@yahoo.com, Tel: 9872045345

COPYRIGHT © 2016, IRANIAN JOURNAL OF PATHOLOGY. This is an open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

## Introduction

Quality assurance in the hematology laboratory is intended to ensure laboratory users of reliable test results. Imprecision and inaccuracy can occur due to pre-analytical, analytical and post analytical variables. With the recent advancement in technology and introduction of automation in hematology laboratories the analytical part of sample analysis is well taken care of if good quality control practices are followed. Post analytical errors like typographical errors can be

avoided with laboratory interface systems and hospital information system in vogue. However, pre-analytical errors remain a challenge for practicing pathologists.

This study was undertaken to evaluate the types of pre-analytical errors in our tertiary care set up (1, 2).

## Materials and Methods

This retrospective analysis was conducted over a period of one year (July 2013-July 2014)

in Hematology Laboratory of Dayanand Medical College and Hospital, Ludhiana, India. All samples received during this period were included in the study. The hospital is 1400-bedded one and sample collection is done using vacutainers by trained technologists and nurses in the out patient department (OPD) collection center and wards respectively. A total of 4,71,006 samples from OPD and in patient department (IPD) were received. The preanalytical variables which hampered the results were classified as:

- Clotted samples
- Quantity not sufficient
- Wrong sample (eg. Coagulation profile in complete blood count vial)
- Without label
- Wrong label

## Results and Discussion

Of the total 4,71,006 samples received in the laboratory, preanalytical errors, as per above mentioned categories was found in 1802 samples. The most common error was clotted samples (1332 samples, 0.28% of the total samples) followed by quantity not sufficient (328 sample, 0.06%), wrong sample (96 samples, 0.02%), without label (24 samples, 0.005%) and wrong label (22 samples, 0.005%) (Table 1). Clotted samples were frequent from the in-patient department however, the exact number could not be ascertained due to paucity of the data.

The haemolysed samples were not included in the study as it was difficult to identify hemolysis

unless we centrifuge sample, conducted more frequently in biochemical tests. Other pre-analytical variables like lipaemic samples and diluted samples were not included in the study.

Lundberg while describing total testing cycle outlined a series of activities, right from arising of doubts in the clinician's mind, test selection, sample collection, sample transport to the laboratory, sample analysis, finally interpretation of reports and decision making by the clinician. These activities have traditionally been divided into three phases as pre-analytical, analytical and post-analytical (1-4).

A laboratory error is usually defined as a defect during the entire testing process (from ordering tests to reporting results) which can influence the quality of laboratory services. Pre analytical and post analytical phases are as important as the analytical phase as correction in the preanalytical variables can reduce the frequency of errors in the later phases (5).

Pre-pre-analytical comprises 46-68% of all errors and included inappropriate test request, patient/specimen misidentification, sample collected from infusion route, inappropriate container, handling, storage and transportation (6). In the present study, the preanalytical errors compound to about 0.38% (1802 samples) which is low. However in the present era of laboratory practice such incompetence can be completely avoided by proper training of nursing staff, phlebotomists and laboratory technical staff (7, 8).

The present study revealed clotted samples

**Table 1**  
Depicting types and frequency of preanalytical errors

	OPD+IPD	Percentage
Total Samples	4,71,006	-
Clotted Samples	1332	0.28
Insufficient quantity	328	0.06
Wrong sample	96	0.02
Wrong label	24	0.005
Without label	22	0.005
Total	1802	0.38

(0.28%) being the main reason of rejecting samples. The common reason of sample being clotted is improper mixing of sample and inadequate EDTA especially if vials are made in house. However in our set up this was attributed to improper mixing. Micro clots if present can be detected on peripheral blood smear but are usually difficult to recognize. No case of microclots was reported in the present study (7).

The insufficient quantity (328 samples, 0.06%) of sample was recognized mainly in pediatric patients where sampling can be some times very difficult especially if done by untrained personnel. Other authors have reported the diluted samples being another cause of rejection in pediatric age group, however this particular variable was not included in our study due to paucity of data (7-9).

Wrong sample (0.005%) and wrong label (0.005 %) were the other two variables identified. These type of errors can have serious adverse effects or can lead to completely wrong treatment of the patient (9).

Diluted samples upto 0.04% is reported and suggested that whenever intravenous fluid is being administered in a patient's arm, blood should be drawn from the opposite arm (10). If an intravenous fluid is running in both arms, sample may be drawn after intravenous infusion is turned off for at least two min before venipuncture and applying the tourniquet below the intravenous infusion site (10).

As laboratories are going for various accreditations, there is lot of emphasis on reducing errors in laboratory practice to minimum possible level. The ISO 15189:2007 also emphasizes the importance of the quality management (QM) and encourages that QM should include internal quality control and inter-laboratory comparisons such as external quality assessment schemes. A list of items that should be included in the quality manual are transportation, collection, handling of samples, reporting of results and communications and other interaction with patients, health professionals, referral laboratories and suppliers

while the monitoring the quality programme. The calibration and instrument function details and analytical system should also be noted. External quality assessment programmes should effectively check the entire examination process, including pre and post-examination procedures (10).

The pre-analytical errors can be avoided by proper training of staff, increased automation in laboratory which may include modern robotic technologies, information systems, computerized order entry, automated phlebotomy tray preparation etc. Barcodes also simplify specimen routing and tracking (11, 12).

## Conclusion

Though there is a lot of development in analytical phase of testing in pathology labs, many errors still occur and they will continue to occur in pre-analytical phase, as there is human intervention in every step, right from filling the requisition form to receiving and preparing the samples for analysis. We believe these can be overcome by better coordination between labs and wards, continuing medical education programmes of laboratory staff, computerization of the labs and competency check of staff.

## Acknowledgements

The authors declare that there is no conflict of interests.

## References

1. Lundberg GD. Acting on significant laboratory results. *JAMA* 1981;245(17):1762-1763
2. Lundberg GD. How clinicians should use the diagnostic laboratory in a changing medical world. *Clin Chim Acta* 1999;280 (1-2):3-11.
3. Laposata M, Dighe A. "Pre-pre" and "post-post" analytical error: high-incidence patient safety hazards involving the clinical laboratory. *Clin Chem Lab Med*

2007;45(6):712-719.

4. Stroobants AK, Goldschmidt HM, Plebani M. Error budget calculations in laboratory medicine: linking the concepts of biological variation and allowable medical errors. Clin Chim Acta 2003;333(2):169-176.

5. Green SF. The cost of poor blood specimen quality and errors in preanalytical processes. Clin Biochem 2013;46(13):1175-1179.

6. Plebani M. The detection and prevention of errors in laboratory medicine. Ann Clin Biochem 2010;47(Pt2):101-110.

7. Ashavaid TF, Dandekar SP, Khodaiji S, Ansari MH, Singh AP. Influence of method of specimen collection on various preanalytical sample quality indicators in EDTA blood collected for cell counting. Ind J Clin Biochem 2009; 24(4):356-60.

8. Uprity S, Upretity S, Bansal R, Jeelani N, Bharat V. Types and Frequency of preanalytical errors in hematology lab. J Clin Diag Res 2013;7(11):2491-93.

9. Favaloro JE, Funk DM, Lippi G. Preanalytical variables in coagulation testing associated with diagnostic errors in hemostasis. Lab Med 2012 ; 43(2): 1-10.

10. International Organization for Standardization. ISO 15189: Medical laboratories - particular requirements for quality and competence. Geneva: International Organization for Standardization; 2007/

11. Hawkins R. Managing the pre and post analytical phases of the total testing process. Ann Lab Med 2012; 32:5-16.

12. Chawla R, Goswami V, Tayal D, Mallika V. Identification of the types of preanalytical errors in the clinical chemistry laboratory: 1 year study of GB Pant Hospital. Lab Med 2010; 41: 89-92.

#### How to cite this article:

Narang V, Kaur H, Kaur Selhi P, Sood N, Singh A. Preanalytical Errors in Hematology Laboratory- an Avoidable Incompetence. Iran J Pathol. 2016;11(2):151-4.

Archive of SID