

Original Article

***Hyalomma aegyptium* on Spur-thighed Tortoise (*Testudo graeca*) in Urmia Region West Azerbaijan, Iran**

*E Tavassoli¹, N Rahimi-Asiabi², M Tavassoli³

¹ Dept. of Biology, College of Science, Urmia University, Urmia, Iran

² Urmia Education Organization, District 2, School of Shahed Dokhtaran 2, Urmia, Iran

³ Dept. of Pathobiology, College of Veterinary Medicine, Urmia University, Urmia, Iran

(Received 28 Jan 2007; accepted 27 Jun 2007)

Abstract

Background: Ticks are obligate blood feeders that parasitize a wide variety of animals. *Hyalomma aegyptium*, parasitize tortoises and other small wild life and livestock. This study was carried out to determine spur-thighed tortoise (*Testudo graeca*) infestation to *H. aegyptium* in Urmia region West Azerbaijan of Iran.

Methods: The study was carried out over a 16 month period from the spring of 2004 to the fall of 2005. A total of 32 tortoises were sampled.

Results: The results indicated that 14 tortoises infected with ticks. A total of 117 ticks were collected from infested animals, the minimum and maximum tick infestation was 1-60. Ticks were attached to the axilla of fore and hind legs of tortoises. All ticks were determined to be *H. aegyptium*.

Conclusion: *H. aegyptium* was the most common tick species in the study area. Due to tendency of some people to keeping tortoise as pet animal, more attention must be done to tortoise's tick infestation. Due to existence of *H. aegyptium* on tortoises in this region more study will need to evaluate presence of this tick on other animal species and its role on transmission of diseases.

Keywords: Ticks, Spur-thighed tortoise, *Hyalomma aegyptium*, Iran

Introduction

Testudo graeca Linnaeus 1758, or the spur-thighed tortoise, is an endangered species with a broad distribution range. It can be found in northern Africa (e.g. Morocco, Algeria, Tunisia and Libya), the Middle East (e.g. Lebanon, Jordan, Syria, and Iraq), Europe (Bulgaria, Romania, Turkey, Greece, and multiple introductions into Spain and Greece), and in Asia (e.g. Armenia, Azerbaijan, Georgia, Turkmenistan, Iran, and possibly Afghanistan) (1-3).

Ticks are obligate blood feeders that parasitize a wide variety of terrestrial and flying vertebrates and a few marine snakes and lizards (4). More than 800 tick species have been described in the world (5, 6). Ticks are vectors of more kinds of microorganisms (5, 7). The vast litera-

ture regarding ticks has centered mostly around 10% of the world tick fauna, which have been well recognized for their medical and veterinary significance. However, a comprehensive knowledge of the ecology of important ticks and tick borne-diseases are best achieved by knowing the biological and physiological similarities and differences between all species, in relation to their hosts and to the environment. Knowledge of biological models of tick parasitism of wildlife is very useful to clarify factors that have permitted a few tick species to become economically important pests and vectors of disease agents to man and animals (4).

Hyalomma ticks are often the most abundant tick parasites of livestock, in warm, arid, and semiarid, generally harsh lowland and middle altitude biotopes, and those with long dry sea-

sons, from central and southwest Asia to southern Europe and southern Africa. Of the 30 known *Hyalomma* spp., ≥ 15 are important vectors of infectious agents to livestock and humans. Hyalommines are mostly moderately large to large ticks with long mouthparts (8). The subgenus *Hyalomma* contains 15 species of veterinary and public health importance. Three of the 15 species have 2, 3, and 4 subspecies, respectively. *H. aegyptium*, parasitize tortoises and small wildlife and livestock from Pakistan to both sides of the Mediterranean basin, and Russia (8, 9). Adults are specific for tortoises. This study was carried out to determine tortoise's infestation to *H. aegyptium* in Urmia region, West Azerbaijan of Iran.

Materials and Methods

This study was carried out over a 16 month period from the spring of 2004 to the fall of 2005, in Urmia West Azerbaijan Province, North West of Iran. This area is semi-humid, with mean rainfall of about 350 mm with the maximum mean temperature of 28.3° C in August and the minimum mean monthly temperature of -5 ° C in January. Tortoise from suburb Urmia

city, were captured and inspected for ticks. Captured tortoises were restrained and given a thorough physical examination. Ticks collected with forceps on the tortoises were immediately placed into screw-capped tubes containing several minute holes. Vials were properly identified and conditioned under room temperatures for few days or weeks, and then they were sent to the laboratory. The purpose of this procedure was to try to maintain ticks alive inside the vials until arriving at the laboratory for taxonomic identification. Adult ticks were morphologically examined under the stereo microscope and identified using the keys of Hoogstraal and Kaiser and Hillyard (10, 11).

Results

A total of 32 tortoises were sampled. The results indicated that 14 tortoises infected with ticks. A total of 117 ticks were collected from infested animals, the minimum and maximum tick infestation was 1-60. Ticks were attached to the axilla of fore and hind legs of tortoises (Fig. 1). All ticks were determined to be *H. aegyptium*, a common tortoise parasite (Fig. 2, 3, 4 & 5).



Fig.1: *H. aegyptium* ticks attached to axilla of hind leg



Fig. 2: *H. aegyptium* male tick, dorsal view



Fig. 3: *H. aegyptium* female tick, dorsal view



Fig. 4: *H. aegyptium* male tick, ventral view, arrow show short external spur of coxa I.



Fig. 5: *H. aegyptium* male tick, ventral view, arrow show broad adanal plates and reduced subanal plates.

Discussion

Ticks biology and their distribution studies in Iran were initiated in 1810 when Dupre visited this country (12). Razi Institute, Pasteur Institute of Iran, Faculties of Veterinary and School of Public Health, continued their works on Iranian ticks (13). In 1935, Brumpt had conducted a study on genus *Ornithodoros* ticks (14). Subsequently, Delpy published a paper on the family of Ixodidae genus *Hyalomma* in 1936 (15). Baltazard explained the characteristics of *Ornithodoros* ticks (16). Abbasian listed the name of Iranian ticks in 1960 (13). The book entitled "Ectoparasites of domestic animals' ticks" was published by Maghami in 1968 (17). Mazlum published his research on the geographical distribution, seasonal activities and host preference of ticks in 1971 (18). Rahbari worked on some ecological aspects of tick fauna of West Azerbaijan, Iran (19). The occurrence of 55 ticks' species was previously reported can be summarized as follow: *Argas* 5 spp., *Ornithodoros* 6 spp., *Hyalomma* 16 spp., *Boophilus* 2 spp., *Haemaphysalis* 11 spp., *Ixodes* 6 spp., *Dermacentor* 4 spp. and *Rhipicephalus* 5 spp.

Ticks are a group of arthropod that can be found on reptiles. These ticks are not usually dangerous to humans, although they can bite humans and family pets, but can carry several different diseases that can infect humans, such as relapsing fever and western equine encephalitis virus. Results of recent studies have shown the ease with which exotic ticks have been introduced into other countries on imported reptiles and disseminated from importers to breeders, zoos, wildlife theme parks, pet stores and private hobbyists (20-22).

It has been known for many years that reptiles imported into other locations were on occasion infested with ticks (23-27). At least eight exotic tick species were being imported into Florida on reptiles (21,28).

New findings demonstrated clearly the potential for at least five of the exotic reptilian ticks to spread rapidly, not only to other reptilian species, but in the case of one also to domestic mammals such as dogs, these ticks have been associated with mammalian diseases, heartwater (29-32) and Q fever (33,34). In addition, these hard ticks species have been reported to be vectors of various haemogregarines of reptiles (35-38),

and in heavy infestation of ticks it cause respiratory distress and even death in monitors (28, 39). Remarkably little data are available on the impact of reptilian ticks on their primary hosts, reptiles. Although parasitaemia may be high and of long duration, there is little documented evidence that haemogregarines are pathogenic to their reptilian hosts (37).

Some studies have documented the frequent occurrence of ticks, specifically *Ornithodoros parkeri*, as ectoparasites of tortoises. Also, tick transmission of mycoplasmas involved in bovine pleuropneumonia disease in Africa has been demonstrated in some earlier experimental studies but not confirmed under natural conditions (16, 40, 41).

In southern Europe the hosts of *H. aegyptium* are primarily tortoises but also lizards, dog, horse, hedgehog, hamster and birds. In Italy *H. aegyptium* has occurred on partridge, in Egypt, on quail, pigeon, chats and warblers (11). *H. aegyptium* were reported from cattle and buffaloes from, Pakistan, Turkey and India (42-44). Vashishta and Mathur and Vashishta *et al.* reported that *H. aegyptium* transmit *Theileria hirci* in goats (45, 46). Our results showed that *H. aegyptium* was the most common tick species in the study area. In our knowledge *H. aegyptium* was not reported from other animals except *Testudo graeca* turtle in Iran (47). As far as we know, *H. aegyptium* is not responsible for any human and domestic animal pathology in North-West Europe (11). In Mediterranean region, it is a vector of protozoa which are blood parasites of land tortoises (38). However, Ece *et al.*, 2003, report a spirochete of the genus *Borrelia* from the hard tick *H. aegyptium* in Turkey (48).

Crimean-Congo hemorrhagic fever virus (CCHFV) is the most often transmitted to man following a tick bite (genus *Hyalomma*) (49). In Iran, *Hyalomma* spp. probably plays the main role in transmitting the infection from animals to humans (50). Due to tendency of some people to keeping tortoise as pet animal, more attention was done to tortoise's tick infestation;

however the risk of contracting a disease from a reptile is generally small, as long as owners practice good hygiene. But people with a suppressed immune system are more at risk than the general population. For example, children under 10 and the elderly are considered to be at higher risk. However, by practicing good sanitation and personal hygiene, keeping tortoise out of the kitchen and food preparation areas, it is possible to minimize the risk. It is also important to have all new tortoise examined and tested prior to introducing them to home.

Many insecticide were used for tick control on tortoise for example Amitraz at a concentration of 2 ml litre⁻¹ of water successfully induced detachment of *Amblyomma marmoratum* and *Amblyomma hebraeum* ticks from the mountain tortoise, *Geochelone pardalis*(51).

In conclusion, due to existence of *H. aegyptium* on tortoises in this region more study will need to evaluate presence of this tick on other animal species and its role on transmission of diseases.

Acknowledgements

Authors would like to thank Mr E Agapour for his technical assistance.

References

1. Boyan P, Vladimir P, Popgeorgiev BG, Plachyiski D. National Action Plan for Tortoises Conservation in Bulgaria, Vers.1, BSPB, NMNHS-BAS, Sofia; 2003.
2. Buskirk JR. Of the absence of spur-thighed tortoises, *Testudo graeca*, from Egypt. Chelonian Conserv Biol. 1996; 2: 118-20.
3. Beshkov VI, Nanev K. Amphibians and Reptiles in Bulgaria.-Pensoft, 2002. p. 120.
4. Hoogstraal H. Argasid and Nuttalliellid ticks as parasites and vectors. Adv Parasitol. 1985; 24: 135-238.
5. Keirans JE. Systematic of the Ixodida (Argasidae, Ixodidae, Nuttalliellidae): an overview and some problems. In: B Fivaz, T Petney, I Horak (Eds), Tick Vector Biol-

- ogy. Medical and Veterinary Aspects, Springer-Verlag, Berlin; 1985. p. 1-21.
6. Camicas JL, Hervy JP, Adam F, Morel PC. Les Tiques du Monde. Nomenclature, Stades Décrits, Hôtes, Répartition (Acarida, Ixodida), Orstom, Paris; 1998.
 7. Oliver Jr JH. Biology and systematic of ticks (Acari: Ixodida). Annu Rev Ecol Syst. 1989; 20: 397-430.
 8. Fraser, CM. The Merck Veterinary Manual, 6th ed. Merck & Co: Rahway, N.J., USA, 1986.
 9. Robbins RG, Karesh WB, Calle PP, Leontyeva OA, Pereshkolnik SL, Rosenberg S. First records of *Hyalomma aegyptium* (Acari: Ixodida: Ixodidae) from the Russian spur-thighed tortoise, *Testudo graeca nikolskii*, with an analysis of tick population dynamics. J Parasitol. 1998; 84(6): 1303-5.
 10. Hoogstraal H, Kaiser MN. Observation on Egyptian *Hyalomma* ticks (*Ixodoidae*, *Ixididae*) 5. Biological notes and differences in identity of *Hyalomma anatolicum* and its subspecies *anatolicum* Koch, and *ex-cavatum* Koch among Russian and other workers. Identity of *H. lusitanicum* Koch. Ann Entomol Soc Am. 1959; 52: 243-61.
 11. Hillyard PD. Ticks of North-West Europe. FSC. UK; 1996.
 12. Janbakhsh B. Report on studies of the tick vectors of relapsing fever in Iran. p. 34. Sofiev 1941. Note préliminaire. Ann Parasit Humaine et compare. 1956; 27: 311- 28.
 13. Abbassian-Lintzen R. A preliminary list of ticks (Acarina: Ixodidae) occurring in Iran, and their distributional data. Acarol. 1960; 2(1): 43-61.
 14. Brumpt E. Presentation de deux Ornithodorus canstrinii Bir 1895, vivants originaires d Isfahan (Perse). Bull Soc Path Exot. 1935; 28: 51- 3.
 15. Deply L. Notes sur les Ixodidés du genre Hyaloma (koch). Ibid. 1936; 14: 206-45.
 16. Baltazard M, Bahmanyar M, Pournaki R, Mofidi Ch, Chama M. *Ornithoderes takovsky* Olenov 1931 et *Borrelia* (Spirochaeta) *latychevi* 1941. Note préliminaire. Ann Parasit Humaine et compare. 1952; 27: 311-28.
 17. Maghami G. External Parasite of Live stocks in Iran. Arch Razi Ins. 1965; 20: 81- 3.
 18. Mazlum Z. Different ticks occurring in Iran (geographical distribution, seasonal activities, hosts). Bull Fac Vet Med. 1971; 27(1): 1-32.
 19. Rahbari S. Studies on some ecological aspects of tick fauna of West Azerbaijan, Iran. J Appl Anim Res. Sup 1995; 7(2): 189-94.
 20. Allan SA, Simmons LA, Burr ridge MJ. Establishment of the tortoise tick *Amblyomma marmoratum* (Acari: Ixodidae) on a reptile-breeding facility in Florida. J Med Entomol. 1998; 35: 621-24.
 21. Burr ridge MJ, Simmons LA, Allan SA. Introduction of potential heartwater vectors and other exotic ticks into Florida on imported reptiles. J Parasitol. 2000a; 86: 700-4.
 22. Simmons LA, Burr ridge MJ. Introduction of the exotic ticks *Amblyomma humerale* Koch and *Amblyomma geoemydae* (Cantor) (Acari: Ixodidae) into the United States on imported reptiles. Inter J Acarol. 2000; 26: 239-42.
 23. Becklund WW. Ticks of veterinary significance found on imports in the United States. J Parasitol. 1968; 54: 622-28.
 24. Anderson JF, Magnarelli LA, Keirans JE. *Aponomma quadricavum* (Acari: Ixodidae) collected from an imported boa, *Epicrates striatus*, in Connecticut. J Med Entomol. 1981; 18: 123-125.
 25. Anderson JF, Magnarelli LA, Keirans JE. Ixodid and argasid ticks in Connecticut, U.S.A.: *Aponomma latum*, *Amblyomma dissimile*, *Haemaphysalis leachi* group, and *Ornithodoros kelleyi* (Acari: Ixodidae, Argasidae). Inter J Acarol. 1984; 10: 149-51.
 26. Wilson N, Barnard SM. Three species of *Aponomma* (Acari: Ixodidae) collected from imported reptiles in the United States. Florida Entomologist. 1985; 68: 478-80.

27. BurrIDGE MJ. Ticks (Acari: Ixodidae) spread by the international trade in reptiles and their potential roles in dissemination of diseases. *Bul Entomol Res.* 2001; 91: 3-23.
28. Norval RAI. The ticks of Zimbabwe. XI. The genus *Aponomma*. *Zimbab Vet J.* 1985; 16: 5-8.
29. Norval RAI, Mackenzie PKI. The transmission of *Cowdria ruminantium* by *Amblyomma sparsum*. *Vet Parasitol.* 1981; 8: 189-91.
30. Peter TF, BurrIDGE MJ, Mahan SM. Competence of the African tortoise tick, *Amblyomma marmoreum* (Acari: Ixodidae), as a vector of heartwater (*Cowdria ruminantium* infection). *J Parasitol.* 2000; 86: 438-41.
31. Ghiretti M, Mwanaumo B. *Amblyomma marmoreum* on tortoises of southern province, Zambia. *J Wildl Dis.* 1989; 25(4): 634-35.
32. BurrIDGE MJ, Simmons LA, Simbi BH, Peter TF, Mahan SM. Evidence of *Cowdria ruminantium* infection (heartwater) in *Amblyomma sparsum* ticks found on tortoises imported into Florida. *J Parasitol.* 2000b; 86: 1135-36.
33. Babudieri B. Q fever: a zoonosis. *Adv Vet Sic.* 1959; 5: 81-182.
34. Arthur DR. Ticks and diseases. Evanston, Illinois, Row, Peterson and Company; 1962.
35. Elbl A, Anastos G. Ixodid ticks (Acarina, Ixodidae) of Central Africa. Volume I. General introduction, genus *Amblyomma* Koch, 1844. Tervuren, Belgium, Annales du Musée Royal de l'Afrique Centrale, Sciences Zoologiques no. 145 ; 1966a.
36. Elbl A, Anastos G. Ixodid ticks (Acarina, Ixodidae) of Central Africa. Volume IV. Genera *Aponomma* Neumann, 1899, *Boophilus* Curtice, 1891, *Dermacentor* Koch, 1844, *Haemaphysalis* Koch, 1844, *Hyalomma* Koch, 1844 and *Rhipicentor* Nuttall and Warburton, 1908, lists and bibliography. 412 pp. Tervuren, Belgium, Annales du Musée Royal de l'Afrique Centrale, Sciences Zoologiques no. 148 ; 1966b.
37. Telford SR. Haemoparasites of reptiles In: Hoff GL, Frye FL, Jacobson ER (Eds) Diseases of amphibians and reptiles. New York, Plenum Press; 1984. p. 385-517.
38. Hoostraal H. African Ixodoidea. Vol. 1. Ticks of the Sudan (with special reference to Equatorial Province and with preliminary reviews of the genera *Boophilus*, *Margaropus* and *Hyalomma*). Department of the Navy, Bureau of Medicine and Surgery, Washington, DC; 1956.
39. Young E. *Aponomma exornatum* (Koch) as a cause of mortality among monitors. *J South Afr Vet Med Assoc.* 1965; 36: 579.
40. Joseph G, Kristin T, Berry H, Henen BT. A Search for *Mycoplasma* in *Ornithodoros parkeri* Ticks collected from the Desert Tortoise (*Gopherus agassizii*) in the Mojave, Colorado, and Sonoran Deserts, Twenty-Third Annual Meeting and Symposium of the Desert Tortoise Council, April 3-5, 1998.
41. Brown MB, Schumacher IM, Klein PA, Harris K, Correll T, Jacobson ER. *Mycoplasma agassizii* causes upper respiratory tract disease in the desert tortoise. *Infect Immun.* 1994; 62: 4580-86.
42. Khan MN. Prevalance of ticks on livestock in Faisalabad (Pakistan). *Vet J.* 1993; 13(4): 182-84.
43. Aydn L. Distribution and species of ticks on ruminants in the southern Marmara Region. *Turkey-parasitol-Dergisi.* 2000; 24(2): 194-200.
44. Sivasankar-v: Rao-PrR. Tick fauna of Andhra Pradesh. *Livestock-Adviser-Bangalore.* 1984; 9(8): 29-33.
45. Vashishta MS, Mathur PD. Observation on a fatal outbreak of theileriosis in goat. *Indian J Anim Health.* 1983; 26(1): 61-2.
46. Vashishta MS, Mathur PD, Goswami SK. Fatal goat theileriosis in India. *Indian J Anim Health.* 1987; 28(7):51-52.
47. Nabian S, Mirsalimi SM. First report of presence of *Hyalomma aegyptium* tick from *Testudo graeca* turtle in Iran. *J Fac Vet Med Univ Tehran.* 2002; 57(3): 61-3.

48. Ece SG, Hashimoto N, Kadosaka T, Imai Y, Masuzawa T. A novel, fast-growing *Borrelia* spp. isolated from the hard tick *Hyalomma aegyptium* in Turkey. *Microbiol.* 2003; 149: 2539-44.
49. Faye O, Cornet JP, Camicas JL. Experimental transmission of Crimean-Congo hemorrhagic fever virus: role of 3 vector species in the maintenance and transmission cycles in Senegal. *Parasite.* 1999; 6: 27- 32.
50. Chinikar S. The specific serological investigation of suspected human and animals to have Crimean-Congo hemorrhagic fever in various parts of Iran using ELISA. *Hakim.* 2002; 4: 294- 300.
51. Petney TN, Knight. The treatment of ticks on tortoises using amitraz, *J S Afr Vet Assoc.* 1988; 59 (4): 206.

Archive of SID