



Tehran University of Medical  
Sciences Publication  
<http://tums.ac.ir>

## Iranian J Parasitol

Open access Journal at  
<http://ijpa.tums.ac.ir>



Iranian Society of  
Parasitology  
<http://isp.tums.ac.ir>

### Original Article

## Onchocerciasis in the Upper Imo River Basin, Nigeria: Prevalence and Comparative Study of Waist and Shoulder Snips from Mesoendemic Communities

EC Uttah

Department of Biological Sciences, Faculty of Science, Cross River University of Technology, Calabar, Nigeria

(Received 21 Feb 2010; accepted 19 May 2010)

### Abstract

**Background:** Onchocerciasis is endemic in the Imo River Basin, Nigeria. This study was aimed at assessing the prevalence and intensity of microfilaria of *Onchocerca volvulus* in the area.

**Methods:** A cross-sectional study was carried out in the Okigwe Local Government Area, Imo State, Nigeria. Two skin snips (one from the waist and another from the shoulder) were taken from 1024 individuals examined. The survey coverage was high (91.8% of the study population). An individual was considered mf positive if either of the waist or shoulder snips or both were mf positive. The SPSS for Windows package was used for entering and analysis of data.

**Results:** Thirty-seven percentage of those examined was positive for *Onchocerca volvulus* microfilariae (39.2% of males and 34.9% of females). The mf prevalence increased steadily with increasing age to reach 70.4% in the oldest age group. The overall mf Geometric Mean Intensity among mf positive individuals was 16 mf/skin snip and was significantly higher among males (18 mf/skin snip) than females (14 mf/skin snip) ( $p < 0.01$ ). A scatter plot of microfilariae numbers in snips from the waist against numbers in snips from the shoulder of the same individuals, showed close correlation (Pearson's correlation coefficient = +0.90;  $p < 0.01$ ), and those with mf intensities below 10 mf/snip had a more scattering tendency away from the regression line than those with higher mf intensities.

**Conclusion:** Onchocerciasis is a public health concern in the area. Perhaps, 10 mf/snip is critical intensity threshold for reliable sampling using corneo-scleral punch.

**Keywords:** Onchocerciasis, *Onchocerca volvulus*, Prevalence, Corneo-Scleral Punch, Nigeria

\*Correspondence: Telephone: +234-806-408-7437, e-mail: drecuttah@yahoo.com

## Introduction

**O**ncchocerciasis is a severe and debilitating parasitic infection of global concern. Its prevalence and the magnitude of associated social and economic effects vary widely in different geographical areas where the disease occurs (1-2). About 90 million people are at risk of which 17.6 million are infected, including 326,000 people who have gone blind, in 34 countries of the world (3). In Africa alone, home to over 96% of all global cases, it has been reported in 26 countries (1).

Onchocerciasis is perhaps the most studied filarial infection in Nigeria. The provisional estimates had suggested that 7-10 million Nigerians are infected with *Onchocerca volvulus*, approximately 40 million are at risk of the disease (4), and 120,000 cases of onchocerciasis-related blindness (1), with many thousands suffering from disabling complications of the disease (5). New foci of onchocerciasis are still being discovered and therefore its distribution could be far more extensive than has been earlier assumed (6). In southeastern Nigeria, there are pockets of endemic foci, as shown by some reported studies (7-13) although there is gross underreporting of the scourge. Arguably, the most significant area in this sub-region as far as onchocerciasis is concerned, is the hilly and undulating Udi-Enugu-Okigwe axis from where some rivers or their tributaries, supporting black fly vector breeding, have their origin. These include rivers such as Oji, Ajali, Mamu, Adada, and Imo (the biggest of them). Unfortunately, studies in this sub-region have been largely cross-sectional. There has not been any comprehensive study on all aspects of the infection: parasitological, clinical, and epidemiological.

This report is the first emanating from this holistic study on meso-endemic onchocerciasis. It is aimed at assessing the prevalence

and intensity of microfilariae of *O. volvulus*, and to compare the microfilariae and mf intensity of skin snips from the waist and shoulders of same individuals with a view to ascertaining the critical threshold for reliable sampling using the corneo-scleral punch.

## Materials and Methods

### *Study area and study population*

The study was conducted in 2005 in two neighboring high altitude communities of Umuowaibu 1 and Ndiorji, in Okigwe Local Government area of Imo State, Nigeria. The two communities with a combined population of 1,116 at the time of this study are socio-culturally similar, both inhabited by Ndiigbo, the majority tribe in southern, Nigeria. A familial settlement pattern was evident in the area with houses arranged in family clusters. A total of 381 houses were recorded in the two communities, 216 in Umuowaibu1 and 165 in Ndiorji, giving an overall average of three persons per house. The area is hilly with characteristic undulating plains. There are a total of seven streams and three rivers in addition to the Imo River. There are two distinct seasons, the dry season (November – May), and the rainy season (June – October). According to data from the Imo State Meteorological Service, the annual rainfall in the area averages 2,840 mm per annum, with most of the rainfall occurring in the months of June through October. The mean relative humidity was between 68.7% and 74.4%. Farming is the main occupation, however, those who are engaged in other occupations engage in subsistence farming.

### *Preparation for the study*

Local Government Area (LGA) health authorities were contacted and their consent

obtained before the actual work began. Furthermore, the local Ezes (traditional rulers of Ndiigbo), chiefs, and leaders of town development unions were briefed about the project, and their cooperation was sought in the mobilization of their people. During the parasitological and clinical surveys, health personnel from the LGA were always present to monitor safety standards.

### **Census and mapping**

All individuals in the selected communities who were more than one year of age were included in the study population, which comprised natives as well as non-natives who had resided there for at least one year. The target population was 1000 persons. During the census, the house registration numbers written on houses by the National Population Commission during the 1993 national census were used. Where such numbers were unavailable, one was provided. The appropriate positions of houses, markets, religious places, major roads and some track roads, as well as water bodies in the communities were noted.

### **Skin snipping**

Two skin snips (one from the shoulder and one from the waist) were taken for parasitological examination from each individual during daytime using a Walser corneo-scleral punch. The size of the biopsies was known to average 0.8 mg., with a range of 0.4 – 1.2 mg (14). The biopsies were placed in micro-titer wells containing 0.2 ml of 0.85% saline solution. When completed, each plate was covered with cellophane tape and taken to the laboratory where it was kept for 24 hours at room temperature (15). At the end of the 24-hour incubation period, the skin biopsies were fixed in formalin solution (35% formaldehyde solution) by adding two drops per micro-well. This was adapted from earlier studies (16-18). Thereafter, emerged microfilariae were observed and counted mi-

croscopically using x 40 magnification. Verification of the microfilariae as *O. volvulus* had been carried out in an earlier study by staining with Giemsa (19).

### **Data analysis**

The Epi-Info version 6.0 was used in entering data from parasitological survey, and SPSS for windows (1995 version) was used for data analysis. The geometric mean intensity (GMI) of microfilariae was calculated as  $\text{antilog} (\sum \log (x+1)/n)$ , with x being the number of mf per ml of blood in microfilaraemic individuals and n the number of microfilaraemic individuals examined.

## **Results**

An individual was considered mf positive if either of the waist or shoulder snips or both were mf positive. The mf intensities presented are averages of the two snips taken from each individual. The survey coverage was high (91.8% of the study population; 93.0% for males and 96.6% for females).

Thirty-seven percentage of those examined were positive for *O. volvulus* microfilariae (39.2% of males and 34.9% of females) and microfilariae appeared early in life. The mf prevalence in relation to age and sex is presented in Fig. 1. The youngest mf positive boy and girl were both four years old. In the youngest age group, 13.9% were positive for microfilaraemia. The mf prevalence increased steadily with increasing age to reach 70.4% in the oldest age group (69.2% for males and 71.4% for females). There was no significant difference in overall mf prevalence between males and females ( $\chi^2$ -test;  $P > 0.05$ ). However, in the age groups 20-39 and 40-59 years, males had significantly higher mf prevalences than females ( $\chi^2$ -test;  $P < 0.05$  for both tests).


The overall mf GMI among mf positive individuals was 16 mf/skin snip (18 mf/skin snip

for males and 14 mf/skin snip for females). Males had higher mf GMI in all age groups except in the 10-19 years age group where it was the same in both sexes. The overall difference in mf GMI between males and females was statistically significant ( $t$ -test;  $P < 0.01$ ). The mf GMI in the first three age groups was comparable. The mf GMI rose significantly from 20-39 to the 40-59 years age group, and also from the 40-59 to the 60+ years age group, to peak at 35 mf/skin snip in the latter ( $t$ -test;  $P < 0.001$  for both tests). The highest individual mf intensity observed was 127 mf/skin snip (119 mf/skin snip from the waist snip and 134 mf/skin snip from the shoulder snip) in a 40 year-old man.

Comparison of microfilariae number of skin snips from the waist and the shoulder of the same individual showed that all who were positive in the shoulder skin snip were also positive in the waist skin snip. Six people

were positive in the waist snip but negative in the shoulder snip. The mf GMI of those who were positive in both the waist and shoulder snippings (GMI = 16 mf/skin snip, ranging from 1 to 127 mf/skin snip) exceeded significantly the mf GMI of those who were positive only in the waist snippings (GMI = 2 mf/skin snip, ranging from 1 to 2 mf/skin snip).

A scatter plot of microfilariae numbers in snips from the waist against numbers in snips from the shoulder of the same individuals, showed that the mf counts from the waist and those from the shoulder were closely correlated (Pearson's correlation coefficient = +0.90;  $P < 0.01$ ). Expectedly, there was a more spread-out scattering of the low mf intensities, and furthermore, those with mf intensities below 10 mf/snip had a more scattering tendency away from the regression line than those with higher mf intensities.



**Fig. 1:** Prevalence of *Onchocerca volvulus* microfilaria in relation to age and sex

## Discussion

The Upper Imo River Basin is traversed by rivers and several streams, and conditions are favorable for transmission of onchocerciasis. Eleven species of *Simulium*, including *S. damnosum s.l.* have thus been reported from the area (9). Since microfilarial density is important in on-set of clinical manifestations (8), the high prevalence of onchocerciasis-related skin manifestations reported in the UIRB (20), may be indicative of higher prevalence than reported here. This area should probably be regarded as the most severe onchocerciasis focus in the eastern part of Nigeria, forming a continuum of high onchocerciasis Endemicity with the Udi Hill range, Oji River Basin and the Anambra River Basin. The prevalence here is higher than reported in other areas in the region such as Ezeagu, Oji River and Uzo-Uwani areas of Enugu State (13), Igwun River Basin (21) and the Niger Delta (8). In the Lower Imo River Basin (LIRB), however, onchocerciasis was found only sporadically, and most infected were immigrants from endemic areas (22). It is worth noting that the skin-dwelling microfilaria of *Mansonella streptocerca* was not found in the Imo River Basin in this study, nor was it found in the neighboring areas of southeastern Nigeria where skin-snip examinations have been carried out for microfilariae (8). This is contrary to the stipulations of Sasa (23) that *M. streptocerca* could be higher in the rainforest Nigeria than in the northern Savannah areas. On the contrary, *M. streptocerca* has been reported mostly in the Savannah parts of Nigeria (24-25), and among Lagos residents (26), a cosmopolitan town, and home to Nigerians from several parts.

The mf prevalence increased steadily with age. This agrees with findings in Nigeria (8, 17, 27-29). This may be because older individuals have been exposed throughout their

lives and that they are more exposed to the vectors because of their occupations, mostly as farmers in the fields, as opposed to the children who are attending school. The peak in biting in the late morning hours in the UIRB (19), thus coincides with adults being in their fields while the children are in school. First vector contact occurs when children visit streams to swim, to fetch water for domestic use or for other purposes. Later in life, exposure continues during farming and other adult occupations. Resistance development is limited and the age-related prevalence and intensity therefore shows a gradual increase. In the highly endemic UIRB, there is frequent exposure to infective vectors from early childhood, and consequently infection is contracted early in life.

The prevalence of microfilaria was not significantly different between the sexes. This is in accordance with the results from the neighboring mesoendemic areas of Ezeagu and Oji-River Basin and Uzo-Uwani in Enugu State (13), and Bali District communities of Taraba State in northern Nigeria (30). It does not agree, however, with the findings of Anosike and Onwuliri (29) in Bauchi State, an area that is predominantly Moslem where traditional restrictions imposed on the women result in less exposure and thereby in lower prevalences than in men.

The overall GMI of 16 mf/skin snip among mf positive individuals in the UIRB study population may be regarded as low when compared with the mean microfilarial density of 50 mf per skin snip in Ezeagu, Orji-River, and Uzo-Uwani areas (13), and 169.2 mf/skin snip and 67.2 mf/skin snip observed in different areas of the Taraba River Valley (30-31). This may be due to ivermectin treatment in the area carried out by the Nigerian Onchocerciasis Control Project (NOCP),

which expectedly resulted in a drastic reduction in the microfilarial density and possibly reduction in the prevalence of microfilariae (32-36). This argument is corroborated by the results in the LIRB where there was no ivermectin treatment, and onchocerciasis is only sporadic, the overall mean mf intensity was 22 mf/skin snip (21). Furthermore, studies have shown that after treatment with ivermectin, microfilariae would be reduced by half after 24 hours, by 85% after 72 hours, by 94% after one week, and by 98–99% after one to two months (37). Multiple treatments with ivermectin have marked effects on embryogenesis (38). Quantitative estimates have ranged from an irreversible decline in microfilarial production of approximately 30% per treatment (39), a reduction in the productivity index of 83% (40) to arrest of development at the single cell stage (41). Thus, when the factors listed above are taken into account, the presence of normal reproductive activity in female worms exposed to multiple treatments is incompatible with a normal response (42). However, there could be significant microfilaridermias despite multiple treatments with ivermectin and is mainly attributable to the non-response of the adult female worms and not to inadequate drug exposure or other factors (42). This was confirmed in a 30-month follow up study (43). Microfilariae in general remained sensitive to Ivermectin (42, 44).

The mf intensity was equal among the males than among females. This is consistent with the observations on another filarial species, *Wuchereria bancrofti* (45) that the microfilaria intensity is lower among women of reproductive age than among males of the same age.

Although the skin snips from the waist had a higher sensitivity for detecting microfilaria than skin snips from the shoulder, the difference presumably does not reflect a site preference. Thus, those who were positive in

both the waist and shoulder skin snippings had a relatively higher GMI (16 mf/skin snip) than those who were positive only for the waist skin snip. A low microfilaria level may lead to a higher level of negatives in skin snipping using corneo-scleral punch, just as low microfilaria levels may lead to more false negatives when using thick smears in detection of *W. bancrofti* microfilariae (46). This is buttressed by the scatterplot of the skin snip results from the waist and shoulder. This showed that those with mf intensities below 10 mf/snip had a more scattering tendency away from the regression line than those with higher intensities. Thus, there was a better agreement between results from the two skin snip sites when the mf intensities increased. The result was the same in the LIRB (19). Perhaps, 10 mf/snip is the critical threshold for parasitological survey methods involving corneo-scleral punch. This means that the method is not ideal for populations with low mf intensities.

## Acknowledgements

We are grateful to Okigwe LGA health officials, Community health assistants of Umuowaibu 1 (Emma Nwaimo), and Ndiroji (Esther), all the Ezes and chiefs of the two communities for their cooperation. The diligence and hard work of our field and laboratory workers: Edwin, Stephen, & Francis Uttah, Jacob, Vernai, Christopher Uzoma, Bassey Cobham, Comfort Nwankwoala, and Elder Akpan, as well as the medical doctors Dom Ibeh and Innocent Chukwuezi are appreciated. The authors declare that they have no conflicts of interest.

The work was funded by Danish International Development Assistance (DANIDA) through the Danish Bilharziasis Laboratory (now DBL-Institute for Health Research and Development), Denmark.

## References

1. World Health Organization. WHO Expert Committee on Onchocerciasis, Third report. *Technical Report Series*, 752; Geneva: WHO; 1987.
2. Remme JHF. The African Programme for Onchocerciasis Control: Preparing to Launch. *Parasitol Today*. 1995; 11: 403-406.
3. World Health Organization. World Report on Tropical Diseases; Geneva: WHO; 1990.
4. World Health Organization. Onchocerciasis. *Weekly Epidemiological Records*, 71; Geneva: WHO; 1996.
5. Anosike JC, Onwuliri CO. Studies on filariasis in Bauchi State, Nigeria. 1. Endemicity of onchocerciasis in Ningi Local Government Area. *Ann Trop Med Parasitol*. 1995; 89 (1): 31-38.
6. Akogun OB, Onwuliri COE. Hyperendemic onchocerciasis in The Taraba river valley of Gongola State (Old Adamawa Province) Nigeria. *Ann de Parasitol Hum et Comp*. 1991; 66: 22-26.
7. Crosskey RW. A review of *Simulium damnosum* s.l. and human onchocerciasis in Nigeria with special reference to geographical distribution and the development of the Nigerian National Control Campaign. *Tropenmed Parasitol*. 1981; 32: 2-16.
8. Udonsi JK. The status of human filariasis in relation to clinical signs in endemic areas of Niger Delta. *Ann Trop Med Parasitol*. 1986; 80: 425-432.
9. Nwoke BEB, Uwazie AU. Studies on the Blackflies, *Simulium* of Imo State, Nigeria: The distribution of immature stages in Isikwuato/Okigwe area. *Nig J Parasitol*. 1991; 12: 29-37.
10. Obikeze DS, Amazigo UO. Socio-cultural dimensions of onchocerciasis in a rural community in Nigeria, WHO/TDR/SER project report number 900353; Geneva: WHO; 1991.
11. Okonkwo P, Akpa A, Iheakwaba A, Nwagbo D, Umeh R, Adibua S, Ezike V, Ogbuokiri J. Studies on onchocerciasis in forest-savannah mosaic areas of Nigeria. 1. Investigations in Gbaragu, Oji River. *Ann Trop Med Parasitol*. 1991; 85: 617-623.
12. Amazigo U. (1993). Onchocerciasis and women's reproductive health: indigenous and biomedical concepts. *Trop Doct*. 1993; 23: 149-151.
13. Nwaorgu OC, Ohaedbulue A, Onweluzo IE, Alo ET, Nweke LN, Agu ML, Eme E. Results of a large scale onchocerciasis survey in Enugu State, Nigeria. *J Helminthol*. 1994; 68: 155-159.
14. Picq JJ, Coz J, Jardel JP. Une method d' evaluation des densites microfilariennes d' *Onchocerca volvulus*, Leuckart 1893, Chez des Onchocerciens technique et temps de lecture des biopsies cutanees. *Bull Wld Hlth Org*. 1971; 45: 517-520.
15. Pedersen EM, Kolstrup N. The epidemiology of Onchocerciasis in the Tukuyu Valley, South West Tanzania. *Trop Med Parasitol*. 1986; 37: 35-38.
16. Nwoke BEB. Studies on the field epidemiology of human onchocerciasis on the Jos Plateau, Nigeria [PhD. Dissertation]. University of Jos; 1986.
17. JICA, Manual of Onchocerciasis (Robles Disease) control in Guatemala. Tokyo: Publication of Japanese International Cooperative Agency; 1983.
18. Tada I, Iwamoto I, Wonde T. Quantitative studies on the emergence of *Onchocerca volvulus* microfilariae from the skin snips. *Jap J Trop Med Hyg*. 1973; 1: 13-24.
19. Uttah EC. Studies on the epidemiology of Filariasis in the Imo River Basin,

- Nigeria. [PhD Dissertation]. Danish Bilharziasis Laboratory Denmark/University of Port Harcourt; 1998.
20. Uttah EC. Clinical Manifestations of Mesoendemic Onchocerciasis in an Area with Multiple Filarial Species. *Ir J Parasitol.* 2009; 4(4): 19-28.
21. Udonsi JK. Filariasis in Igwun River [Basin, Nigeria: an epidemiological and clinical study with note on the vectors. *Ann Trop Med Parasitol.* 1988; 82: 75-82.
22. Uttah EC, Simonsen PE, Pedersen EM, Udonsi JK. Sporadic onchocerciasis in the Lower Imo River Basin, Nigeria. *Afr J Appl Zool Environ Biol.* 2004; 6: 76-85.
23. Sasa M. *Human Filariasis. A global survey of epidemiology and control.* Tokyo: University of Tokyo Press; 1976. p. 1- 468.
24. Anosike JC. Studies on filariasis in Bauchi State, Nigeria V. The distribution and prevalence of mansonellosis with special references to clinical signs. *Appl Parasitol.* 1994; 35, 189-192.
25. Wijeyaratne PM, Verma OP, Singha P, Osuhor PC, Motha B, Saha AL, Slotboom AB, DeLeon A, Bandipo AB. Epidemiology of filariasis in Malumfashi district of northern Nigeria. *Ind J Med Res.* 1982; 76: 534-544.
26. Hunponu-Wosu OO, Somorin AO. Onchocerciasis in Nigeria: observations on the geographical basis of the Endemicity in Lagos. *J Trop Med Hyg.* 1977; 80: 129-131.
27. Edungbola LD. Onchocerciasis control in Nigeria. *Parasitol Tod.* 1991; 7: 97-99.
28. Anosike JC, Onwuliri COE. A probable case of vertical transmission of *Onchocerca volvulus* microfilariae. *J Helminthol.* 1993; 67: 83-84.
29. Anosike JC, Onwuliri COE. Studies on filariasis in Bauchi State, Nigeria II. The prevalence of human filariasis in Darazo Local Government Area. *Appl Parasitol.* 1994; 35, 242-250.
30. Akogun OB, Ako JI, Hellandendu H. Onchocerciasis in Taraba State, Nigeria: Intensity, rate of infection and associated symptoms in 14 communities of Bali District. *Appl Parasitol.* 1992; 35: 125-132.
31. Akogun OB. Eye lesions, blindness and visual impairment in the Taraba river valley, Nigeria and their relation to onchocercal microfilariae in skin. *Act Trop.* 1992; 51: 143-149.
32. Alley ES, Plaiser AP, Boatın BA, Dadzie KY, Remone J, Zerobo G, Samba EM. The impact of five years of annual ivermectin treatment on skin microfilarial loads in the onchocerciasis focus of Asubende, Ghana. *Trans R. Soc Trop Med Hyg.* 1994; 88: 581-584.
33. Rodríguez-Pérez MA, Rodríguez MH, Margeli-Pérez H, Rivas-Alcalá AR. Effect of semiannual treatments of ivermectin on the prevalence and intensity of *Onchocerca volvulus* skin infection, ocular lesions, and infectivity of *Simulium ochraceum* populations in southern Mexico. *Am J Trop Med Hyg.* 1995; 52: 429-434.
34. Oyibo WA, Fagbenro-Beyioku AF. Reduced prevalence of onchocerciasis following mass treatment with Ivermectin. *E Afr Med J.* 1997; 74: 326-330.
35. World Health Organization. Onchocerciasis (river blindness). *Weekly Epidemiological Records*, 81; Geneva: WHO; 2006.
36. Osei-Atweneboana MY, Eng JK, Boakye DA, Gyapong JO, Prichard RK. Prevalence and intensity of *Onchocerca volvulus* infection and efficacy of Ivermectin in endemic communities in Ghana: a two-phase epidemiological study. *Lanc.* 2007; 369: 2021-2029.



37. Basanez MG, Pion SD, Boakes E, Filipe JA, Churcher TS, Boussinesq M. Effect of single-dose Ivermectin on *Onchocerca volvulus*: a systematic review and meta-analysis. *Lancet Inf Dis*. 2008; 8:310-322.
38. Gardon J, Boussinesq M, Kamgno J, Gardon-Wendel N, Demanga N, Duke BO. Effects of standard and high doses of Ivermectin on adult worms of *Onchocerca volvulus*: a randomized controlled trial. *Lancet*. 2002; 360: 203-210.
39. Plaisier AP, Alley ES, Boatn BA, Van Oortmarssen GJ, Remme H, De Vlas SJ, Bonneux L, Habbema JD. Irreversible effects of Ivermectin on adult parasites in onchocerciasis patients in the Onchocerciasis Control Programme in West Africa. *J Inf Dis*. 1995; 172: 204-210.
40. Klager S, Whitworth JA, Post RJ, Chavasse DC, Downham MD. How long do the effects of Ivermectin on adult *Onchocerca volvulus* persist? *Trop Med Parasitol*. 1993; 44: 305-310.
41. Chavasse DC, Post RJ, Lemoh PA, Whitworth JA. The effect of repeated doses of Ivermectin on adult female *Onchocerca volvulus* in Sierra Leone. *Trop Med Parasitol*. 1992; 43: 256-262.
42. Taylor MJ, Awadzi K, Basáñez M, Biritwum N, Boakye D, Boatn B, Bockarie M, Churcher TS, Debrah A, Edwards G, Hoerauf A, Mand S, Matthews G, Osei-Atweneboana M, Prichard RK, Wanji S, Adjei O. Onchocerciasis Control: Vision for the Future from a Ghanaian perspective. *Parasit Vect*. 2009; 2:7 doi:10.1186/1756-3305-2-7.
43. Awadzi K, Boakye DA, Edwards G, Opoku NO, Attah SK, Osei-Atweneboana MY, Lazdins-Helds, JK, Ardrey AE, Addy ET, Quartey BT. An investigation of persistent microfilaridermias despite multiple treatments with Ivermectin, in two onchocerciasis-endemic foci in Ghana. *Ann Trop Med Parasitol*. 2004; 98: 231-249.
44. Ali MM, Mukhtar MM, Baraka OZ, Homeida MM, Kheir MM, Mackenzie CD. Immunocompetence may be important in the effectiveness of Mectizan (Ivermectin) in the treatment of human onchocerciasis. *Act Trop*. 2002; 84: 49-53.
45. Lammie PJ, Hitch WL, Walker-Allen EM, Hightower AW, Eberhard ML. Maternal filarial infection as a risk factor for infection in children. *Lanc*. 1991; 337: 1005-1006.
46. Gubler DJ, Inui TS, Black HR, Bhattacharya NC. Comparisons of microfilaria density in blood sampled by finger prick, venipuncture, and ingestion by mosquitoes. *Am J Trop Med Hyg*. 1973; 22: 174-178.