



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iranian J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Original Article

Amphistome Species in Cattle in South Coast of Caspian Sea

*SZ Coskun¹, A Eslami², A Halajian³, A Nikpey⁴

¹Dept. of Parasitology, Faculty of Veterinary Medicine, University of Uludag, Turkey

²Dept. of Parasitology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

³Dept. Parasitology, Veterinary Faculty, Science and Research Branch, Islamic Azad University, Tehran, Iran

⁴Dept. of Biology, Basic Sciences Faculty, University of Guilan, Rasht, Iran

(Received 11 May 2011; accepted 18 Dec 2011)

ABSTRACT

Background: Knowledge about the amphistomid fauna in Iranian domestic ruminants depends on the studies conducted almost 30 years ago. The last situation in cattle is introduced here in the provinces in south coast of Caspian Sea.

Methods: Amphistomid species were collected from cattle at slaughter houses of the provinces Gilan and Mazandaran in 2010. Median sagittal sections were prepared by the conventional method. Species were identified by the histomorphological peculiarities of the muscular organs.

Results: Five amphistomid species, *Paramphistomum cevri*, *P. gotoi*, *Calicophoron calicophorum*, *Carmyerius spatiosus* and *Gastrothylax compressus*, recovered. *Calicophoron calicophorum* is a new species for Iran. Criteria used in identification of the species were illustrated.

Conclusion: Iran has a rich amphistomid fauna and mostly under the influence of oriental conditions.

Keywords: Parasitic Fauna, Amphistome, Cattle, Iran

*Corresponding author: Tel: +90 224 2941311, E-mail: scoskun@uludag.edu.tr

Introduction

The last comprehensive study on the amphistome species of Iranian ruminants was published almost 30 years ago by Sey and Eslami (1). In that study, 10 amphistomid species were described in ruminants in detail and illustrated as *Paramphistomum cervi*, *P. gotoi*, *P. gracile*, *P. Microbothrium* (now *Calicophoron microbothrium*), *Calicophoron papillosum*, *Gastrothylax crumenifer*, *G. compressus*, *Gigantocotyle explanatum* and *Orthocoelium scolicoelium*). Additionally, *Cotylophoron cotylophorum* and *Orthocoelium orthocoelium* appeared in the list given by Rafyi et al. (2). The present situation of amphistomid fauna of Iranian cattle in the provinces in south of Caspian Sea was determined in this study.

Materials and Methods

Amphistomid species were collected from cattle, at slaughter houses of the provinces Gilan and Mazandaran in south coast of Caspian sea in 2010. Median sagittal sections from the samples fixed in 70% alcohol were prepared by the conventional method. Histomorphological features, especially those of muscular organs such as pharynx, genital opening and acetabulum were examined under the light of the literatures (3-5). Most important criteria used in the identification of the species were illustrated.

Results

In the samples available five species recovered, including four species described by earlier authors. *Calicophoron calicophorum* was described for the first time in Iran.

Paramphistomum cervi Zeder, 1790

It has a *Liorchis* type of pharynx, *Gracile* type of genital opening and *Paramphisto-*

mum type of acetabulum (Fig. 1). This species is closely related to *P. gotoi* which is also found in Iran. It differs from *P. gotoi* by having smaller papillae found in the pharynx and by the position of the blind ceaca which do not meet dorso medially.

Paramphistomum gotoi Fukui, 1922

It has a *Liorchis* type of pharynx, *Gracile* type of genital opening and *Paramphistomum* type of acetabulum (Fig. 2). This species is closely related to preceding one. It differs from *P. cervi* by having larger papillae found in the pharynx and by the position of the blind ceaca which usually meet dorso medially.

Calicophoron calicophorum (Fischöeder, 1901) Näsmark, 1937

It is a new species for Iran. It has *Calicophoron* type of pharynx, genital opening and acetabulum (note the absence of dorsal and ventral exterior circular 2 muscle units) (Fig. 3). This species is found characteristically in oriental region. Iran is the most western distributional area for *C. calicophorum* in Asia.

Carmyerius spatiosus Brandes, 1898

It has *Gastrothylax* type of pharynx, *Gracile* type of genital opening and *Gastrothylax* type of acetabulum (Fig. 4). Ventral pouch is triangular near to circular with blunt angles.

Gastrothylax compressus Brandes, 1898

It has *Gastrothylax* type of pharynx, *Gracile* type of genital opening and *Carmyerius* type of acetabulum (Fig. 5). Ventral pouch usually triangular with apex dorsally directed. This species accepted as synonym of *G. crumenifer* for a long time. However, they differs easily by having different type of acetabulum.




Fig.1: *Liorchis* type of pharynx , *Gracile* type of genital opening and *Paramphistomum* type of acetabulum in *P. Cervi*




Fig. 2: *Liorchis* type of pharynx, *Gracile* type of genital opening and *Paramphistomum* type of acetabulum in *P. gotoi*




Fig. 3: *Calicophoron* type of pharynx, *Calicophoron* type of genital opening and *Calicophoron* type of acetabulum in *C. Calicophorum*




Fig. 4: *Gastrothylax* type of pharynx, *Gracile* type of genital opening and *Gastrothylax* type of acetabulum in *C. spatiosus*




Fig. 5: *Gastrothylax* type of pharynx, *Gracile* type of genital opening and *Carmyerius* type of acetabulum in *G. compressus*

Discussion

Findings of this study confirmed once more that Iran has a wide variety of amphistomid species as reported by previous researchers (1,2,4) and present authors. Five species have been identified in two provinces in the north of the country. Whereas, amphistomid fauna in ruminates in Turkey, the west neighbor country of Iran, consists only 3 species (*P. cervi*, *P. ichikawai*, *Calicophoron daubneyi*) as seen in Europe (6-8). Amphistomid fauna becomes richer in India, the east neighbor country (4, 9). *Paramphistomum cervi* and *P. gotoi* is wide spread in palaearctic region. Iran is the most west country where *C. calicophorum* and *G. compressus* recovered. *Carmyerius spatiosus* belongs to the Ethiopian region. South coast of the Caspian Sea is the most north point for this species (1). Its obviously seems that the amphistome fauna of Iran is strongly influenced by oriental and to a lesser extent by pale arctic and Ethiopian elements.

Distributional area of amphistomids is closely related to the climatic and microenvironmental conditions which affect the viability of the intermediate snail host. Further studies are necessary to elucidate the intermediate snail hosts of that species in the studied regions.

Acknowledgments

The authors declare that there is no conflict of interests.

References

1. Sey O, Eslami A. Review of amphistomes (Trematoda: Paramphistomata) of Iranian domestic ruminants. *Parasit Hung.* 1981-1982; 14: 61-65.
2. Rafyi A, Alavi A, Maghami G. Etat actuel de nos connaissances sur les Maladies parasitaires les plus importantes en Iran et Lutte contre ces Maladies. *Bull Off Int Epiz.* 1968; 69: 195-201.
3. Näsmark KE. A revision of the trematoda family Paramphistomidae. *Zool Bidr Upps.* 1937; 16: 301-565.
4. Eduardo SL. The taxonomy of the family Paramphistomidae Fiscoeder, 1901 with special reference to the morphology of species occurring in ruminants. Doctorate thesis, Univ London. 1980; 1-563.
5. Sey O. Revision of the family Gastrothylacidae Stiles et Goldberger, 1910 (Trematoda: Paramphistomata). *Acta Zool Acad Sci Hung.* 1983; 29: 223-252.
6. Coşkun ŞZ. Ruminantlarda *Paramphistomum* türlerinin bulunuş ve yayılışı. *DOĞA TU Vet ve Hay D.* 1988; 12:168-179.
7. Coşkun ŞZ. *Paramphistomum ichikawai* Fukui, 1922 (Trematoda: Paramphistomidae) in sheep in Turkey. *AÜ Vet Fak Derg.* 1988; 35:118-123.
8. Sey O. Revision of the Amphistomes of European ruminants. *Parasit Hung.* 1980; 13: 13-25.
9. Sey O. Examination of the validity and systematic position of some paramphistomids of indian ruminants. *Parasit Hung.* 1979; 12: 31-35.