



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Short Communication

Epidemiological and Clinical Characteristics of Patients with Hydatid Cysts in Khorasan Razavi Province, from 2011 to 2014

Salman KHAZAEI¹, Shahab REZAEIAN², Zaher KHAZAEI³, Elham GOODARZI³, Somayeh KHAZAEI⁴, Mahdi MOHAMMADIAN⁵, Hamid SALEHINIYA^{6,7}, Erfan AYUBI⁸, *Abdollah MOHAMMADIAN-HAFSHEJANI^{9,10}

1. Dept. of Epidemiology and Biostatistics, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran
2. Dept. of Epidemiology, School of Health, Shiraz University of Medical Sciences, Shiraz, Iran
3. Dept. of Epidemiology and Biostatistics, Kurdistan University of Medical Sciences, Kurdistan, Iran
4. Dept. of Paramedicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
5. Dept. of Social Medicine, School of Medicine, Dezfoul University of Medical Sciences, Dezfoul, Iran
6. Dept. of Public Health, School of Public Health, Zabol University of Medical Sciences, Zabol, Iran
7. Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, Iran
8. Dept. of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran
9. Social Determinants in Health Promotion Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
10. Dept. of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Received 14 Dec 2015

Accepted 06 Mar 2016

Keywords:

Human hydatidosis,
Echinococcosis,
Epidemiology,
Iran

*Correspondence Email:

amohamadii1361@gmail.com

Abstract

Background: This study aimed to investigate the epidemiological and clinical aspects of patients with hydatid cyst during 2011 to 2014.

Methods: This cross-sectional study was conducted in Khorasan Razavi Province, the Northeast of Iran, from 2011 to 2014. The study population was all cases with hydatid cyst who diagnosed in governmental and private laboratories, hospitals and health centers (HC) in Khorasan Razavi Province during 2011-14.

Results: The prevalence rate of hydatidosis was 1.44 per 100000 individuals. Of 357 cases, 54.9% were women, 40.3% rural, 45.8% housewives, and 3.4% were Afghan. The mean age of women was higher than that of men (39.13 ± 18.9 compared to 34.7 ± 17.9 yr, respectively, P -value=0.025). The highest proportion of cases (39.2%) was in the age group of 21-40 yr old. Abdominal pain was reported in 42.3% of cases. Liver involvement was the most common localization of hydatid cyst reported in 59.4% of patients, and 8.4% had multiple organ involvement. The common diagnosis methods of the disease were radiology (42.3%) followed by CT scan (37.8%). 45.9% of patients had domestic dog and hygiene principles of washing the vegetables was adhered by 6.7% of patients.

Conclusion: The prevalence of human hydatidosis, as a most important neglected disease, should be considered by health policy-makers in public health domain. In addition, educational programs to better recognition of the disease symptoms, and to identify the infection sources are needed in high risk group of population.

Introduction

Hydatid cyst disease is one of the most important zoonotic diseases, which occur, in larval stage of *Echinococcus granulosus* parasite. In this disease, canines such as dog, wolf, fox, and jackal (the final host) are infected by eating contaminated viscera, and small adult worms with size of 2-7 mm are formed in their intestines and begin to spawn. Canines repel the eggs of adult worms along with feces and infect the soil, water, and vegetables. Herbivores as the main intermediate host are infected by eating forage contaminated with the larvae of this parasite. Besides, human as the intermediate host is infected accidentally either by eating contaminated water and vegetables, or direct contact with contaminated dogs is infected to parasite egg of this disease; and hydatid cyst is usually involved the lung and liver in human (1, 2).

Epidemiological, clinical, and pathological symptoms of this disease depend on some factors such as age, gender, infection severity, cyst size, and the involved organ in the body (1, 2). This disease is common in most parts of the world especially in countries where animal husbandry is current. Contamination to this parasite is spreading in most tropical regions of the world and it is globally distributed in many countries such as Argentina, Peru, East Africa, Central Asia, China, and South America (3).

In endemic regions, the incidence rate of disease can be reached over 50 per 100,000 person-years with the prevalence rate of 5%-10% in human (3, 4). On the other hand, hydatidosis is among the most important human diseases with the high health and economic burden (5, 6). Iran is one of the areas that has been considered as hyper-endemic area by WHO in terms of close relationship of a high proportion of society with animals, traditional animal husbandry, and then contact with the sources of infection (7, 8). The prevalence

rate of this disease in the intermediate host has reported as 11.5%-34.6% (9), and the prevalence rate of human hydatid cyst has been reported from 1.1% to 13.7% in different parts of Iran (10-12).

Assessment of the epidemiological and clinical aspects of the disease can help health policy-makers to focus on the main public health problems in the community and to assess the efficacy of control and preventive programs to limit the spread and transmission of the parasite in humans.

Hence, the present study aimed to determine the epidemiological and clinical status of human hydatidosis in Khorasan Razavi Province during 2011-14.

Materials and Methods

This cross-sectional study was conducted in Khorasan Razavi Province, the Northeast of Iran, in 2014. The study population was all cases with hydatid cyst who diagnosed in governmental and private laboratories, hospitals and health centers in Khorasan Razavi Province during 2011-14.

The data collection was performed using a checklist including age, sex, nationality, place of residence, occupation, infected organ, the number of cysts, the type of diagnosis (CT scan, MRI, radiology, sonography, Immunofluorescence, other), clinical symptom, and having risk factors such as how to wash vegetables and contact with dog.

Descriptive statistics including frequency tables and diagrams as well as analytical statistics including Chi square and T-student tests were used for data analysis. All statistical analyses were performed at 0.05 significance levels using Stata software, version 12 (Stata Corp, College Station, TX, USA). In addition, mapping the percentage of hydatid cyst cases was created using GIS software by cities. For considering ethical issues in data collection, it

should be noted that all information of the patients has been confidential and the information has been recorded with code and without mentioning name.

Results

In total, 357 hydatid cyst cases had been registered in the health centers during the periods 2011-14 in all parts of Province. The prevalence of hydatidosis was 1.44 per 100000 individuals. Fig. 1 shows the percentage of hydatid cyst cases by county. The most percentage of the cases is related to the

northern cities. The most percentage of cases was related to the cities Mashhad (51.8%), Ghoochan (7.8%), Neishabour (7.0%), and Fariman (4.8%), respectively.

Demographic characteristics of studied patients by sex are shown in Table 1. Of 357 cases, 54.9% were women, 40.3% rural, 45.8% housewives, and 3.4% were Afghan. The mean age of women was higher than that of men (39.13 ± 18.9 compared to 34.7 ± 17.9 , respectively, P -value=0.025). About 21% of cases were younger than 20 years old. Farmers and ranchers were included the small number of cases.

- Torbatheidarieh 1
- Neishaboore 2
- Sabzevar 3
- Bardskan 4
- Kashmar 5
- Joghtai 6
- Jovin 7
- Khoshab 8
- Firoozeh 9
- Ghoochan 10
- Dargaz 11
- Chenaran 12
- Torghabeh 13
- shandiz 14
- Mashhad 15
- Kalat 16
- Sarakhs 17
- Fariman 18
- Torbatjam 19
- Zaveh 20
- Bakhazar 21
- Taibad 22
- Khavaf 23
- Mahvelat 24
- Rashtkhar 25
- Gonabad 26
- Bajestan 27
- Khalilabad 28




Fig. 1: Percentage of hydatid cyst cases by county in Khorasan Razavi Province in 2011-14

Table 1: Demographic characteristics of patients with hydatid cyst by sex in Khorasan Razavi Province in 2011-14

| Variable | Subgroup | Male (n=161) | Female (n=196) | P-value ^a |
|-----------|--------------------------|-----------------|-------------------|----------------------|
| Location | Urban | 87(54.0) | 126(64.3) | 0.050 |
| | Rural | 74(46.0) | 70(35.7) | |
| Age group | 0-20 | 35(21.7) | 40(20.4) | 0.209 |
| | 21-40 | 69(42.8) | 71(36.2) | |
| | 41-60 | 41(25.5) | 51(26.0) | |
| | >61 | 16(10.0) | 34(17.4) | |
| | Continuous ^b | 34.7±17.9 | 39.1±18.9 | 0.025 |
| Job | Employed ² | 2(1.2) | 4(2.1) | 0.001 |
| | Farmer ³ | 26(16.2) | 3(1.5) | |
| | Housekeeper ⁴ | 7(4.4) | 156(80.0) | |
| | Rancher ⁶ | 12(7.4) | 0(0.0) | |
| | Student ⁷ | 22(13.7) | 21(10.8) | |
| | Other ⁵ | 92(57.1) | 11(5.6) | |

*Data are presented as number (%). ^aStatistical analysis was done using Chi-square. ^b Comparing mean age was done using *t*-test.

The most common symptom of cases was related to the cities Mashhad (51.8%), Ghoochan (7.8%), Neishabour (7.0%), and Fariman (4.8%), respectively. Moreover, 45.9% of patients had domestic dog and hygiene principles of washing the vegetables was adhered by 6.7% of patients and 64.4% were washed the vegetables with water alone (Table 2).

The clinical characteristics of the patients are presented in Table 3. The majority of patients (70.5%) were infected with a single cyst, and 24.5% of them had multiple cysts. One hundred fifty one (42.3%) of patients were diagnosed by radiology method and 37.8% by CT scan method. The most common clinical symptom in the hydatidosis patients was abdominal pain (42.3%), after that hepatomegaly (27.4%), chest pain (21.6%), and cough (8.7%) were other common symptoms. Liver

involvement was reported in 59.4% of patients, and 8.4% had multiple organ involvement (Fig. 2).




Fig. 2: Status of organ involvement in the patients with hydatid cysts in Khorasan Razavi province during 2011-14

Table 2: Risk factors of hydatid cysts in the patients in Khorasan Razavi Province in 2011-14

| Variable | Subgroup | Number | Percent |
|--------------------------------|--------------------------------------|--------|---------|
| Contact with dog | Yes | 164 | 45.9 |
| | No | 193 | 54.1 |
| Consumption type of vegetables | Washing with water alone | 230 | 64.4 |
| | Washing with water and detergent | 87 | 24.4 |
| | Washing with water and disinfectants | 16 | 4.5 |
| | All three methods | 24 | 6.7 |

Discussion

The prevalence of human hydatidosis was 1.44 per 100,000 individuals in Khorasan Razavi Province. Previous studies conducted from different parts of Iran found the annual incidence of 1.33 in Hamadan (13), and 0.54 in Kermanshah (14) per 100,000 population. In addition, the rate of infection varied from 1.79% in Ardabil (11) to 13.8% in Khozestan (12).

Our findings showed that the highest proportion of hydatidosis (39.2%) in the age group of 21-40 years old. According to some factors such as geographical region, culture, and workforce; age can play an important role on the occurrence of the infection. In this regard, previous studies have reported the highest proportion of hydatidosis in the different age groups. For example, the age group of 60-69 in Isfahan (10), 40-49 year in Ardabil (11), and 30-60 year in Qom (15). Based on a review study, the range of 20-40 years old as the age group of the highest proportion has been reported (7).

The findings of this study showed the most percentage of infection in women than men (54.9% compared with 45.1%, respectively). This might be due to more contribution of women to agricultural and animal husbandry in this region. The high frequency of the disease in women than men is consistent with the obtained results of other Iranian studies (16, 17) and other countries (18, 19). However, some studies reported the higher rate of infection in men (10, 11, 15). Moreover, as stated in the previous studies conducted in Iran (7, 20), the housewives had the highest proportion of infection. It may be due to the more likely to contact with the source of contamination especially vegetables contaminated to parasitic eggs.

In our study, the highest proportion of hydatidosis was urbanite. There are some reasons for this finding. For example, the most percentage of the province's population

is urbanite and another might be a good-quality disease reporting. There are conflicting results in this regard; some studies reported the same result with our study (11, 15, 20, 21), and other reported inconsistent result (8, 22).

We found that abdominal pain was the main symptom with a proportion of 42.3% in the patients. Studies conducted in Turkey (23) and Iran (24) has reported a proportion of 77.6% and 66% for this symptom, respectively. In our study in line with previous reports (24, 25), liver was the most common localization of hydatid cyst. About 8.4% of the patients in this study had multiple organs involved and it was 12.3% in another study in Iran (24).

The common diagnosis methods of the disease were radiology (42.3%) followed by CT scan (37.8%). In a study to determine the pattern of hydatid cyst during 2000-10 in Tehran revealed that ultrasonography and CT scan were the most used imaging methods (24). Our findings demonstrate that these methods are the available tools for detecting hydatid cyst in Iran.

Concerning risk factors of infection with hydatidosis, our results found that 45.9% of individuals were in contact with dog, 64.4% of them have washed the vegetables with the water alone and 6.7% have washed the vegetables by hygiene principles of washing. Consequently, focus of improvement in education level of the society could be better. In Bulgaria (25) 90% of the affected children had sustained contact with dogs.

Conclusion

The prevalence of human hydatidosis, as a most important neglected disease, should be considered by health policy-makers in public health domain. In addition, educational programs to better recognition of the disease symptoms, and to identify the infection sources are needed in high risk group of population.

Acknowledgements

The authors would like to express their gratitude to the students who helped in gathering of data for the study. The authors declare that there is no conflict of interest.

References

- Zhang T, Zhao W, Yang D, Piao D, Huang S, Mi Y, et al. Human cystic echinococcosis in Heilongjiang Province, China: a retrospective study. *BMC Gastroenterol*. 2015;15:29.
- Budke CM, Carabin H, Ndimubanzi PC, Nguyen H, Rainwater E, Dickey M, et al. A systematic review of the literature on cystic echinococcosis frequency worldwide and its associated clinical manifestations. *Am J Trop Med Hyg*. 2013;88(6):1011-27.
- World Health Organization. Echinococcosis. Geneva: WHO; 2015 [cited 2015]; Available from: <http://www.who.int/mediacentre/factsheets/fs377/en/>.
- Craig PS, McManus DP, Lightowlers MW, Chabalgoity JA, Garcia HH, Gavidia CM, et al. Prevention and control of cystic echinococcosis. *Lancet Infect Dis*. 2007;7(6):385-94.
- Budke CM, Deplazes P, Torgerson PR. Global socioeconomic impact of cystic echinococcosis. *Emerg Infect Dis*. 2006;12(2):296-303.
- Schurer JM, Rafferty E, Farag M, Zeng W, Jenkins EJ. Echinococcosis: An Economic Evaluation of a Veterinary Public Health Intervention in Rural Canada. *PLoS Negl Trop Dis*. 2015;9(7):e0003883.
- Rokni MB. Echinococcosis /hydatidosis in Iran. *Iran J Parasitol*. 2009;4(2):1-16.
- Kalili B, Shahrani M, Moradi MT. Study of hydatid cyst in hospitalized patients with operation in Chaharmahal va Bakhtiary province (1988-2007). *J Shahrekord Univ Med Sci*. 2010;12(1):69-74.
- Yousofi H. Situation of hydatid cyst infection during last two decades (1985-2005) in Iran (Review of articles). *J Shahrekord Univ Med Sci*. 2008;10(1):78-88.
- Ilbeigi P, Mohebbali M, Kia EB, Saber-Inasab M, Aryaeipour M, Bizhani N, et al. Seroepidemiology of Human Hydatidosis Using AgB-ELISA Test in Isfahan City and Suburb Areas, Isfahan Province, Central Iran. *Iran J Public Health*. 2015;44(9):1219-24.
- Heidari Z, Mohebbali M, Zarei Z, Aryayipour M, Eshraghian M, Kia E, et al. Seroepidemiological study of human hydatidosis in meshkinshahr district, ardabil province, iran. *Iran J Parasitol*. 2011;6(3):19-25.
- Rafiei A, Hemadi A, Maraghi S, Kaikhaei B, Craig PS. Human cystic echinococcosis in nomads of south-west Islamic Republic of Iran. *East Mediterr Health J*. 2007;13(1):41-8.
- Ahmadi NA, Hamidi M. A retrospective analysis of human cystic echinococcosis in Hamedan province, an endemic region of Iran. *Ann Trop Med Parasitol*. 2008;102(7):603-9.
- Hamzavi Y, Vejdani M, Nazari N, Mikaeili A. The trend of hydatidosis in kermanshah province, Western iran (1986-2008). *Iran J Parasitol*. 2011;6(4):33-40.
- Rakhshanpour A, Harandi MF, Moazezi S, Rahimi M, Mohebbali M, Mowlavi G, et al. Seroprevalence of human hydatidosis using ELISA method in qom province, central iran. *Iran J Parasitol*. 2012;7(3):10-5.
- Harandi MF, Moazezi SS, Saba M, Grimm F, Kamyabi H, Sheikhzadeh F, et al. Sonographical and serological survey of human cystic echinococcosis and analysis of risk factors associated with seroconversion in rural communities of Kerman, Iran. *Zoonoses Public Health*. 2011;58(8):582-8.
- Amouian S, Tayebi N, Mohamadian Roshan N. A retrospective study of 1759 cases of Hydatid cyst in Mashad University hospitals. *Hakim Health Sys Res*. 2005;4(7):7-13.
- Yaghan RJ, Bani-Hani KE, Heis HA. The clinical and epidemiological features of hydatid disease in Northern Jordan. *Saudi Med J*. 2004;25(7):886-9.
- Ibrahim B, Haridy F, Hegazi M, Morsy T. Human hydatidosis granulosis in greater Cairo, Egypt: with general review. *J Egypt Soc Parasitol*. 2007;37(2):681-8.
- Rezaei F, Saghafipour A, Zia Sheikholeslami N, KaramiJooshi M. Investigation of demographic and clinical status of sufferers of hydatidcysts referred to hospitals affiliated to Qom University of Medical Science during a 12-Year

- Period (2002-2013), Iran. Qom Univ Med Sci J. 2014;8(5):63-7.
21. Asghari M, Mohebbali M, Kia EB, Farahnak A, Aryaeipour M, Asadian S, et al. Seroepidemiology of Human Hydatidosis Using AgB-ELISA Test in Arak, Central Iran. Iran J Public Health. 2013;42(4):391-6.
 22. Cobzaru RG, Dumitrescu AM, Ciobotaru M, Rîpă C, Leon M, Luca M, et al. Epidemiological aspects of hydatidosis in children, in some areas of north-eastern Romania. Rev Med Chir Soc Med Nat Iasi. 2013;117(3):754-7.
 23. Akcam AT, Ulku A, Koltas IS, Izol V, Bicer OS, Kilicbagir E, et al. Clinical characterization of unusual cystic echinococcosis in southern part of Turkey. Ann Saudi Med. 2014;34(6):508-16.
 24. Mousavi SR, Samsami M, Fallah M, Zirakzadeh H. A retrospective survey of human hydatidosis based on hospital records during the period of 10 years. J Parasit Dis. 2012;36(1):7-9.
 25. Jordanova DP, Harizanov RN, Kaftandjiev IT, Rainova IG, Kantardjiev TV. Cystic echinococcosis in Bulgaria 1996-2013, with emphasis on childhood infections. Eur J Clin Microbiol Infect Dis. 2015;34(7):1423-8.