



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Short Communication

Genetic Diversity of Hepatic/Non-Hepatic Cystic Echinococcosis in Baqiyatallah Hospital, Tehran, Iran

Hamidreza NEYSI¹, *Tahereh MOHAMMADZADEH^{1,2}, *Seyed Mahmoud SADJJADI³,
Jamal AKHAVANMOGHADDAM⁴, Alireza SHAMSAEI⁵

1. Department of Parasitology and Mycology, School of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran
2. Health Research Center, Life Style Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran
3. Department of Parasitology and Mycology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
4. Department of Surgery, School of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran
5. The Laboratory of Pathology, Baqiyatallah Hospital, Tehran, Iran

Received 15 Apr 2019
Accepted 14 Jun 2019

Keywords:

Genetic characterization;
Echinococcus granulosus;
Paraffin-embedded
tissue;
Iran

*Correspondence Email:

smsadjjadi@sums.ac.ir
yasint80@bmsu.ac.ir

Abstract

Background: Cystic echinococcosis (CE) is a worldwide zoonotic helminthic disease caused by the larval stage of *Echinococcus granulosus*. The infection is particularly important in terms of economic and medico-veterinary aspects in endemic areas including Iran. Considering the possibility of organ-tropism in *E. granulosus* strains, the present study was aimed to identify the genotypes of *E. granulosus* in different organs involved in patients, undergone surgery in Baqiyatallah Hospital, Tehran, Iran from 2005-2015.

Methods: Overall, 29 formalin-fixed paraffin-embedded tissues (FFPT) from patients with histologically confirmed CE including liver (N: 14) lungs (N: 6) abdomen (N: 2), pancreas (N: 2) and each of spleen, gallbladder and, muscles (N: 1) plus unknown organs (N: 2) were used and genetically characterized using polymerase chain reaction, followed by partial sequencing of mitochondrial cytochrome c oxidase gene subunit 1 (cox1) and analyzed.

Results: Nineteen out of 29 isolates including liver (N: 6) lungs (N: 4) abdomen (N: 2), pancreas (N: 2) and each of spleen, gallbladder, and muscle (N: 1), unknown organs (N: 2) obtained from paraffin-embedded blocks of human CE created an acceptable sequence in two directions. All 19 isolates regardless of the organ involved were recognized as *E. granulosus* sensu stricto (G1).

Conclusion: The sequence alignments of the isolates displayed two profiles. All sequenced samples showed *E. granulosus* sensu stricto (G1) with no organ-related genotype.

Introduction

Cystic echinococcosis (CE), is an important zoonotic parasitic disease with global distribution including Iran (1, 2). The infection is particularly important in terms of economic, medical and veterinary aspects in endemic areas (3). Human is usually infected following ingestion of parasite eggs via contaminated foods. The eggs develop to cysts in any organ especially in the liver and lungs (4). Diagnosis of hydatid disease is initially based on clinical signs followed by the imaging of suspected organs combined with serological tests (5-7). New techniques such as proteomics are also used (8). Diagnosis of the disease in human is difficult due to the absence of specific clinical signs; hence imaging methods may not be able to differentiate between hydatid cysts, tumors, and other lesions (9). Medical and surgical treatments are undertaken for the disease but surgery has been the most important and effective method for treating hydatid disease. However, the rupture of the cyst during surgery has been one of the major problems leading to the formation of secondary hydatid cysts in the abdominal cavity. Therefore, surgeons usually utilize several synthetic or natural protoscolicidal agents in treatment attempts to prevent this important complication (10-13).

Iran is an important region in the Middle East where various species of animals become infected by *E. granulosus* (2). Moreover, human CE is also documented in many studies in different parts of Iran (2, 6, 14,15).

During recent years, the molecular characterization of several organisms such as different parasites has been developed in many areas. In this context, genotype characterization of *E. granulosus* could help different aspects of the echinococcosis including epidemiology, diagnosis, treatments and control strategies of this important disease (4, 15).

Five genotypes of *E. granulosus* including *E. granulosus sensu stricto* (G1–G3 complex), *E. ortleppi* (G5), and *E. canadensis* (G6) have been reported from animal and/or human in Iran (4, 15-21).

Considering the possibility of organ-tropism in *E. granulosus* strains, the present study was aimed to identify the genotypes of *E. granulosus* in different organs of patients' undergone surgery in Baqiyatallah Hospital, Tehran, Iran.

Materials and Methods

Overall, 29 out of 104 formalin-fixed paraffin-embedded tissues (FFPT) including liver (N: 14) lungs (N: 6) abdomen (N: 2), pancreas (N: 2) and each from spleen, gallbladder and muscles (N: 1) and also unknown organ (N: 2) from patients (19 males and 10 females) with histologically confirmed CE obtained from Baqiyatallah Hospital, Tehran, Iran from 2005-2015.

After xylene de-paraffinization and methanol rehydration (100%, 90%, 80%, 70 % and 60%) of the tissue sections from all samples, DNA was extracted using Tissue DNA Extraction Micro Kit (Favorgen, Taiwan) according to the manufacturer's instructions. The quality of DNA was checked by NanoDrop (Biotek, USA, version: Epoch 2) and DNA was subsequently used as a template for polymerase chain reaction (PCR) using amplification of a partial mitochondrial DNA fragment of *cox1*. JB3 (TTTTTTGGG-CATCCTGAGGTTTAT) and JB4.5 (TAAA-GAAAGAACATAATGAAAATG) sequences were utilized as forward and reverse primers respectively (22). PCR was carried out in the final volume of 50 μ L, including 4 μ L (50-100 ng) of genomic DNA, 2 μ L (25 p. mol) of each primers and 25 μ L of master mix including Taq DNA polymerase, Mgcl₂, dNTP, PCR buffer and loading dye (Cinaclon, Iran) and 17 μ L of DDW under the following con-

ditions: 5 min at 94 °C as an initial denaturation step, followed by 35 cycles of 30 sec at 94 °C, 45 sec at 50 °C, 35 sec at 72 °C and a final extension step of 10 min at 72 °C. Negative (no added DNA) and positive controls were included in each PCR cycle. The amplification products were analyzed by electrophoresis in ethidium bromide-stained 1% agarose gel prepared in TAE buffered medium (65 mM Tris-HCl, 22.5 mM boric acid, 1.25 mM EDTA, pH 8.3) and subsequently visualized using an UV trans-illuminator (UVitec, Cambridge, UK).

A panel of 29 PCR amplicons for the *cox1* gene was purified using FavorPrep™ GEL/ PCR Purification Kit (Favorgen, Taiwan) and subjected to sequencing in two directions, using the same PCR primer set (First BASE Laboratories Sdn Bhd-604944X, Malaysia).

The sequences of the *cox1* gene were deposited in the GenBank database. Blast software was applied in order to preliminary identification and comparison of our sequences with other deposited ones in GenBank (<http://www.ncbi.nlm.nih.gov>). A Phylogenetic tree was drawn using our sequences and a few cases obtained from GenBank (Table 1). Alignment was carried out using ClustalW and the aligned sequences manually refined in BioEdit software (ver. 5.0.6) (23); maximum likelihood (ML) was inferred by MEGA 5 software for phylogenetic tree construction (24).

Approval of the study protocol was received from the Ethical Committee of Baqiyatallah University of Medical Sciences.

Results

Twenty-nine fragments of human CE isolates with around 450 bp long were successfully amplified within *cox1* gene (Fig. 1).




Fig. 1: Gel electrophoresis of *cox1*-PCR products. Lanes 1-4: *E. granulosis* isolates; Lane N: negative control; Lane P: positive control; Lane M, 100 bp DNA ladder

Liver and lungs were the most organs involved with 14 (48.27%) and 6 (20.68%) cases respectively. Other organs have been shown to be 9(31.04%) including two in abdomen (6.9%), two in pancreas (6.9%), one (3.45%) in gallbladder, one (3.45%) in muscle and one(3.45%) in heart. Furthermore, the organ of the two (6.9%) isolates have not been reported.

Nineteen out of 29 isolates including liver (N: 6) lungs (N: 4) abdomen (N: 2), pancreas (N: 2) and each of spleen, gallbladder and muscles (N: 1), and unknown organ (N: 2) obtained from paraffin embedded blocks of human CE resulted reasonable sequences in both directions. All 19 isolates regardless of their organs involved were recognized as *E. granulosis* sensu stricto (G1) (Table 1). The sequence alignments of the isolates displayed two profiles. Sequence profiles were deposited in GenBank with accession numbers MH244469-87. A consensus tree was made and the polymorphism points were shown (Fig. 2).

Table 1: The genotype of *Echinococcus granulosus* isolates identified by partial mitochondrial cox1 sequence in Baqiyatallah hospital (Tehran, Iran) and relevant information pertaining to the origins of sequences used for subsequent phylogenetic analyses (Fig. 2)

Code (No)	Identification number of isolates	Geographic origin Country (City or County)	Organ localization	Accession Number	Strain (Genotype)	References
1	6451	Iran (Tehran)	Liver	MH244469	G1	Present study
2	13910	Iran (Melard)	Unknown	MH244470	G1	Present study
3	14207	Iran (Tehran)	Liver	MH244471	G1	Present study
4	3455	Iraq (not reported)	Liver	MH244472	G1	Present study
5	4119	Iran (Rasht)	Lung	MH244473	G1	Present study
6	6493	Iran (Kermanshah)	Lung	MH244474	G1	Present study
7	3865	Iran (Karaj)	Lung	MH244475	G1	Present study
8	1611	Iran (not reported)	Pancreas	MH244476	G1	Present study
9	2875	Iran (not reported)	Pancreas	MH244477	G1	Present study
10	510	Iran (not reported)	Abdomen	MH244478	G1	Present study
11	5262	Iran (not reported)	Unknown	MH244479	G1	Present study
12	560	Iran (not reported)	Muscle	MH244480	G1	Present study
13	360	Iran (Golestan -Bandar-e-Gaz)	Liver	MH244481	G1	Present study
14	644	Iran (Khomein)	Lung	MH244482	G1	Present study
15	7840	Iran (Pakdasht)	Liver	MH244483	G1	Present study
16	6355	Iran (Tehran)	Abdomen	MH244484	G1	Present study
17	2703	Iran (Baghmalek)	Liver	MH244485	G1	Present study
18	603	Iran (Tehran)	Gallbladder	MH244486	G1	Present study
19	1583	Iran (Tehran)	Spleen	MH244487	G1	Present study
20	KU37	Iraq (Kurdistan)	Not reported	MF004309*	G1	(34)
21	EG44	Peru	Not reported	AB688621*	G1	(10)
22	Not reported	Australia (Tasmania)	Not reported	M84662*	G2	(22)
23	Not reported	India	Not reported	M84663*	G3	(22)
24	KH10	Iran (Birjand)	Liver	KP751430*	G6	(21)
25	(Out group)	China: Inner Mongolia	Not reported	AB461420*	<i>E.m</i>	(35)

*GenBank samples for comparison




Fig. 2: Genetic relationships of nineteen *Echinococcus granulosus* isolates from the Baqiyatallah hospital and selected GenBank sequence samples based on phylogenetic analysis of partial cox1 sequence data (Table 1). AB461420: *E. multilocularis* isolate from GenBank as the out-group

Discussion

Despite many efforts in control and prevention of CE, the disease is one of the most important zoonotic tissue helminthic infections in the world especially in Mediterranean regions (25).

Different aspects of the disease including the molecular perspective of human and animal CE is useful for control programs. Re-

cently many researchers have utilized molecular tools to access the information about parasite genotypic diversity (16-21, 26). Several genes including both coding and non-coding regions have used for intra-species differentiation; however, mitochondrial markers such as cox1 have shown great power in genetic discrimination (25-28).

The G6 genotype, which has already been reported as a main causative agent of brain CE

in Iran (16-17); is widespread in Birjand, Eastern Iran (21).

Application of formalin-fixed, paraffin-embedded tissue samples have been used for discrimination of genotypes in different localities to show the geographical distribution of the genotypes (29).

In this context, in the present study 29 human CE samples obtained from various organs were successfully amplified within *cox1* gene of which only 19 isolates including liver (31.58%) lungs (21.05%) abdomen (10.52%), pancreas (10.52%) and each of spleen, gallbladder and muscle (5.27%) plus unknown organ (10.52%) were shown an acceptable bi-directional sequences. All isolates regardless of the organ involved were recognized as *E. granulosus* sensu stricto (G1) which supports previous reports as G1 as the more prevalent strain in Iran (15, 18-19, 29,30).

A study on formalin-fixed, paraffin-embedded tissue samples from Alborz, Tehran, and Kerman provinces on 125 samples reported 56%, 40%, 3.2% and 0.8% of isolates as G1, G6, G3, and G2 genotypes, respectively. Despite the high number of samples, there is no report of the organ involved in that investigation (29).

A study on 334 liver and lungs CE, originated from sheep, cattle, goats and human by sequencing of ITS1 genes in Khuzestan province showed only G1 strain in all tested samples (30). However, another study on human brain CE showed them as G6 while other samples from the human liver were reported as G1 strain which showed organ tropism in *E. granulosus* (16).

Molecular study on 30 samples from hospitalized patients in Isfahan showed them as G1 which is similar to our results (31). A study on the paraffin embedded of CE in Mazandaran Province showed the human isolates as *E. granulosus* sensu stricto (32).

Overall, 47 human CE from two hospitals in Tehran were amplified based on *cox1* and *nad1* genes. Sixteen (53.3%), 13 (43.3%), and 1 (3.3%) samples were related to three organs

including lung, liver and spleen, respectively. Twenty-six and 3 of the 29 isolates were related to G1, and G3 strains, respectively. Only one G6 isolate was obtained from the lungs (33).

The lack of diversity in the isolates of our study could be because G1 is a common strain worldwide consequently, Iran and the number of samples although from different geographical origin (Tehran, Golestan, Kermanshah, Central provinces and even Iraq) is parallel to the most prevalent strain in those areas.

Conclusion

Genetic diversity of hepatic and non-hepatic cystic echinococcosis in Baqiyatallah Hospital, Tehran, during ten years showed that *E. granulosus* sensu stricto is predominant. This means that more attention should be directed to the diagnosis and treatment protocols to this important genotype in this geographic region.

Acknowledgements

The authors thank the "Clinical Research Development Center of Baqiyatallah Hospital" personnel for their kindly cooperation. Results presented in this article are part of the MS. thesis (Mr. Neysii) from the Baqiyatallah University of Medical Sciences, Tehran, Iran, (Ethical code: IR. BMSU.REC.1396.535). The authors appreciate the generous cooperation of Mrs. Haghghi, Mr. Aziz-Zadeh, Mr. Ebrahimi, Dr. Mahabadi, Dr. Esmaeili, Dr. Karamian, Dr. Davood Esmaeili and Mrs. Barzgar.

Conflict of interests

The author declares that there is no conflict of interest.

References

1. Deplazes P, Rinaldi L, Alvarez Rojas CA et al. Global Distribution of Alveolar and Cystic

- Echinococcosis. Adv Parasitol. 2017; 95:315-493.
2. Rokni M. Echinococcosis/hydatidosis in Iran. Iran J Parasitol. 2009; 4(2): 1-16.
 3. Fallah M, Taherkhani H, Sadjjadi M. Echinococcosis in stray dogs in Hamedan, west of Iran. Iran J Med Sci. 1995; 29:170-72.
 4. Alvarez Rojas CA, Romig T, Lightowlers MW. *Echinococcus granulosus* sensu lato genotypes infecting humans--review of current knowledge. Int J Parasitol. 2014; 44(1):9-18.
 5. Sadjjadi SM, Sedaghat F, Hosseini SV et al. Serum antigen and antibody detection in echinococcosis: application in serodiagnosis of human hydatidosis. Korean J Parasitol. 2009; 47(2):153-57.
 6. Mohammadzadeh T, Sako Y, Sadjjadi SM et al. Comparison of the usefulness of hydatid cyst fluid, native antigen B and recombinant antigen B8/1 for serological diagnosis of cystic echinococcosis. Trans R Soc Trop Med Hyg. 2012; 106(6):371-75.
 7. Khalilpour A, Sadjjadi SM, Moghadam ZK et al. Lateral flow test using *Echinococcus granulosus* native antigen B and comparison of IgG and IgG4 dipsticks for detection of human cystic echinococcosis. Am J Trop Med Hyg. 2014; 91(5):994-99.
 8. Sadjjadi FS, Rezaie-Tavirani M, Ahmadi NA et al. Proteome evaluation of human cystic echinococcosis sera using two dimensional gel electrophoresis. Gastroenterol Hepatol Bed Bench. 2018; 11(1):75-82.
 9. Babba H, Messedi A, Masmoudi S et al. Diagnosis of human hydatidosis - comparison between imagery and 6 serologic techniques. Am J Trop Med Hyg. 1994; 50(1): 64-68.
 10. Gökçe O, Gökçe C, Yilmaz M et al. Povidone iodine in experimental peritoneal hydatidosis. Br J Surg. 1991; 78(4):495-96.
 11. Sadjjadi SM, Zoharizadeh MR, Panjeshahin MR. In vitro screening of different *Allium sativum* extracts on hydatid cysts protoscoleces. J Invest Surg. 2008; 21(6):318-22.
 12. Hosseini SV, Ghanbarzadeh K, Barzin J et al. In vitro protoscolicidal effects of hypertonic glucose on protoscolices of hydatid cyst. Korean J Parasitol. 2006; 44(3):239-42.
 13. Tabatabaei ZS, Dehshahri S, Taghi MM et al. In vitro study on protoscolicidal effect of methanolic extract of *Allium birtifolium* on protoscoleces of cystic echinococcosis. Infect Disord Drug Targets. 2018;9 [Epub ahead of print].
 14. Sadjjadi SM, Ardehali S, Noman-Pour B et al. Diagnosis of cystic echinococcosis: ultrasound imaging or countercurrent immunoelectrophoresis? East Mediterr Health J. 2001; 7(6):907-11.
 15. Yanagida T, Mohammadzadeh T, Kamhawi S et al. Genetic polymorphisms of *Echinococcus granulosus* sensu stricto in the Middle East. Parasitol Int. 2012; 61(4):599-603.
 16. Sadjjadi SM, Mikaeili F, Karamian M et al. Evidence that the *Echinococcus granulosus* G6 genotype has an affinity for the brain in humans. Int J Parasitol. 2013; 43(11):875-77.
 17. Sadjjadi SM, Ebrahimipour M, Sadjjadi FS. Comparison between *Echinococcus granulosus* sensu stricto (G1) and *E. canadensis* (G6) mitochondrial genes (cox1 and nad1) and their related protein models using experimental and bioinformatics analysis. Comput Biol Chem. 2019;79 : 103-109.
 18. Gholami S, Jahandar H, Abastabar M et al. *Echinococcus granulosus* Sensu stricto in Dogs and Jackals from Caspian Sea Region, Northern Iran. Iran J Parasitol. 2016; 11(2):186-94.
 19. Hosseini-Safa A, Mohaghegh MA, Pestechian N et al. First report of Tasmanian sheep strain (G2) genotype isolated from Iranian goat using the high resolution melting (HRM) analysis. Gastroenterol Hepatol Bed Bench. 2016; 9(Suppl1):S70-S74.
 20. Ebrahimipour M, Sadjjadi S.M, Yousofi Darani H et al. Molecular Studies on Cystic Echinococcosis of camel (*Camelus dromedarius*) and report of *Echinococcus ortleppi* in Iran. Iran J Parasitol. 2017; 12(3):323-31.
 21. Karamian M, Haghighi F, Hemmati M et al. Heterogeneity of *Echinococcus canadensis* genotype 6 - the main causative agent of cystic echinococcosis in Birjand, Eastern Iran. Vet Parasitol. 2017; 245:78-85.
 22. Bowles J, Blair D, McManus DP (1992) Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. Mol Biochem Parasitol. 1992; 54(2):165-173.
 23. Hall, T. BioEdit version 5.0.6. 2001; <http://darwin.uvigo.es/dposada/repository/pdfs/BioEdit.help.pdf>

24. Tamura K, Peterson D, Peterson N, et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 2011; 28(10):2731-39.
25. Grosso G, Gruttadauria S, Biondi A, et al. Worldwide epidemiology of liver hydatidosis including the Mediterranean area. *World J Gastroenterol.* 2012; 18(13):1425-37.
26. Thompson RCA, McManus. Towards a taxonomic revision of the genus *Echinococcus*. *Trends Parasitol.* 2002; 18(10):452-57.
27. Nakao M, McManus D, Schantz P, et al. A molecular phylogeny of the genus *Echinococcus* inferred from complete mitochondrial genomes. *Parasitol.* 2007; 134(Pt 5):713-22.
28. Sharbatkhori M, Mirhendi H, Jex AR et al. Genetic categorization of *Echinococcus granulosus* from humans and herbivorous hosts in Iran using an integrated mutation scanning-phylogenetic approach. *Electrophoresis.* 2009; 30(15):2648-55.
29. Rostami S, Torbaghan SS, Dabiri S et al. Genetic characterization of *Echinococcus granulosus* from a large number of formalin-fixed, paraffin-embedded tissue samples of human isolates in Iran. *Am J Trop Med Hyg.* 2015; 92(3):588-94.
30. Khademvatan S, Yousefi E, Rafiei A, et al. Molecular characterization of livestock and human isolates of *Echinococcus granulosus* from south-west Iran. *J Helminthol.* 2013; 87(2):240-44.
31. Kia EB, Rahimi H, Sharbatkhori M et al. Genotype identification of human cystic echinococcosis in Isfahan, central Iran. *Parasitol Res.* 2010; 107(3):757-60.
32. Hedayati Z, Daryani A, Sarvi S et al. Molecular Genotyping of the Human Cystic Echinococcosis in Mazandaran Province, North of Iran. *Iran J Parasitol.* 2019; 14(1): 151–158.
33. Nikmanesh B, Mirhendi H, Ghalavand Z et al. Genotyping of *Echinococcus granulosus* Isolates from Human Clinical Samples Based on Sequencing of Mitochondrial Genes in Iran, Tehran. *Iran J Parasitol.* 2014; 9(1):20-27.
34. Hassan ZI, Meerkhan AA, Boufana B et al. Two haplotype clusters of *Echinococcus granulosus sensu stricto* in northern Iraq (Kurdistan region) support the hypothesis of a parasite cradle in the Middle East. *Acta Trop.* 2017; 172:201-207.
35. Nakao M, Xiao N, Okamoto M et al. Geographic pattern of genetic variation in the fox tapeworm *Echinococcus multilocularis*. *Parasitol Int.* 2009; 58(4):384-9.