



## ORIGINAL ARTICLE

## Ultrasonographic Assessment of Caudal Vena Cava to Aorta Ratio as a Novel Endpoint in Hemorrhagic Shock Resuscitation in Dogs

Reza Azargoun<sup>1\*</sup>, Reza Avizeh<sup>2</sup>, Alireza Ghadiri<sup>2</sup>, Hadi Imani Rastabi<sup>2</sup>, Mahdi Pourmahdi<sup>3</sup>

<sup>1</sup> Graduated of Small Animal Internal Medicine, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

<sup>2</sup> Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

<sup>3</sup> Department of Food Hygiene, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

Received: 19 March 2019  
Accepted: 11 August 2019  
Online: 19 August 2019

### Keywords:

Ultrasonography;  
Caudal vena cava to  
aorta ratio;  
Haemorrhagic shock;  
Dog.

### Abstract

**Objective-** The aim of this study was to assess ultrasonography-derived caudal vena cava to aorta ratio (CVC/Ao) as a novel endpoint in the resuscitation of experimental hemorrhagic shock in dogs.

**Design-** Experimental study.

**Animals-** Ten adult mongrel healthy dogs.

**Procedures-** After induction of anesthesia (control assessments), hemorrhagic shock was induced by blood withdrawal to a mean arterial pressure of 40 to 50 mmHg within 30 minutes and then maintained in a hypovolemic situation for an additional 30 minutes (second and third stages of assessments). Afterward, the dogs were randomly assigned to two groups which received 20 ml/kg lactated Ringer's solution or 5 ml/kg hydroxyethyl starch, in four consecutive 15 minutes intervals (fourth stage of assessments). One hour after the last resuscitation step, final ultrasonographic assessments were performed.

**Results-** Hemorrhagic shock caused a significant decrease in the CVC and Ao diameters as well as the CVC/Ao ( $2.14 \pm 0.28$  cm,  $0.85 \pm 0.07$  cm and  $0.4 \pm 0.06$ , respectively) ( $p < 0.05$ ). Following the fluid resuscitation CVC/Ao increased and returned to pre-shock values in both groups.

**Conclusion and clinical relevance-** Findings indicated that serial ultrasonographic assessment of the CVC/Ao can be a useful endpoint in the resuscitation of dogs with hemorrhagic shock.

\* Correspondence to: Reza Azargoun, Graduated of Small Animal Internal Medicine, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran. E-mail: [azargoun\\_vet@yahoo.com](mailto:azargoun_vet@yahoo.com)

## 1. Introduction SID

Hemorrhagic shock as the most common types of shock, is associated with morbidity and mortality in surgery and trauma patients.<sup>1</sup> Intravenous fluids therapy is the cornerstone of managing patients with hemorrhagic shock.<sup>2</sup> Reliable assessment of intravascular volume status during fluid resuscitation is crucial, because of either over- or under-resuscitation, can be perilous.<sup>3</sup> When to stop fluid resuscitation would ideally be directed by universally established endpoints based on documented evidence. Unfortunately, such endpoints do not yet exist in veterinary medicine, and time to cessation of fluid therapy is often based on clinical parameters and laboratory data that can be challenging.<sup>3,4</sup>

Ultrasonography has garnered attention as a rapid, repeatable, and generally reliable method for assessment of fluid status, particularly in the care of patients in shock.<sup>5-7</sup> Ultrasound may be used as an efficient tool in the identification of hypovolemia and to guide fluid management because it is safe, non-invasive, and applicable at the bedside.<sup>8</sup> One of the routine ultrasonographic method that usually used to assess intravascular volume status in adult humans is measurement of the inferior vena cava (IVC) diameter and comparing it with reference values. But, in order to eliminate the problems arising from various body weight and size in pediatric patients a new ultrasonographic variable has been introduced recently.<sup>6</sup>

The inferior vena cava/aorta diameter is a useful variable for evaluation of intravascular volume that can predict fluid responsiveness in critical care patients.<sup>9</sup> Similar to human subjects, there is also different dog sizes, therefore, use of an index of volemia such as those described for pediatric patients can be effective. Recent studies have shown that assessment of caudal vena cava to aorta ratio (CVC/Ao), as a marker of cardiac preload, is feasible method for estimation of hydration status in dogs,<sup>6,10</sup> However, to our knowledge, assessment of CVC/Ao in the

response to fluid therapy was not investigated previously. Therefore, the purpose of this study was to assess the CVC/Ao as a novel endpoint in the resuscitation phases of experimental induced hemorrhagic shock dogs with lactated Ringer's solution or 6% hydroxyethyl starch.

## 2. Materials and Methods

### *Animals*

Ten healthy male mongrel dogs, aged 1.5 to 3.5 years (weighing  $18.56 \pm 4.80$  kg) were experimentally studied. Animals were determined to be in good health based on clinical and laboratory parameters and excluded if they had a relevant cardiac disease determined by the echocardiographic and electrocardiographic examinations. Food was withheld from each dog for 12 hours prior to experiment, but had free access to water. No splenectomy was performed in this study.

### *Animal preparation and instrumentation*

After cannulation of left cephalic vein, with an 18-gauge catheter (for drugs and fluid administration), the anesthesia was induced with an intravenous bolus dose of propofol (Lipuro 1%, Braun, Melsungen, Germany) (6 mg/kg) and fentanyl (Caspian Tamin, Rasht, Iran) (5  $\mu$ g/kg).<sup>11</sup> Followed by intubation with an 8.0 to 8.5 mm cuffed endotracheal tube the animals were placed in left lateral recumbency and anesthesia was maintained with isoflurane (1.8%) in 100% oxygen.<sup>12</sup>

The left femoral artery was exposed by dissection and cannulated with a 16-gauge catheter, which was connected to a 3-way stopcock for induction of bleeding and to measure direct arterial blood pressure. Body temperature was maintained at 37° to 38°C with a heating blanket.<sup>13</sup> Also, an experienced investigator evaluated physical parameters including mucous membrane color, capillary refill time and peripheral pulse quality routinely at the end of each step.

## Experimental protocol

Ultrasonographic assessment of CVC/Ao was performed at the end of each step (A1 to A8) in five distinct stages as follows:

Control stage (A1): Baseline assessment was obtained after induction of anesthesia and instrumentation.

Hemorrhagic stage (A2): Each dog was then hemorrhaged to a mean arterial pressure (MAP) ranging from 40 to 50 mmHg.<sup>11</sup> The procedure lasted approximately 30 min and blood was collected into sterile empty blood bags.

Hypovolemic stage (A3): The animals were left in shock situation for an additional 30 min period during which no fluid was administered. If a physiologic compensatory mechanism developed and MAP increased above the purpose values, more blood was removed to restore the MAP back to 40 to 50 mmHg.

Resuscitation stage (A4–A7): Animals were then randomly allocated into two equal groups. Group A was resuscitated with lactated Ringer's solution (LR) (Iranian Parenteral and Pharmaceutical Co., Tehran, Iran) at 20 ml/kg in 15 min for four consecutive times. Group B was resuscitated with 6% Hydroxyethyl Starch (HES) (Voluven, Fresenius Kabi, Homburg, Germany) at 5 ml/kg in a similar manner to group A.<sup>14</sup>

Post-resuscitation stage (A8): The last assessment was performed one hour after termination of resuscitation. Then, animals were allowed to recover from anesthesia.

## Ultrasonographic technique

Ultrasonographic assessment was performed using commercially available ultrasound machine (Landwind Medical, Model: Mirror 2, China) with a 5–7 MHz microconvex probe.

Our ultrasonographic technique has been described in detail previously by Meneghini and coworkers.<sup>6</sup> In brief, the probe was located in a transverse position in the tenth to twelfth intercostal space (just few centimeters below to the spinal column), in order to find the porta hepatis. Upon

visualization of the CVC and Ao, the image was frozen and cine loops were recorded. A single veterinary radiologist performed all ultrasonographic assessments and CVC/Ao calculations. All values were calculated three times, to minimize errors in measurements.

## Statistical analysis

Data of this study were evaluated by repeated measures analysis of variance and LSD post hoc using SPSS 16.0 statistical software (SPSS, Inc., Chicago, USA). All data were presented as mean  $\pm$  standard deviation (SD). A *p* value of less than 0.05 was considered as statistically significant.

## 3. Results

The average blood volume loss was  $54 \pm 4$  ml/kg, which corresponded to approximately  $62\% \pm 4\%$  of the estimated circulating blood (88 ml/kg) volume; so that there were no significant differences between the groups ( $p > 0.05$ ). In both groups, the resuscitation were completed successfully with no mortality. There was no considerable difference between the studied groups according to MAP during all study stages ( $p > 0.05$ ). The mean time needed to ultrasonographic assessment after each step was  $53 \pm 10$  seconds.


Hemorrhagic shock led to significant reduction in CVC and Ao diameters from  $1.23 \pm 0.08$  and  $2.36 \pm 0.39$  cm, respectively, in A1 to  $0.85 \pm 0.07$  and  $2.14 \pm 0.28$  cm, respectively, in A2 ( $p < 0.05$ ); that increased after fluids administration (Table 1).

The data in Figure 1 represent the CVC/Ao changes in the studied stages. The mean  $\pm$  SD of CVC/Ao ratio in both groups before induction of shock was  $0.53 \pm 0.1$ , which was significantly reduced in A2 ( $0.04 \pm 0.06$ ) ( $p < 0.05$ ). Following the fluid therapy, this ratio increased and approximately returned to pre-shock values, at steps 4 and 7 in groups A and B, respectively. It is worth noting, the statistical analysis indicated no significant difference between solution types ( $p > 0.05$ ).

**Table 1.** Mean  $\pm$  SD values of CVC and Ao diameters in dogs with experimental hemorrhagic shock after resuscitation with LR and HES.

Factors	Caudal vena cava diameter (cm)		Aorta diameter (cm)	
	LR	HES	LR	HES
Steps				
A1	1.22 $\pm$ 0.07 <sup>a</sup>	1.25 $\pm$ 0.08 <sup>a</sup>	2.28 $\pm$ 0.28 <sup>a</sup>	2.44 $\pm$ 0.5 <sup>a</sup>
A2	0.85 $\pm$ 0.09 <sup>c</sup>	0.85 $\pm$ 0.06 <sup>d</sup>	2.11 $\pm$ 0.21 <sup>bc</sup>	2.16 $\pm$ 0.36 <sup>bc</sup>
A3	0.91 $\pm$ 0.09 <sup>c</sup>	0.89 $\pm$ 0.05 <sup>d</sup>	2.07 $\pm$ 0.24 <sup>c</sup>	2.10 $\pm$ 0.31 <sup>bc</sup>
A4	1.16 $\pm$ 0.08 <sup>b</sup>	0.99 $\pm$ 0.07 <sup>d</sup>	2.15 $\pm$ 0.25 <sup>b</sup>	1.96 $\pm$ 0.17 <sup>c</sup>
A5	1.23 $\pm$ 0.07 <sup>a</sup>	1.05 $\pm$ 0.06 <sup>c</sup>	2.24 $\pm$ 0.2 <sup>a</sup>	2.28 $\pm$ 0.38 <sup>bc</sup>
A6	1.29 $\pm$ 0.06 <sup>a</sup>	1.13 $\pm$ 0.07 <sup>b</sup>	2.3 $\pm$ 0.17 <sup>a</sup>	2.39 $\pm$ 0.42 <sup>b</sup>
A7	1.35 $\pm$ 0.04 <sup>a</sup>	1.26 $\pm$ 0.08 <sup>a</sup>	2.36 $\pm$ 0.18 <sup>a</sup>	2.48 $\pm$ 0.48 <sup>a</sup>
A8	1.24 $\pm$ 0.04 <sup>a</sup>	1.3 $\pm$ 0.07 <sup>a</sup>	2.21 $\pm$ 0.25 <sup>a</sup>	2.43 $\pm$ 0.46 <sup>a</sup>

Note: Different lowercase in each column denote significant differences ( $p < 0.05$ ).



**Figure 1.** Mean  $\pm$  SD values of CVC/AO in dogs with experimental hemorrhagic shock after resuscitation with LR and HES.

#### 4. Discussion

In this study, after induction of severe hemorrhagic shock, animals were resuscitated with LR and HES as the most frequently used crystalloid and synthetic colloid fluids in veterinary medicine.

The endpoint of resuscitation occurs when a patient has been fluid resuscitated adequately. Traditionally, resuscitation endpoints including restoration of normal clinical perfusion parameters (i.e., heart rate, capillary refill time, peripheral pulse quality, and rectal temperature), blood pressure, and urine output remain the standard of care. However, reliance on these traditional endpoints of resuscitation, due to body's compensatory mechanisms for initial volume loss, may not provide a reliable enough estimate of volume status and occult shock can occur

eventually.<sup>2,15</sup> For example, blood pressure measurement, as the main marker in evaluating emergency situations, is often used to gauge the adequacy of resuscitation.<sup>16</sup> Interestingly, Wo et al. demonstrated that blood pressure is an unreliable indicator of acute blood loss and accordingly is poor marker of the severity of the shock as well as response to treatment.<sup>17</sup>

The CVC-to-Ao ratio have been demonstrated to be strongly correlated with hemodynamic parameters and the CVC diameter has been proposed to reflect volume status more closely than other variables based on the arterial system, such as blood pressure, pulse rate or quality, and Ao diameter.<sup>3,6</sup> Furthermore, Akilli *et al.* proved that ultrasonographic measurement of the IVC diameter, as a marker of early hemorrhagic shock, is more accurate than the shock index and other routinely used non-invasive predictors of acute blood loss such as heart rate, serum lactate level, and base deficit.<sup>18</sup> In another study, IVC diameter strongly reflect the central venous pressure which is currently considered the most precise method in assessing patients' fluid status.<sup>19,20</sup>

In our study, as expected, hypovolemia caused by hemorrhagic shock led to decrease in CVC diameter. However, in contrast to the recent study by Cambournac *et al.*, Ao diameter also showed a significant decrease after blood loss which can be attributed to severe hypovolemia, than milder one that was reported.<sup>3</sup> As intravenous fluids were administered, the CVC diameter increased a little more than the aorta. This may be due to differences in vessel characteristics. Histologically, the CVC has a thinner wall than the aorta, with less defined layers, as well as decreased number of smooth myocytes. In addition, veins have less elasticity than arteries. The venous system acts as a fluid reservoir that contains 65–75% of total blood volume, with the most contribution created by splanchnic venous system.<sup>21</sup>

In light of the mentioned issues, assessment of CVC/Ao may be a reasonable method to monitor the response to fluid resuscitation. In our experiment, significantly

hypovolemic dogs showed a decrease in mean value of CVC/Ao, that was lower than the results of another study that evaluate CVC/Ao after a blood donation of  $9.8 \pm 2.2$  mL/kg (0.4 versus 1.01).<sup>3</sup> The CVC/Ao increased slightly after fluid resuscitation and almost returned to control values that can indicate successful resuscitation in both groups. In normovolemic dogs that were admitted for surgical repair of skin wounds, the mean value of CVC/Ao after bolus administration of Hartmann solution ( $14 \pm 7$  mL/kg) was 0.91 that was higher than our results.<sup>6</sup> The difference in the results may be associated with the differences in anesthetic protocols, intravascular volume status, fluid therapy methods, and ultrasonographic techniques.

The lack of significant difference in CVC/Ao values between the two types of solutions that were used, despite different dosage, probably is due to the discrepancy in mechanisms of action as well as the stability time in the vessels of infused fluids.<sup>13</sup>

Ultrasonographic assessment of CVC/Ao to guide fluid resuscitation also makes it possible to assess abdominal (liver, spleen, kidneys) and thoracic organs (heart, lungs) in daily practice. However, Ultrasound has some disadvantages. The results are operator dependent, and the image quality is affected by anatomical obstacles such as large fat deposits and excessive abdominal gas.<sup>22</sup> According to Tan et al. study, some situations including right ventricular dysfunction, tricuspid valve disease, liver cirrhosis, tension pneumothorax, asthma, pericardial tamponade, increasing age, and ethnicity of patients can affect the IVC diameter and making it less reliable.<sup>23</sup>

There were some limitations in our study. Firstly, the dogs were under anesthesia and it alters hemodynamic variables by reducing cardiac contractility, vascular tonicity, and changing cardiovascular responses to variation in blood pressure and volume.<sup>21</sup> However, in the study reported here, anesthesia was maintained using isoflurane to restrict potential confounding hemodynamic effects.<sup>24</sup> Secondly, the animal position may affect the CVC diameter.

Although not investigated in dogs, but it is demonstrated that the position of the human patients can influence IVC diameter. So that, left lateral recumbency making the smallest diameter and right lateral recumbency making the largest IVC diameter.<sup>25</sup> Finally, ultrasonographic assessments were performed by a single operator and involving more operators could improve the validity of the results.

In conclusion, ultrasonographic assessment of CVC/Ao can provide an affordable and simple method for estimation the resuscitation endpoint beside a comprehensive physical examination in dogs with hemorrhagic shock. Further studies are now needed to establish reference intervals for CVC/Ao, in order to guide the resuscitation of referral dogs in daily practice that have uncertain pre-shock values.

### Acknowledgment

This survey was financially supported by the Research Council of Veterinary Faculty, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

### Conflict of interest

The authors declare no conflict of interest.

### References

1. Fülöp A, Turóczy Z, Garbaisz D, Harsányi L, Sziártó A. Experimental models of hemorrhagic shock: A review. *European Surgical Research*, 2013; 50: 57-70.
2. Sawe HR, Haeffele C, Mfinanga JA, Mwafongo VG, Reynolds TA. Predicting fluid responsiveness using bedside ultrasound measurements of the inferior vena cava and physician gestalt in the emergency department of an urban public hospital in Sub-Saharan Africa. *PLOS One*, 2016; 11: e0162772.
3. Cambournac M, Goy-Thollot I, Violé A, Boisvineau C, Pouzot-Nevoret C, Barthélemy A. Sonographic assessment of volaemia: development and validation of a new method in dogs. *Journal of Small Animal Practice*, 2018; 59: 174-182.



4. Keefe J. Shock and Initial Stabilization. In: Norkus C, Ed. *Veterinary Technician's Manual for Small Animal Emergency and Critical Care*. 1st ed. Iowa: Wiley-Blackwell, 2012; 25-43.
5. Seif D, Mailhot T, Perera P, Mandavia D. Caval sonography in shock: a noninvasive method for evaluating intravascular volume in critically ill patients. *Journal of Ultrasound in Medicine*, 2012; 31: 1885-1890.
6. Meneghini C, Rabozzi R, Franci P. Correlation of the ratio of caudal vena cava diameter and aorta diameter with systolic pressure variation in anesthetized dogs. *American Journal of Veterinary Research*, 2016; 77: 137-143.
7. Lee CW, Kory PD, Arntfield RT. Development of a fluid resuscitation protocol using inferior vena cava and lung ultrasound. *Journal of Critical Care*, 2016; 31: 96-100.
8. Zengin S, Al B, Genc S, Yildirim C, Ercan S, Dogan M, Altunbas G. Role of inferior vena cava and right ventricular diameter in assessment of volume status: a comparative study. *American Journal of Emergency Medicine*, 2013; 31: 763-767.
9. Darnis E, Boysen S, Merveille AC, Desquilbet L, Chalhoub S, Gommeren K. Establishment of reference values of the caudal vena cava by fast-ultrasonography through different views in healthy dogs. *Journal of Veterinary Internal Medicine*, 2018; 32: 1308-1318.
10. Kwak J, Yoon H, Kim J, Kim M, Eom K. Ultrasonographic measurement of caudal vena cava to aorta ratios for determination of volume depletion in normal beagle dogs. *Veterinary Radiology & Ultrasound*, 2018; 59: 203-211.
11. Braz JR, do Nascimento P Jr, Paiva Filho O, Braz LG, Vane LA, Vianna PT, Rodrigues GR Jr. The early systemic and gastrointestinal oxygenation effects of hemorrhagic shock resuscitation with hypertonic saline and hypertonic saline 6% dextran-70: A comparative study in dogs. *Anesthesia & Analgesia*, 2004; 99: 536-546.
12. Dyson D, Sinclair M. Impact of dopamine or dobutamine infusions on cardiovascular variables after rapid blood loss and volume replacement during isoflurane-induced anesthesia in dogs. *American Journal of Veterinary Research*, 2006; 67: 1121-1130.
13. Nascimento P Jr, de Paiva Filho O, de Carvalho LR, Braz JR. Early hemodynamic and renal effects of hemorrhagic shock resuscitation with lactated ringer's solution, hydroxyethyl starch, and hypertonic saline with or without 6% dextran-70. *Journal of Surgical Research*, 2006; 136: 98-105.
14. Mazzaferro E, Powell LL. Fluid Therapy for the Emergent Small Animal Patient. *Veterinary Clinics of North America: Small Animal Practice*, 2013; 43: 721-734.
15. Dipti A, Soucy Z, Surana A, Chandra S. Role of inferior vena cava diameter in assessment of volume status: a meta-analysis. *American Journal of Emergency Medicine*, 2012; 30: 1414-1419.
16. Lyon M, Blaivas M, Brannam L. Sonographic measurement of the inferior vena cava as a marker of blood loss. *American Journal of Emergency Medicine*, 2005; 23: 45-50.
17. Wo CC, Shoemaker WC, Appel PL, Bishop MH, Kram HB, Hardin E. Unreliability of blood pressure and heart rate to evaluate cardiac output in emergency resuscitation and critical illness. *Critical Care Medicine*, 1993; 21: 218-223.
18. Akilli B, Bayir A, Kara F, Ak A, Cander B. Inferior vena cava diameter as a marker of early hemorrhagic shock: a comparative study. *Turkish Journal of Trauma and Emergency Surgery*, 2010; 16: 113-118.
19. Marcelino P, Borba A, Fernandes AP, Marum S, Germano N, Lopes MR. Non invasive evaluation of central venous pressure using echocardiography in the intensive care-specific features of patients with right ventricular enlargement and chronic exacerbated pulmonary disease. *Revista Portuguesa de Pneumologia*, 2006; 12: 637-658.
20. Durajska K, Januszkiewicz E, Szmygel Ł, Kosiak W. Inferior vena cava/aorta diameter index in the assessment of the body fluid status - a comparative study of measurements performed by experienced and inexperienced examiners in a group of young adults. *Journal of Ultrasonography*, 2014; 14: 273-279.
21. Nelson NC, Drost WT, Lerche P, Bonagura JD. Noninvasive estimation of central venous pressure in anesthetized dogs by measurement of hepatic venous blood flow velocity and abdominal venous diameter. *Veterinary Radiology & Ultrasound*, 2010; 51: 313-323.
22. Karacabey S, Sanri E, Guneysel O. A non-invasive method for assessment of intravascular fluid status: Inferior vena cava diameters and collapsibility index. *Pakistan Journal of Medical Sciences*, 2016; 32: 836-840.
23. Tan HL, Wijeweera O, Onigkei J. Inferior vena cava guided fluid resuscitation – Fact or fiction? *Trends in Anaesthesia and Critical Care*, 2015; 5: 70-75.

24. Cheung AT, Driessen B, Jahr JS, Duong PL, Ramanujam S, Chen PC, Gunther RA. Blood substitute resuscitation as a treatment modality for moderate hypovolemia. *Artificial Cells Blood Substitutes and Biotechnology*, 2004; 32: 189-207.

25. Ciozda W, Kedan I, Kehl DW, Zimmer R, Khandwalla R, Kimchi A. The efficacy of sonographic measurement of inferior vena cava diameter as an estimate of central venous pressure. *Cardiovascular Ultrasound*, 2016; 14: 33.

## چکیده

ارزیابی اولتراسونوگرافی نسبت بزرگ سیاهرگ پسین به آئورت به عنوان نقطه جدید در احیای شوک هموراژیک در سگ

رضا آذرگون<sup>۱</sup>، رضا آویزه<sup>۲</sup>، علیرضا غدیری<sup>۳</sup>، هادی ایمانی راستابی<sup>۴</sup>، مهدی پورمهدی<sup>۳</sup>

<sup>۱</sup> دانش‌آموخته طب داخلی دام‌های کوچک، دانشکده دامپزشکی، دانشگاه شهید چمران اهواز، اهواز، ایران.

<sup>۲</sup> گروه علوم درمانگاهی، دانشکده دامپزشکی، دانشگاه شهید چمران اهواز، اهواز، ایران.

<sup>۳</sup> گروه بهداشت مواد غذایی، دانشکده دامپزشکی، دانشگاه شهید چمران اهواز، اهواز، ایران.

**هدف-** هدف از این مطالعه ارزیابی اولتراسونوگرافی نسبت بزرگ سیاهرگ پسین به آئورت به عنوان یک نقطه جدید در خاتمه احیای سگ‌های دچار شوک هموراژیک تجربی است.

**طرح مطالعه-** مطالعه تجربی.

**حیوانات-** ده قلاده سگ بالغ سالم از نژاد مخلوط.

**روش کار-** پس از القای بیهوشی (ارزیابی‌های کنترل)، شوک هموراژیک از طریق خون‌گیری تا رسیدن به فشار متوسط سرخرگی ۴۰ تا ۵۰ میلی‌متر جیوه در عرض ۳۰ دقیقه ایجاد و سپس در وضعیت هیپوولمی به مدت ۳۰ دقیقه حفظ شد (مراحل دوم و سوم ارزیابی). پس از آن، سگ‌ها به‌طور تصادفی به دو گروه تقسیم شدند و محلول رینگر لاکتات با دوز ۲۰ میلی‌لیتر به ازای هر کیلوگرم وزن یا هیدروکسی‌اتیل استارچ با دوز ۵ میلی‌لیتر به ازای هر کیلوگرم وزن را در چهار دوره متوالی ۱۵ دقیقه‌ای دریافت کردند (مرحله چهارم ارزیابی). یک ساعت پس از آخرین مرحله احیاء، ارزیابی نهایی اولتراسونوگرافی انجام گرفت.

**نتایج-** شوک هموراژیک موجب کاهش معنادار در قطر بزرگ سیاهرگ پسین و آئورت و همچنین نسبت بزرگ سیاهرگ پسین به آئورت (به ترتیب ۲/۱۴±۰/۲۸ سانتیمتر، ۰/۸۵±۰/۰۷ سانتیمتر و ۰/۴±۰/۰۶) گردید ( $p < 0/05$ ). پس از احیاء نسبت بزرگ سیاهرگ پسین به آئورت افزایش یافته و به مقادیر پیش از شوک در هر دو گروه بازگشت.

**نتیجه‌گیری و کاربرد بالینی-** یافته‌ها نشان داد که ارزیابی پیاپی اولتراسونوگرافی نسبت بزرگ سیاهرگ پسین به آئورت می‌تواند نقطه پایانی مفید در احیای سگ‌های دچار شوک هموراژیک باشد.

**واژه‌های کلیدی-** اولتراسونوگرافی، نسبت بزرگ سیاهرگ پسین به آئورت، شوک هموراژیک، سگ.