

A Case of Acrorenal Syndrome

Mehdi Esmaeili Jadidi,¹ Farshid Oliaei^{2,3,4}

¹Student Research Committee, Babol University of Medical Science, Babol, Iran

²Cellular and Molecular Biology Research Center; Cancer Research Center; Health Research Institute, Babol University of Medical Sciences, Babol, Iran

³Clinical Research Development Center, Shahid Beheshti Hospital, Babol University of Medical Sciences, Babol, Iran

⁴Department of Internal Medicine, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Urinary tract anomalies are common and comprise about 20% to 30% of total congenital anomalies. This spectrum consists of many different anomalies of the urinary tract that may be syndromic or nonsyndromic with different etiologies. In this case report, a patient with single kidney and urinary tract signs is introduced that was diagnosed accidentally. The finding of different anomalies in different organ systems should lead us to examination of the intactness of the urinary tract. In these disorders, if there is no need for immediate intervention, long-term follow-up can be helpful to postpone chronic kidney disease progression.

IJKD 2018;12:247-9
www.ijkd.org

Keywords. congenital anomalies, urinary tract, acrorenal syndrome

INTRODUCTION

Urinary tract anomalies are common and comprise about 20% to 30% of total congenital anomalies.¹ This spectrum consists of unilateral or bilateral renal agenesis, hypoplasia, dysplasia, multicystic kidneys, horseshoe or duplex kidneys, vesicoureteral reflux, hydroureter, hydronephrosis, and ureteropelvic or ureterovesical junction obstruction.^{2,3}

Congenital renal anomalies can be sporadic, familial, syndromic (with disorders in other organs), or nonsyndromic. The etiology can be environmental (nutrition and drugs) or genetic.^{4,5} In this case report, a single-kidney patient with urinary tract signs is introduced that was diagnosed with a syndrome.

CASE REPORT

A 29-year-old man was referred to hospital with severe left flank pain and reduced urine output. On physical examination, in addition to left costovertebral angle tenderness, loss of right thumb was prominent as a congenital anomaly

(Figure). Abdominal ultrasonography represented a single left kidney with moderate hydronephrosis and a small lower calyceal calculus.

The patient was hydrated and then he passed a calculus that was lost. The history of urinary calculus was positive in his father. He was azotemic that recovered completely after calculus passage. The patient was unaware about his single kidney and there was no similar history in his family. Repeated history taking revealed right undescended testis that had been surgically corrected in childhood. In order to examine other anomalies associated with acrorenal syndrome, vesicocystoureterography was done that was negative. He was discharged with a recommendation for follow up of the left kidney function.

DISCUSSION

Acrorenal syndrome is part of many different syndromes occurring with congenital anomalies of the kidney and urinary tract, and it can be due to chromosomal, teratogenic, or dysplastic disorders.⁶ In this syndrome, limb disorders can be unilateral or



Radiographic image of the right hand shows oligodactyly (no thumb).

bilateral; upper or lower; polydactyly, oligodactyly, syndactyly, brachydactyly, or ectrodactyly (loss of a part or whole of a finger like this case); and shortness of radial bone or forearm. This problem can be seen in a wide range of disorders of oromandibular area, trachea, lungs, skin and appendages, uterus, testes and vas deferens, heart, eyes, and spine. Simultaneous formation of the urinary tract and other systems in 4 to 12 weeks of embryonic period is the main reason of these syndromes.⁷

It seems that many acrorenal syndromes are transmitted as autosomal recessive inheritance, but the reported case did not match with syndromes like Dieker,^{8,9} Johnson-Munson,¹⁰ or Siegler.¹¹ In these syndromes, the probable gene (Formin gene) is located at 15q13-14.¹¹ In reported case, due to no similar patient in his family, it must be

a sporadic mutation.

Unilateral renal agenesis occurs in 1 in 1500 to 3200 live births. This is more common in men and more prevalent in the left side.¹² Regardless of morbidities from other nonurinary system involvement and their impact on the quality of life, unilateral renal agenesis alone with making hyperfiltration, glomerulosclerosis, and chronic kidney disease can be impressive on life.

The finding of different anomalies in different organ systems should lead us to examination of the intactness of the urinary tract. In these disorders, if there is no need for immediate intervention, long-term follow-up can be helpful to postpone chronic kidney disease progression.

ACKNOWLEDGEMENTS

We thank Miss Mehdinia for typing and editing of the paper and the Clinical Research Development Center staff of Shahid Beheshti Hospital, Babol, for their cooperation.

CONFLICT OF INTEREST

None declared.

REFERENCES

1. Hakan RT, Okan T, Ali H et al . Congenital Anomalies of kidney and Urinary tract. *Semin Nephrol.* 2010;30:374-86.
2. Piscione TD, Rosenblum N. The molecular control of renal branching morphogenesis: current knowledge and emerging insights. *Differentiation.* 2002;70:227-46.
3. Zaffanello M, Brugnara M, Zuffante M, et al. Are children with congenital solitary kidney at risk for life long complications? A lack of prediction demands caution. *Int Urol Nephrol.* 2009;41:127-35.
4. Weber S, Moriniere V, Knuppel T et al. Prevalence of mutations in renal developmental genes in children with renal hypodysplasia: results of the Escap study. *J Am Soc Nephrol.* 2006;17:2864-70.
5. Nielson GL, Norgard B, Puho E, et al. Risk of specific congenital abnormalities in offspring of women with diabetes. *Diabet Med.* 2005;22:693-6.
6. Rascher W, Rosch WH. Congenital abnormalities of the urinary tract. In: Davison AM, editor. *Oxford textbook of clinical nephrology.* Oxford University press; 2005. p. 2471-94.
7. Krzemieln G, Blaim MR, Kostro I, et al. Urological anomalies in children with renal agenesis or multicystic dysplastic kidney. *J Appl Genet.* 2006;47:171-6.
8. Evans JA, Vitez M, Czeizel A. Patterns of acrorenal malformation associations. *Am J Med Genet.* 1992;44:413-9.
9. Zeier M, Ritz E. No stinkfinger and renal failure-do you see a link? *Nephrol Dial Transplant.* 1999;14:2763-5.

10. Curran AS, Curran JP. Associated acral and renal malformations: a new syndrome? *Pediatrics*. 1972;49:716-25.
11. Natarajan G, Jeyachandran D, Subramaniyan B, et al. Congenital anomalies of kidney and hand: a review. *Clin Kidney J*. 2013;6:144-9.
12. Robson WL, Leung AK, Rogers RC. Unilateral renal agenesis. *Adv Pediatr* 1995;42:575-592.

Correspondence to:
Farshid Oliaei, MD
Department of Internal Medicine, School of Medicine, Babol
University of Medical Sciences, Babol, Iran
Tel: 011-32254899
Fax: 011- 32251664
E-mail: ol_1964@yahoo.com

Received November 2017

Accepted March 2018

Archive of SID