

Iranian Journal of Mathematical Sciences and Informatics  
Vol.1, No.2 (2006),pp 65-77

## 4-PLACEMENT OF ROOTED TREES

H. YOUSEFI-AZARI AND A. GOODARZI

DEPARTMENT OF MATHEMATICS , STATISTICS AND COMPUTER SCIENCE,  
UNIVERSITY OF TEHRAN, TEHRAN, IRAN

EMAIL: HYOUSEFI@UT.AC.IR

ABSTRACT. A tree  $T$  of order  $n$  is called  $k$ -placeable if there are  $k$  edge-disjoint copies of  $T$  into  $K_n$ . In this paper we prove some results about 4-placement of rooted trees.

**Keywords:** Embedding, packing,  $k$ -placement, star-path, star-path-star.

**2000 Mathematics subject classification:**05C70, 05C05.

### 1. INTRODUCTION

We use standard graph theory notation. We consider in our study only simple undirected graphs. Suppose  $G_1, G_2, \dots, G_k$  are graphs of order  $n$ . We say that there is a packing of  $G_1, G_2, \dots, G_k$  (into the complete graph  $K_n$ ) if there is injections  $\alpha_i : V(G_i) \rightarrow V(K_n), i = 1, 2, \dots, k$  such that  $\alpha_i^*(E(G_i)) \cap \alpha_j^*(E(G_j)) = \emptyset$  for  $i \neq j$  where the map  $\alpha_i^* : E(G_i) \rightarrow E(K_n)$  is the one induced by  $\alpha_i$ .

A packing of  $k$  copies of a graph  $G$  is called a  $k$ -placement of  $G$ . Then we say  $G$  is  $k$ -placeable (into  $K_n$ ). The main references of the paper are the last chapter of Bollobás book [1] and the survey paper [10], (cf. also [4,7,8]).

Let  $P_n$  and  $S_n$  be the path and star on  $n$  vertices respectively. A tree of order  $n$  obtained from one path and one star by joining two end-vertices of theirs is called path-star and denoted by  $(PS)_n$ . A  $(PS)_n$  in which the order of path is  $r$  denoted by  $(P_rS)_n$ . Similarly a tree of order  $n$  is obtained from two stars by joining their centers by a path is called star-path-star and denoted by  $(SPS)_n$ .

---

This paper was supported by a grant No.6103007/1/01 by the research council of the University of Tehran

©2006 Academic Center for Education, Culture and Research TMU.

A rooted tree  $T$  of order  $n$  with root  $v_0 \in V(T)$ , by three branches  $T_1, T_2, T_3$  of  $T$  (the components of  $T - v_0$ ) is denoted by  $RT_n(T_1, T_2, T_3)$ . In this paper we prove any  $RT_{n-2}(T_1, T_2, T_3)$ ,  $T_i = (PS)_{n_i}$   $i = 1, 2, 3$ ,  $n_1 + n_2 + n_3 = n - 3$  is 4-placeable.

## 2. PRELIMINARIES

The following conjecture stated by Bollobás and Eldridge [2].

**Conjecture 1.** Let  $G_1, G_2, \dots, G_k$  be  $k$  graphs of order  $n$ . If  $|E(G_i)| \leq n - k$ ,  $i = 1, \dots, k$  then  $G_1, G_2, \dots, G_k$  are packable into  $K_n$ .

The cases  $k = 2, 3$  of the above-mentioned conjecture, was proved in [6] and [5], respectively.

Let us mention that some related problems have already been considered. For instance in the case of trees the hypothesis on the size may be improved. The first theorem concerning the packing of three trees was probably proved in connection with the following conjecture stated by Gyárfás in [3].

**Conjecture 2.** Let  $T_i$  denote a tree of order  $i$ . The sequence of trees  $T_2, T_3, \dots, T_n$  can be packed into  $K_n$ .

This conjecture is sometimes called tree packing conjecture (TPC). The TPC has been proved valid for quite a few cases. We list some of the results below.

**Theorem 2.1.** (Gyárfás and Lehel[3] ) Any sequence of trees  $T_2, T_3, \dots, T_n$  in which all but two are stars, can be packed into  $K_n$ .

**Theorem 2.2.** (Gyárfás and Lehel[3] ) Any sequence of trees  $T_2, T_3, \dots, T_n$  where  $T_i \in \{S_i, P_i\}$   $i = 2, 3, \dots, n$ , can be packed into  $K_n$ .

Hobbs in [4] proved that :

**Theorem 2.3.** Any three trees of order  $n_1 < n_2 < n_3 \leq n$ , respectively, can be packed into  $K_n$ .

Inspired by the above theorem, a similar result was obtained in [9] :

**Theorem 2.4.** Any three trees of order  $n - 1$  can be packed into  $K_n$ .

The authors in [11] proved the following Theorem :

**Theorem 2.5.** Suppose  $T$  is a tree of order  $n-2$  where  $T \in \{P_{n-2}, S_{n-2}, (PS)_{n-2}, (SPS)_{n-2}\}$ . Then four copies of  $T$  can be packed into  $K_n$ .

## 3. PACKING FOUR COPIES OF ROOTED TREES

In this paper we suppose  $c_i$   $i = 1, 2, 3, 4$  are four different colors. The notation  $(T)_{c_i}$  means that the edges of tree  $T$  are colored by  $c_i$ . In the Figures we used four different lines instead of colors. Suppose  $N = \{1, 2, \dots, n\}$  is the




FIGURE 1

vertex set of  $K_n$ . The star  $S_n$  with center  $v$  is denoted by  $S_n(V)$ , in particular if  $A = V(S_n) - \{v\}$  then we use the notation  $S_n(v, A)$ .

The main result of this paper is as follows :

**Theorem 3.1.** *Every  $RT_{n-2}(T_1, T_2, T_3)$ ,  $T_i = (PS)_{n_i}$ ,  $i = 1, 2, 3$ ,  $n_1 + n_2 + n_3 = n - 3$  (Figure 16) is 4-placeable.*

We prove this theorem in the end of this section.

**Lemma 3.2.** *Every  $RT_{n-2}(T_1, T_2, T_3)$  with children  $v_i$ ,  $i = 1, 2, 3$ ,  $T_i = S_{n_i+1}(v_i)$ ,  $n_1 + n_2 + n_3 = n - 6$  (Figure 1) is 4-placeable.*

**Proof.** Suppose  $A_1 = \{3, 4, 5\}$ ,  $A_2 = \{2, 3, 5\}$ ,  $A_3 = \{1, 3, 6\}$ ,  $A_4 = \{2, 5, 6\}$  be subsets of  $V(K_6)$ . Then the edge-disjoint stars  $(T)_{c_1} = S_4(2, A_1)$ ,  $(T)_{c_2} = S_4(6, A_2)$ ,  $(T)_{c_3} = S_4(4, A_3)$ ,  $(T)_{c_4} = S_4(1, A_4)$  are a 4-placement of  $S_4$  into  $K_6$  (Figure 2). Now we partition the  $n - 6$  other vertices of  $K_n$  (except the above six vertices) to three sets  $B_i$  such that  $|B_i| = n_i$ ,  $i = 1, 2, 3$ . Then color the edges of  $T_1 = S_{n_1+1}(4, B_1)$ ,  $T_2 = S_{n_2+1}(3, B_2)$ ,  $T_3 = S_{n_3+1}(5, B_3)$  with color  $c_1$ . So the  $RT_{n-2}(T_1, T_2, T_3)$  with children  $v_1 = 3$ ,  $v_2 = 4$ ,  $v_3 = 5$  and root  $v_0 = 2$  is embedded into  $K_n$ . By the same way we embed three another copies of  $RT_{n-2}(T_1, T_2, T_3)$  into  $K_n$  as follows :

1. Let  $T_1 = S_{n_1+1}(2, B_1)$ ,  $T_2 = S_{n_1+1}(5, B_2)$ ,  $T_3 = S_{n_3+1}(3, B_3)$  colored by  $c_2$ .
2. Let  $T_1 = S_{n_1+1}(5, B_1)$ ,  $T_2 = S_{n_1+1}(2, B_2)$ ,  $T_3 = S_{n_3+1}(6, B_3)$  colored by  $c_3$ .
3. Let  $T_1 = S_{n_1+1}(3, B_1)$ ,  $T_2 = S_{n_1+1}(6, B_2)$ ,  $T_3 = S_{n_3+1}(1, B_3)$  colored by  $c_4$ .

For example we show packing of two copies (colored by  $c_2, c_4$ ) of these trees into  $K_n$  in Figure 3. Clearly these four copies of  $RT_{n-2}(T_1, T_2, T_3)$  are edge-disjoint and so it is 4-placeable.




FIGURE 2




FIGURE 3

**Lemma 3.3.** *Every  $RT_{n-2}(T_1, T_2, T_3)$  with children  $v, v_i$ ,  $i = 1, 2$ ,  $T_i = S_{n_i+1}(v_i)$ ,  $i = 1, 2$ ,  $T_3 = S_{n_3+1}(v)$ ,  $v \neq v_3$ ,  $n_1 + n_2 + n_3 = n - 7$  (Figure 4) is 4-placeable.*

**Proof.** We consider the rooted tree of Figure 5. There is a packing of four copies of it into  $K_7$  (Figure 6). Now we partition the  $n - 7$  other vertices of  $K_n$  (except the above seven vertices) to three sets  $B_i$  such that  $|B_i| = n_i$ ,  $i = 1, 2, 3$ . Then we use the method of Lemma 1 as follows:




FIGURE 4




FIGURE 5

1. Let  $T_1 = S_{n_1+1}(4, B_1)$  ,  $T_2 = S_{n_2+1}(3, B_2)$  ,  $T_3 = S_{n_3+1}(5, B_3)$  colored by  $c_1$ .
2. Let  $T_1 = S_{n_1+1}(2, B_1)$  ,  $T_2 = S_{n_2+1}(5, B_2)$  ,  $T_3 = S_{n_3+1}(3, B_3)$  colored by  $c_2$ .
3. Let  $T_1 = S_{n_1+1}(5, B_1)$  ,  $T_2 = S_{n_2+1}(2, B_2)$  ,  $T_3 = S_{n_3+1}(6, B_3)$  colored by




FIGURE 6

$c_3$ .

4. Let  $T_1 = S_{n_1+1}(3, B_1)$ ,  $T_2 = S_{n_2+1}(6, B_2)$ ,  $T_3 = S_{n_3+1}(1, B_3)$  colored by  $c_4$ .

Then four copies of  $RT_{n-2}(T_1, T_2, T_3)$  are edge-disjoint and so they form a packing (into  $K_n$ ).

**Lemma 3.4.** Every  $RT_{n-2}(T_1, T_2, T_3)$  with children  $v_i$ ,  $i = 1, 2, 3$ ,  $T_i = S_{n_i+1}(v_i)$ ,  $i = 1, 2$ ,  $T_3 = (P_4S)_{n_3+1}$ ,  $n_1 + n_2 + n_3 = n - 9$  (Figure 7) is 4-placeable.

**Proof.** By Figure 9, the tree pictured in Figure 8 is 4-placeable into  $K_9$ . We partition  $n - 9$  vertices of  $K_n$  (except the above 9 vertices) into three sets  $B_i$  such that  $|B_i| = n_i$   $i = 1, 2, 3$ . Set

1.  $T_1 = S_{n_1+1}(5, B_1)$ ,  $T_2 = S_{n_2+1}(1, B_2)$ ,  $T_3 = S_{n_3+1}(9, B_3)$  colored by  $c_1$ .
2.  $T_1 = S_{n_1+1}(9, B_1)$ ,  $T_2 = S_{n_2+1}(4, B_2)$ ,  $T_3 = S_{n_3+1}(5, B_3)$  colored by  $c_2$ .
3.  $T_1 = S_{n_1+1}(4, B_1)$ ,  $T_2 = S_{n_2+1}(2, B_2)$ ,  $T_3 = S_{n_3+1}(1, B_3)$  colored by  $c_3$ .
4.  $T_1 = S_{n_1+1}(2, B_1)$ ,  $T_2 = S_{n_2+1}(5, B_2)$ ,  $T_3 = S_{n_3+1}(3, B_3)$  colored by  $c_4$ .

Clearly these four copies of  $RT_{n-2}(T_1, T_2, T_3)$  are edge-disjoint, so they form a 4-placement.

**Lemma 3.5.** Every  $RT_{n-2}(T_1, T_2, T_3)$  with children  $v_i$ ,  $i = 1, 2, 3$ ,  $T_i = S_{n_i+1}(v_i)$ ,  $i = 1, 2$ ,  $T_3 = (P_mS)_{n_3+1}$ ,  $m > 8$ ,  $n_1 + n_2 + n_3 = n - m - 6$  (Figure 10) is 4-placeable.

**Proof.** It is well known that every  $K_n$  contains  $\lfloor \frac{n-1}{2} \rfloor$  edge-disjoint Hamiltonian paths. So there is a 4-placement of  $P_3$  into  $K_m$  for  $m > 8$ . There are




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10

also 4-placement of  $S_2$  into  $K_6$ . Suppose  $K_6, K_m$  be above complete graphs such that  $V(K_6) = \{1, 2, \dots, 6\}$ ,  $V(K_m) = \{7, 8, \dots, m + 6\}$ . If we join the center of every above  $S_2$  to end-vertex of one of the four paths then we get a 4-placement of tree in Figure 11 into  $K_{m+6}$  (Figure 12). Now we partition  $n - m - 6$  other vertices of  $k_n$  (except  $m + 6$  above vertices) to three sets  $B_i$   $i = 1, 2, 3$  such that  $|B_i| = n_i$ ,  $n_1 + n_2 + n_3 = n - m - 6$ . Finally let:




FIGURE 11




FIGURE 12

1.  $(T)_{c_1} = S_{n_1+1}(5, B_1)$   $(T)_{c_1} = S_{n_1+1}(4, B_2)$   $(T)_{c_1} = S_{n_3+1}(6, B_3)$
2.  $(T)_{c_2} = S_{n_1+1}(3, B_1)$   $(T)_{c_2} = S_{n_1+1}(6, B_2)$   $(T)_{c_2} = S_{n_3+1}(4, B_3)$
3.  $(T)_{c_3} = S_{n_1+1}(6, B_1)$   $(T)_{c_3} = S_{n_1+1}(3, B_2)$   $(T)_{c_3} = S_{n_3+1}(1, B_3)$




FIGURE 13




FIGURE 14

$$4. (T)_{c_4} = S_{n_1+1}(4, B_1) \quad (T)_{c_4} = S_{n_1+1}(2, B_2) \quad (T)_{c_4} = S_{n_3+1}(3, B_3)$$

So we have embedded four edge-disjoint copies of tree in Figure 10 into  $K_n$ .

**Lemma 3.6.** *Every  $RT_{n-2}(T_1, T_2, T_3)$  with children  $v_i$ ,  $i = 1, 2, 3$ ,  $T_1 = S_{n_1+1}(v_1)$ ,  $T_i = (P_{m_i}S)_{n_i+1}$ ,  $m_i > 8$ ,  $i = 1, 2$ ,  $n_1+n_2+n_3 = n - (m_1+m_2) - 6$  (Figure 13) is 4-placeable.*

**Proof.** The Figure 15 shows there are 4-placement of tree in Figure 14 into  $K_{m_1+m_2+6}$  with the following vertices and colors:  
 Copy 1 :  $u = 3$   $v_1 = 6$   $v_2 = 4$   $v_3 = 5$  colored by  $c_1$ .




FIGURE 15

- Copy 2 :  $u = 1 \ v_1 = 4 \ v_2 = 6 \ v_3 = 3$  colored by  $c_2$ .
- Copy 3 :  $u = 2 \ v_1 = 3 \ v_2 = 1 \ v_3 = 6$  colored by  $c_3$ .
- Copy 4 :  $u = 5 \ v_1 = 1 \ v_2 = 2 \ v_3 = 4$  colored by  $c_4$ .

Now we partition  $n - (m_1 + m_2) - 6$  other vertices of  $k_n$  (except  $m_1 + m_2 + 6$  above vertices) to three sets  $B_i \ i = 1, 2, 3$  such that  $|B_i| = n_i$  ,  $n_1 + n_2 + n_3 = n - (m_1 + m_2) - 6$ . Then we consider four edge-disjoint trees.

1.  $S_{n_1+1}(5, B_1)$  ,  $S_{n_2+1}(4, B_2)$  ,  $S_{n_3+1}(6, B_3)$  colored by  $c_1$ .
2.  $S_{n_1+1}(3, B_1)$  ,  $S_{n_2+1}(6, B_2)$  ,  $S_{n_3+1}(4, B_3)$  colored by  $c_2$ .




FIGURE 16




FIGURE 17

3.  $S_{n_1+1}(6, B_1)$  ,  $S_{n_2+1}(1, B_2)$  ,  $S_{n_3+1}(3, B_3)$  colored by  $c_3$ .
4.  $S_{n_1+1}(4, B_1)$  ,  $S_{n_2+1}(2, B_2)$  ,  $S_{n_3+1}(1, B_3)$  colored by  $c_4$ .

So four copies of tree in Figure 13 can be packed into  $K_n$ .

**Proof of Theorem 6.** As the proof of Lemma 5 there are a 4-placement of tree in Figure 17 into  $K_{m_1+m_2+m_3+6}$  with the following vertices and colors :

- Copy 1 :  $u = 2$   $v_1 = 4$   $v_2 = 3$   $v_3 = 5$  colored by  $c_1$ .
- Copy 2 :  $u = 6$   $v_1 = 2$   $v_2 = 5$   $v_3 = 3$  colored by  $c_2$ .
- Copy 3 :  $u = 1$   $v_1 = 5$   $v_2 = 2$   $v_3 = 6$  colored by  $c_3$ .
- Copy 4 :  $u = 4$   $v_1 = 3$   $v_2 = 6$   $v_3 = 1$  colored by  $c_4$ . Now we partition  $n - (m_1 + m_2 + m_3) - 6$  other vertices of  $K_n$  (except  $m_1 + m_2 + m_3 + 6$  above

vertices) to three sets  $B_i$   $i = 1, 2, 3$  such that  $|B_i| = n_i$  ,  $n_1 + n_2 + n_3 = n - (m_1 - m_2 - m_3) - 6$ . Then let :

1.  $S_{n_1+1}(4, B_1)$  ,  $S_{n_2+1}(3, B_2)$  ,  $S_{n_3+1}(5, B_3)$  colored by  $c_1$ .
2.  $S_{n_1+1}(2, B_1)$  ,  $S_{n_2+1}(5, B_2)$  ,  $S_{n_3+1}(3, B_3)$  colored by  $c_2$ .
3.  $S_{n_1+1}(5, B_1)$  ,  $S_{n_2+1}(2, B_2)$  ,  $S_{n_3+1}(6, B_3)$  colored by  $c_3$ .
4.  $S_{n_1+1}(3, B_1)$  ,  $S_{n_2+1}(6, B_2)$  ,  $S_{n_3+1}(1, B_3)$  colored by  $c_4$ .

Therefore  $RT_{n-2}(T_1, T_2, T_3)$  of Figure 15 is 4-placeable and we have proved the main Theorem.

#### REFERENCES

- [1] B. Bollobás, *Extremal Graph Theory*, Academic Press, London, 1978.
- [2] B. Bollobas and S. E. Eldridge, Packing of graphs and applications to computational complexity, *J. Combin. Theory Ser. B* 25 (1978) 105-124.
- [3] A. Gyárfás and J. Lehel, Packing trees of different order into  $K_n$ , *Colloq. Math. Soc. J. Bolyai*, 18 (1976), 463-469.
- [4] A. M. Hobbas, B. A. Bourgeois and J. Kasiraj, Packing trees in complete graphs, *Discrete Math.*, 67 (1987), 27-42.
- [5] H. Kheddouci, S. Marshall, J. S. Sacle and M. Wozniak, On the packing of three graphs, *Discrete Math.*, 236 (2001), 197-225.
- [6] N. Sauer and J. Spencer. Edge disjoint placement of graphs, *J. Combin. Theory Ser. B*, 25 (1978), 295-302.
- [7] H. Wang and N. Sauer, Packing three copies of a tree into a complete graph, *European J. Combin.*, 14 (1993), 137-142.
- [8] M. Wozniak, Embedding graphs of small size, *Discrete Appl. Math.*, 51 (1994), 233-241.
- [9] M. Wozniak, Packing three trees, *Discrete Math.*, 150 (1996), 393-402.
- [10] H. P. Yap, Packing of graphs-a survey, *Discrete Math.*, 72 (1988), 395-404.
- [11] H. Yousefi-Azari and A. Goodarzi, *On the packing of four trees*, Global J. Math. Math. Sci. (to appear).