A Note on Tensor Product of Graphs

Sirous Moradi

Department of Mathematics, Faculty of Science, Arak University, Arak, 38156-8-8349, Iran

ABSTRACT. Let G and H be graphs. The tensor product $G \otimes H$ of G and H has vertex set $V(G \otimes H) = V(G) \times V(H)$ and edge set $E(G \otimes H) = \{(a,b)(c,d) | ac \in E(G) \text{ and } bd \in E(H)\}$. In this paper, some results on this product are obtained by which it is possible to compute the Wiener and Hyper Wiener indices of $K_n \otimes G$.

Keywords: Tensor product, connected graph, bipartite graph, Wiener index, hyper Wiener index.

2000 Mathematics subject classification: 05C35, 05C12, 05A20, 05C05.

1. INTRODUCTION

A graph G consists of a set of vertices V(G) and a set of edges E(G). For every vertices $a, b \in V(G)$, the edge connecting a and b is denoted by ab. The distance between two vertices in a connected graph G is the number of edges in a shortest path between them. For vertices a and b of G, their distance is shown by $d_G(a, b)$. This concept has been known for a very long time and has received considerable attention as a subject of research in metric graph theory. Graph operations play an important role in the study of graph decompositions into isomorphic subgraphs. For more details about graph operations see [1, 4, 6, 7, 8, 9, 10, 11, 15]. For any two simple graphs G and H, the tensor product of G and H has vertex set $V(G \otimes H) = V(G) \times V(H)$ and edge set $E(G \otimes H) = \{(a,b)(c,d) | ac \in E(G) \text{ and } bd \in E(H)\}$. We refer the reader to

E-mail address: s-moradi@araku.ac.ir (S. Moradi)

Received 12 May 2011; Accepted 17 December 2011 ©2012 Academic Center for Education, Culture and Research TMU

[2] for the proof of this fact that $|E(G \otimes H)| = 2|E(G)||E(H)|$. The distance between two vertices under tensor product of graphs is studied in [13]. Here we obtain useful equality for distance of vertices in tensor product of graphs by a new method and using simple concepts. Then we use these results for obtaining simple proofs for some classical theorems [4]. One of the concepts related to distance in graphs is the Wiener index [14]. It is not only an early index which correlates well with many physico-chemical properties of organic compounds but also a subject that has been studied by many mathematicians and chemists. The Wiener index is the sum of distances between all vertex pairs in a connected graph, $W(G) = \sum_{\{a,b\} \subseteq V(G)} d_G(a,b)$. The hyper-Wiener index of acyclic graphs was introduced by Milan Randic in 1993. Then Klein et al. [12], generalized Randics definition for all connected graphs, as a generalization of the Wiener index. It is defined as $WW(G) = \frac{1}{2}W(G) + \frac{1}{2}\sum_{\{a,b\} \subseteq V(G)} d^2(a,b)$. In last section of this paper, formula for the Wiener and hyper-Wiener indices of the tensor product of complete graph K_n by a graph G are obtained. For more details about Wiener and hyper-Wiener index see for example [3, 5].

2. MAIN RESULT AND DISCUSSION

In this section some new concepts are presented by which it is possible to find new proof for some classical results.

Definition 1. Let G be a graph and $x, y \in V(G)$. Define $d'_G(x, y)$ as follows: (i) If $d_G(x, y)$ is odd then $d'_G(x, y)$ is defined as the length of a shortest even walk joining x and y in G, and if there is no shortest even walk then $d'_G(x, y) = +\infty$

(ii) If $d_G(x, y)$ is even then $d'_G(x, y)$ is defined as the length of a shortest odd walk joining x and y in G, and if there is no shortest odd walk then $d'_G(x, y) = +\infty$.

(iii) If
$$d_G(x, y) = +\infty$$
, then $d'_G(x, y) = +\infty$.

Example 1. Let K_n be a complete graph with $n \ge 3$. For each $a, b \in V(K_n)$, if $a \ne b$ then $d_{K_n}(a, b) = 1$ and $d'_{K_n}(a, b) = 2$ and if a = b then $d_{K_n}(a, b) = 0$ and $d'_{K_n}(a, b) = 3$. Also let C_{2n+1} be a cycle of order 2n + 1, then for each two vertices $a, b \in V(C_{2n+1})$, it is easy to see that, $d'_{C_{2n+1}}(a, b) = 2n + 1 - d_{C_{2n+1}}(a, b)$. In every even cycle C_{2n} , $d'_{C_{2n}}(a, b) = +\infty$, where $a, b \in V(C_{2n})$.

Proposition 1. Let G be a graph. Then G is not bipartite graph if and only if there exists $x, y \in V(G)$ such that $d'_G(x, y) < +\infty$.

Proof. Let G is not bipartite, then there is an odd cycle C in G. Suppose x and y are two vertices on C. If $d_G(x, y)$ is even (odd) then there exists an odd (even) path between x and y in C, so $d'_G(x, y) < +\infty$. Conversely, let $x, y \in V(G)$ and $d'_G(x, y) < +\infty$. Therefore is odd and so there exists an odd

closed walk contains x. Thus there is an odd cycle in G and this proves G is non-bipartite.

Note. If G is a connected and non-bipartite graph, then $d'_G(x, y) < +\infty$, for all $x, y \in V(G)$.

Lemma 1. Let G and H be graphs and $(a,b)(c,d) \in V(G \otimes H)$. Then

(2.1)
$$Max\{d_G(a,c), d_H(b,d)\} \le d_{G\otimes H}((a,b), (c,d)).$$

Proof. If $d_{G\otimes H}((a,b),(c,d)) = +\infty$, then inequality (2.1) holds. Suppose that $d_{G\otimes H}((a,b),(c,d)) = n$. So the following path between (a,b) and (c,d) in $G\otimes H$ exists:

$$P: (a,b) = (x_0, y_0), (x_1, y_1), ..., (x_n, y_n) = (c, d).$$

Obviously $a = x_0, x_1, ..., x_n = c$ is a walk with length n between a and c in G and $b = y_1, y_2, ..., y_n = d$ is a walk with length n between b and d in H. Hence $d_G(a, c), d_H(b, d) \leq n$ and then,

$$Max\{d_G(a,c), d_H(b,d)\} \le n = d_{G\otimes H}((a,b), (c,d)).$$

Definition 2. Let G and H be two graphs and $(a, b), (c, d) \in V(G \otimes H)$. The relation \sim on the vertices of $G \otimes H$ is defined as follows:

 $(a,b)\sim (c,d)$ if and only if $d_G(a,c), d_H(b,d)<+\infty$ and $d_G(a,c)+d_H(b,d)$ is even.

Proposition 2. Let G and H be graphs and $(a,b), (c,d) \in V(G \otimes H)$. If $(a,b) \sim (c,d)$, then (a,b) and (c,d) are in the same component of $G \otimes H$. Moreover

$$Max\{d_{G}(a, c), d_{H}(b, d)\} = d_{G \otimes H}((a, b), (c, d)).$$

Proof. Suppose $d_G(a,c) = m$, $d_H(b,d) = n$ and $m \leq n$. The following paths between a and c in G and b and d in H exist:

$$P_1: a = x_0, x_1, ..., x_m = c \text{ and } P_2: b = y_1, y_2, ..., y_n = d.$$

Since $(a, b) \sim (c, d)$, then m + n and n - m are even. Therefore there exists the following path between (a, b) and (c, d) in $G \otimes H$, with length n:

$$P: (a,b) = (x_0, y_0), (x_1, y_1), \dots, (x_m, y_m) = (c, y_m), (x_{m-1}, y_{m+1}), (x_m, y_{m+2}), \dots, (x_m, y_n) = (c, d).$$

So $d_{G\otimes H}((a,b),(c,d)) \leq n = d_H(b,d) = Max\{d_G(a,c),d_H(b,d)\}$. Similarly, if m > n, then $d_{G\otimes H}((a,b),(c,d)) \leq n = d_G(a,c) = Max\{d_G(a,c),d_H(b,d)\}$. By using Lemma $1,Max\{d_G(a,c),d_H(b,d)\} \leq d_{G\otimes H}((a,b),(c,d))$, and this completes the proof. **Proposition 3.** Let G and H be graphs and $(a,b), (c,d) \in V(G \otimes H)$. If $(a,b) \nsim (c,d)$ then,

$$d_{G\otimes H}((a,b),(c,d)) = Min\Big\{Max\{d_G(a,c),d'_H(b,d)\}, Max\{d'_G(a,c),d_H(b,d)\}\Big\}.$$

Proof. At first we show that the following inequality is holds:

(2.2)
$$d_{G\otimes H}((a,b),(c,d)) \leq Min \left\{ Max\{d_G(a,c),d'_H(b,d)\}, \\ Max\{d'_G(a,c),d_H(b,d)\} \right\}.$$

If $d_G(a,c) = +\infty$ $(d_H(b,d) = +\infty)$, then $d'_G(a,c) = +\infty$ $(d'_G(a,c) = +\infty)$, then the above inequality is satisfied.

We now assume that $d_G(a, c), d_H(b, d) < +\infty$. It is enough to show the following inequalities are hold:

(2.3)
$$d_{G\otimes H}((a,b),(c,d)) \le Max\{d_G(a,c),d'_H(b,d)\},\$$

(2.4)
$$d_{G\otimes H}((a,b),(c,d)) \le Max\{d'_G(a,c),d_H(b,d)\}$$

If $d'_H(b,d) = +\infty$ then inequality (2.3) holds. Suppose that $d'_H(b,d) = n$ and $d_G(a,c) = m$ and $m \le n$. There exist the shortest path $P: a = x_0, x_1, ..., x_m = c$ in G and the walk $W: b = y_0, y_1, ..., y_n = d$ in H. Since $(a,b) \nsim (c,d)$ then n + m and n - m are even. Therefore there exist the path P' between (a,b) and (c,d) in as follows:

$$\begin{array}{lll} P':(a,b) &=& (x_0,y_0), (x_1,y_1), ..., (x_m,y_m) = (c,y_m), \\ && (x_{m-1},y_{m+1}), (x_m,y_{m+2}), ..., (x_m,y_n) = (c,d). \end{array}$$

Hence, $d_{G\otimes H} \leq n = d'_H(b,d) = Max\{d_G(a,c), d'_H(b,d)\}$. Similarly, if n < m then the inequality (2.3) is satisfied. By a similar argument, one can prove the inequality (2.4). By above argument we have shown inequality (2.2) is satisfied. We now prove the following:

(2.5)
$$d_{G\otimes H}((a,b),(c,d)) \ge Min \left\{ Max\{d_G(a,c),d'_H(b,d)\}, Max\{d'_G(a,c),d_H(b,d)\} \right\}$$

If (a, b) and (c, d) are not in the same component of tensor product of G and H then, $d_{G\otimes H}((a, b), (c, d)) = +\infty$ and (2.5) is satisfied. Suppose (a, b) and (c, d) are in the same component. Then a and c (b and d) are in the same component of G (H). If $d_{G\otimes H}((a, b), (c, d)) = n$ then the path P'' exists, P'': $(a, b) = (x_0, y_0), (x_1, y_1), ..., (x_n, y_n) = (c, d)$. Since $(a, b) \nsim (c, d)$ then $d_G(a, c) + d_H(b, d)$ is odd. We can suppose that $d_G(a, c)$ is odd and $d_H(b, d)$ is even. Our main proof consider the following cases:

Case 1: If n is odd, then the walk W_1 : $b = y_0, y_1, ..., y_n = d$ is an odd walk with length n between b and d in H. So by definition of $d'_H(b, d)$,

it is clear that $d'_H(b,d) \leq n$. It is easy to see $d_G(a,c) \leq n$. Therefore $Max\{d_G(a,c), d'_H(b,d)\} \leq n$.

Case 2: If n is even, then by a similar method we can see

$$Max\{d'_G(a,c), d_H(b,d)\} \le n$$

Hence

 $Min\{Max\{d_G(a,c), d'_H(b,d)\}, Max\{d'_G(a,c), d_H(b,d)\}\} \le d_{G\otimes H}((a,b), (c,d)).$ and the proof is complete. \Box

Corollary 1. Let G and H be graphs and $(a,b), (c,d) \in V(G \otimes H)$. Then

$$d_{G\otimes H}((a,b),(c,d)) = \begin{cases} d_1((a,b),(c,d)) & (a,b) \sim (c,d) \\ d_1((a,b),(c,d)) & (a,b) \nsim (c,d) \end{cases}$$

where,

$$d_1((a,b), (c,d)) = Max\{d_G(a,c), d_H(b,d)\}$$

and

$$d_2((a,b),(c,d)) = Min\{Max\{d_G(a,c),d'_H(b,d)\}, Max\{d'_G(a,c),d_H(b,d)\}\}.$$

3. Application

In this section at first we present new simple proofs for some classical theorems related to the tensor product of graphs and then as an application of previous results, we compute the Wiener and Hyper Wiener indices of the tensor product of K_n and G, where K_n is the complete graph of order n and G is a connected graph.

Proposition 4. ([4]) Let G and H be connected graphs with at least two vertices. Then $G \otimes H$ is connected if and only if G or H is not bipartite.

If G or H are not bipartite then by Proposition 4, $G \otimes H$ is connected and has just one component. Now suppose G and H are bipartite. If |V(G)| = |V(H)| = 2, then $G \otimes H$ also is connected and has just one component. Let |V(G)| > 2 or |V(H)| > 2. Since G is bipartite, there exists the path P : a_1, a_2, a_3 in G. Obviously $d_G(a_1, a_3) = 2$. Let b_1 and b_2 are adjacent vertices in H. Since $(a_1, b_1) \not\approx (a_2, b_1)$, then by using Proposition 3 or Corollary 1, $d_{G \otimes H}((a_1, b_1), (a_2, b_1)) = +\infty$. Then (a_1, b_1) and (a_2, b_1) are in different components. For each $(a, b) \in V(G \otimes H)$, two cases are considered:

Case 1: If $d_G(a, a_1) + d_H(b, b_1)$ is even, then $(a, b) \sim (a_1, b_1)$ and by Corollary 1, (a, b) and (a_1, b_1) are in the same component.

Case 2: If $d_G(a, a_2) + d_H(b, b_1)$ is even, then $(a, b) \sim (a_2, b_1)$ and by Corollary 1, (a, b) and (a_2, b_1) are in the same component. Notice that if $d_G(a, a_1) + d_H(b, b_1)$ and $d_G(a, a_2) + d_H(b, b_1)$ are odd, then both of $d_G(a, a_1)$ and $d_G(a, a_2)$ are even or odd. So $d_G(a, a_1) + d_G(a, a_2)$ is even, and hence there exists an even walk between a_1 and a_2 . So there exists an odd closed walk in G containing

which is a contradiction.

78

Therefore we can conclude the following well known result.

Proposition 5. ([4]) Suppose G and H are connected graphs with at least two vertices, then $G \otimes H$ has at most two components.

In the following propositions the Wiener and Hyper Wiener indices of the tensor product of K_n and a connected graph G are computed.

Proposition 6. Let G be connected graph of order m. Then the Wiener index of the tensor product of K_n and G is given by:

$$W(K_n \otimes G) = n^2 W(G) + 2\binom{n}{2} |V(G)| + 2n|E(G)| - n|T_G|,$$

where $n \geq 3$ and $T_G = \{ab \in E(G) | ab \text{ is contained in a triangle} \}$.

Proof. By definition of the Wiener index,

$$W(K_n \otimes G) = \sum_{\{(a,b),(c,d)\} \subseteq V(K_n \otimes G)} d_{K_n \otimes G} ((a,b), (c,d))$$

We consider a partition of $V(K_n \otimes G)$ into the following parts:

$$A_{1} = \left\{ \{(a,b), (c,d)\} | a \neq c, b \neq d, (a,b) \sim (c,d) \right\},\$$

$$A_{2} = \left\{ \{(a,b), (c,d)\} | a \neq c, b \neq d, (a,b) \nsim (c,d) \right\},\$$

$$A_{3} = \left\{ \{(a,b), (c,d)\} | a \neq c, b = d \right\},\$$

$$A_{4} = \left\{ \{(a,b), (c,d)\} | a = c, b \neq d, (a,b) \sim (c,d) \right\},\$$

$$A_{5} = \left\{ \{(a,b), (c,d)\} | a = c, b \neq d, (a,b) \nsim (c,d) \right\}.$$

Therefore,

 $W(K_n \otimes G) =$

$$\sum_{\{(a,b),(c,d)\}\in A_1} d_{K_n\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_2} d_{K_n\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_3} d_{K_n\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_4} d_{K_n\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_5} d_{K_n\otimes G}((a,b),(c,d)).$$

We evaluate each summation separately. It is obvious that if $\{a, c\} \subseteq V(K_n)$, then $d_{K_n}(a, c) = 1$ and $d'_{K_n}(a, c) = 2$. Apply Proposition 3, if $(a, b) \sim (c, d)$ and $a \neq c, b \neq d$ then,

 $Min\{Max\{d_{K_n}(a,c), d'_G(b,d)\}, Max\{d'_{K_n}(a,c), d_G(b,d)\}\} = d_G(b,d)$

and

$$Max\{d_{K_n}(a,c), d_G(b,d)\} = d_G(b,d)$$

Therefore,

$$\sum_{\{(a,b),(c,d)\}\in A_1\cup A_2} d_{K_n\otimes G}((a,b),(c,d)) = \sum_{\{(a,b),(c,d)\}\in A_1} d_{K_n\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_2} d_{K_n\otimes G}((a,b),(c,d))$$
$$= 2\sum_{\{a,c\}\subseteq V(K_n)} \sum_{\{b,d\}\}\subseteq V(G)} d_G(b,d)$$
$$= 2\binom{n}{2} W(G).$$

By considering the set A_3 , we have:

$$\sum_{\{(a,b),(c,d)\}\in A_3} d_{K_n\otimes G}((a,b),(c,d)) = \sum_{\substack{\{a,c\}\subseteq V(K_n)\\b\in V(G)}} 2 = 2|V(G)|\binom{n}{2}.$$

For computing our summation over the set A_4 , we notice that,

$$A_4 = \left\{ \{(a,b)(a,d)\} | a \in V(K_n), \{b,d\} \subseteq V(G) \text{ and } 2 | d_G(b,d) \right\}$$

Hence,

$$\sum_{\{(a,b),(c,d)\}\in A_4} d_{K_n\otimes G}((a,b),(c,d)) = \sum_{\substack{\{b,d\}\subseteq V(G)\\a\in V(K_n)\\2|d_G(b,d)}} d_{K_n\otimes G}((a,b),(c,d))$$

$$= \sum_{\substack{\{b,d\} \subseteq V(G) \\ a \in V(K_n) \\ 2|d_G(b,d)}} d_G(b,d) = n \sum_{\substack{\{b,d\} \subseteq V(G) \\ 2|d_G(b,d)}} d_G(b,d).$$

Compute our summation over A_5 , we have:

$$\sum_{\{(a,b),(c,d)\}\in A_5} d_{K_n\otimes G}\big((a,b),(c,d)\big) = \sum_{\substack{\{b,d\}\subseteq V(G)\\a\in V(K_n)\\2\notin G_{(b,d)}}} d_{K_n\otimes G}\big((a,b),(c,d)\big).$$

If $d_G(b, d)$ is odd then by Proposition 3,

$$d_{K_n \otimes G}((a, b), (c, d)) = Min \Big\{ Max\{d_{K_n}(a, a), d'_G(b, d)\}, \\ Max\{d'_{K_n}(a, a), d_G(b, d)\} \Big\} \\ = Min \Big\{ d'_G(b, d), Max\{3, d_G(b, d)\} \Big\}.$$

By a case by case calculations, one can see that

$$Min\left\{d'_{G}(b,d), Max\{3, d_{G}(b,d)\}\right\} = \left\{\begin{array}{rr} d_{G}(b,d) & d_{G}(b,d) \ge 3\\ 2 & d_{G}(b,d) = 1 \text{ and } d'_{G}(b,d) = 2\\ 3 & d_{G}(b,d) = 1 \text{ and } d'_{G}(b,d) \ge 4\end{array}\right.$$

Define:

$$\begin{aligned} A_5' &= \Big\{ \{(a,b), (c,d)\} | a \in V(K_n), d_G(b,d) \ge 3 \text{ and } d_G(b,d) \text{ is odd} \Big\}, \\ A_5'' &= \Big\{ \{(a,b), (c,d)\} | a \in V(K_n), d_G(b,d) = 1 \text{ and } d_G'(b,d) = 2 \Big\}, \\ A_5''' &= \Big\{ \{(a,b), (c,d)\} | a \in V(K_n), d_G(b,d) = 1 \text{ and } d_G'(b,d) \ge \Big\}. \end{aligned}$$

Such that $A_5 = A'_5 \cup A''_5 \cup A''_5$. Hence,

$$\sum_{\{(a,b),(c,d)\}\in A_{5}} d_{K_{n}\otimes G}((a,b),(c,d)) = \sum_{\{(a,b),(c,d)\}\in A_{5}'} d_{K_{n}\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_{5}''} d_{K_{n}\otimes G}((a,b),(c,d)) + \sum_{\{(a,b),(c,d)\}\in A_{5}''} d_{K_{n}\otimes G}((a,b),(c,d)) = \sum_{\{(a,b),(c,d)\}\in A_{5}''} d_{G}(b,d) + \sum_{\{(a,b),(c,d)\}\in A_{5}''} 2 + \sum_{\{(a,b),(c,d)\}\in A_{5}''} 3 = n\Big(\sum_{\substack{\{(a,b),(c,d)\}\in A_{5}''\\ 2 \nmid d_{G}(b,d)}} d_{G}(b,d) - |E(G)|\Big) + 2n|T_{G}| + 3n(|E(G)| - |T_{G}|).$$

By the above calculations, one can see that,

$$W(K_n \otimes G) = n^2 W(G) + 2\binom{n}{2} |V(G)| + 2n|E(G)| - n|T_G|.$$

Corollary 2. The following statements are hold:

 $a) \ Let \ G \ be \ a \ triangle-free \ graph \ then,$

$$W(K_n \otimes G) = n^2 W(G) + 2\binom{n}{2} |V(G)| + 2n|E(G)|.$$

b) The Wiener index of two complete graphs of orders m and n is computed as follows:

$$W(K_n \otimes K_m) = n(n+1)\binom{m}{2} + n(n-1)m.$$

Proof. a) It is immediate by Proposition 6.

b) In complete graph K_n , we can see that $T_{K_m} = {\cal E}(K_m)$ and so by Proposition 6,

$$W(K_n \otimes K_m) = n^2 W(K_m) + 2\binom{n}{2}m + 2n\binom{m}{2} - n\binom{m}{2}$$
$$= n(n+1)\binom{m}{2} + n(n-1)m.$$

Proposition 7. Let G be connected graph of order m. Then the hyper Wiener index of the tensor product of K_n and G is given by:

$$WW(K_n \otimes G) = \binom{n+1}{2} WW(G) + 3\binom{n}{2} |V(G)| + 5n|E(G)| - 3n|T_G|.$$

Proof. The proof is similar to the proof of Proposition 6.

Corollary 3. The hyper Wiener index of two complete graphs of orders m and n is computed as follows:

$$WW(K_n \otimes K_m) = \binom{n+1}{2}m(m-1) + 3\binom{n}{2}m + 5n\frac{m(m-1)}{2} - 3nm(m-1).$$

Acknowledgments

The authors sincerely thank the referees for their careful reading of this paper and valuable suggestions and comments.

References

- 1. J. Bosak, Decompositions of Graphs, Kluwer Academic Publishers, 1990.
- 2. A. Bottreou and Y. Metivier, Some remarks on the Kronecker product of graphs, Inform. Process. Lett., 63 (1998), 55–61.
- G. H. Fath-Tabar and A. R. Ashrafi, The Hyper-Wiener Polynomial of Graphs, Iranian Journal of Mathematical Sciences and Informatics, 6(2) (2011), 67–74.
- W. Imrich and S. Klavsar, Product Graphs, Structure and Recognition, Discrete Appl. Math., John Wiley and Sons, New York, USA, 2000.
- A. Iranmanesh and Y. Alizadeh, Computing Wiener Index of HAC₅C₇[p,q] Nanotubes by Gap Program, Iranian Journal of Mathematical Sciences and Informatics, 3(1) (2008), 1–12.
- M. H. Khalifeh, H. Yousefi-Azari, A. R. Ashrafi and I. Gutman, The edge Szeged index of product graphs, *Croat. Chem. Acta*, 81(2) (2008), 277–281.
- M. H. Khalifeh, H. Yousefi-Azari and Ashrafi, The Hyper-Wiener index of graph operations, Comput. Math. Appl., 56 (2008), 1402-1407.
- M. H. Khalifeh, H. Yousefi-Azari, A. R. Ashrafi and S. Wagner, Some new results on distance-based graph invariants, *Europ. J. Combin.*, 30(5) (2009), 1149–1163.
- M. H. Khalifeh, H. Yousefi-Azari and A. R. Ashrafi, Vertex and edge PI indices of Cartesian product graph, *Discrete Appl. Math.*, **156** (2008), 1780–1789.
- M. H. Khalifeh, H. Yousefi-Azari and A. R. Ashrafi, The first and second Zagreb indices of graph operations, *Discrete Appl. Math.*, 157(4) (2009), 804–811.
- M. H. Khalifeh, H. Yousefi-Azari and A. R. Ashrafi, A matrix method for computing Szeged and vertex PI indices of join and composition of graphs, *Linear Alg. Appl.*, 429(11,12) (2008), 2702–2709.
- D. J. Klein and I. Lukovits, On the definition of the hyper-Wiener index for cyclecontaining structures, J. Chem. Inf. Comput. Sci., 35 (1995), 50–52.
- D. Stevanovic, Distance regularity of compositions of graphs, Appl. Math. Lett., 17 (2004), 337–343.
- H. Wiener, Structural determination of the paraffin boiling points, J. Am. Chem. Soc., 69 (1974), 17–20.
- H. Yousefi-Azari, B. Manoochehrian and A. R. Ashrafi, The PI index of product graphs, Appl. Math. Lett., 21(6) (2008), 624–627.

81