Iranian Journal of Mathematical Sciences and Informatics Vol. 8, No. 2 (2013), pp 39-47

Generalization of α -Centroidal Mean and its Dual

K. M. Nagaraja^a, P. Siva Kota Reddy^{b,*} and Sudhir Kumar Sahu^c ^aDepartment of Mathematics, J. S. S. Academy of Technical Education, Bangalore-560 060, India ^bDepartment of Mathematics, Siddaganga Institute of Technology, Tumkur-572103, India ^cPost Graduate Department of Statistics, Sambalpur University, Sambalpur-768 019, India

E-mail: reddy_math@yahoo.com E-mail: pskreddy@sit.ac.in

ABSTRACT. In this paper, the generalized α -centroidal mean and its dual form in 2 variables are introduced. Also, we will study some properties and prove their monotonicity. Further, shown that various means are particular cases of generalized α -centroidal mean.

Keywords: Monotonicity, Inequality, Power Oscillatory mean, Dual.

2000 Mathematics subject classification: Primary 26D15, 26D10.

1. INTRODUCTION

If a, b > 0 are two real numbers, then

$$A(a,b) = F_1(a,b) = \frac{a+b}{2}$$
 (1.1)

$$G(a,b) = F_2(a,b) = \sqrt{ab}$$
(1.2)

$$H(a,b) = F_2(a,b) = \frac{2ab}{a+b}.$$
 (1.3)

$$L(a,b) = \begin{cases} \frac{a-b}{\ln a - \ln b} & a \neq b\\ a & a = b \end{cases}$$
(1.4)

*Corresponding author

Received 16 October 2012; Accepted 28 August 2013 ©2013 Academic Center for Education, Culture and Research TMU

$$I(a,b) = \begin{cases} e^{\left(\frac{a\ln a - b\ln b}{a - b} - 1\right)} & a \neq b \\ a & a = b \end{cases}$$
(1.5)

$$M_{r}(a,b) = \begin{cases} \left(\frac{a^{r}+b^{r}}{2}\right)^{\frac{1}{r}} & r \neq 0\\ \sqrt{ab} & r = 0 \end{cases}$$
(1.6)

$$H_p(a,b) = \left\{ \frac{a^p + (\sqrt{ab})^p + b^p}{3} \right\}^{1/p}$$
(1.7)

are respectively called arithmetic mean, geometric mean, harmonic mean, logarithmic mean, identric mean, power mean and power type heron mean thoroughly studied by various researchers having their own importance in the literature. In [12], the definition of contra-harmonic mean is given by;

$$C(a,b) = \frac{a^2 + b^2}{a+b}$$
(1.8)

and its generalization form is gives as;

$$C_n(a,b) = \frac{a^n + b^n}{a^{n-1} + b^{n-1}}$$
(1.9)

All these means defined above have been studied extensively. For more details the interested reader is referred to [1-10]. In [11], the authors established new Čebyš type integral inequalities involving functions whose derivatives belong to L_p spaces via certain integral identities.

In [7,10], the authors defined oscillatory mean and its dual forms. Also, established some new inequalities involving logarithmic mean, identric mean and power mean. Further, the authors also obtained best possible values for the equalities as follows:

Definition 1.1. For a, b > 0 and $\alpha \in (0, 1)$, then Oscillatory mean and its dual form are defined as follows;

$$O(a,b;\alpha) = \alpha G(a,b) + (1-\alpha)A(a,b)$$
(1.10)

and

$$Q^{(d)}(a,b;\alpha) = G(a,b)^{\alpha} A(a,b)^{1-\alpha}.$$
(1.11)

Definition 1.2. For a, b > 0 and $\alpha \in (0, 1)$, then r^{th} Oscillatory mean and its dual form are defined as follows;

$$O(a,b;\alpha,r) = \alpha M_r(a,b) + (1-\alpha)A(a,b)$$
(1.12)

and

$$O^{(d)}(a,b;\alpha,r) = M_r(a,b)^{\alpha} A(a,b)^{1-\alpha}.$$
(1.13)

In [8], K. M. Nagaraja and P. Siva Kota Reddy introduced α -centroidal mean and its dual and studied some important results.

Definition 1.3. For a, b > 0 and $\alpha \in (0, 1)$, α -centroidal mean and its dual form are defined as follows;

$$CT(a,b;\alpha) = \alpha H(a,b) + (1-\alpha)C(a,b)$$
(1.14)

and

$$CT^{(d)}(a,b;\alpha) = H(a,b)^{\alpha}C(a,b)^{1-\alpha}.$$
 (1.15)

and the extended mean values given as below:

$$S_{s,t}(a,b) = \begin{cases} \left(\frac{t(a^s - b^s)}{s(a^t - b^t)}\right)^{\frac{1}{s-t}} & if(s-t)st \neq 0, a \neq b\\ exp\left(-\frac{1}{s} + \frac{a^s loga - b^s logb}{a^s - b^s}\right) & ifs = t \neq 0, a \neq b\\ exp\left(\frac{a^s - b^s}{s(a^s loga - b^s logb)}\right)^{\frac{1}{s}} & ifs \neq 0, t = 0, a \neq b\\ \sqrt{ab} & ifs = t = 0\\ a & ifa = b \end{cases}$$
(1.16)

By the motivation of work done by the authors [8], in this paper we make an attempt to generalize the α -centroidal mean and its dual. In the next section, we established monotonicity results for generalized α -centroidal mean and some inter-related inequalities among several means. In concluding section, some consequent examples are appended. The new initiated mean in this paper is the generalization of oscillatory mean, r^{th} oscillatory mean, α -centroidal mean and some and some other means.

2. Definitions and Properties

In this section, the generalized α -centroidal mean and its dual are defined as follows:

Definition 2.1. For a, b > 0, r, n are real numbers and $\alpha \in (0, 1)$, then the generalized α -centroidal mean and its dual form are defined as follows:

$$CT(a,b;\alpha,r,n) = \alpha M_r(a,b) + (1-\alpha)C_n(a,b)$$
(2.1)

and

$$CT^{(d)}(a,b;\alpha,r,n) = M_r(a,b)^{\alpha} C_n(a,b)^{1-\alpha}.$$
 (2.2)

For $\alpha \in (0,1)$, the generalized α -centroidal mean and its dual satisfy the following properties:

Property 2.2. The generalized α -centroidal mean and its dual are means. That is

$$Min\{a,b\} \le \{CT(a,b;\alpha,r,n), CT^d(a,b;\alpha,r,n)\} \le Max\{a,b\}.$$

Property 2.3. The means $CT(a, b; \alpha, r, n)$ and $CT^{d}(a, b; \alpha, r, n)$ are:

(1) **Symmetric** :

 $CT(a,b;\alpha,r,n) = CT(b,a;\alpha,r,n)$ and $CT^{(d)}(a,b;\alpha,r,n) = CT^{(d)}(b,a;\alpha,r,n)$.

(2) Homogeneous :

 $CT(at, bt; \alpha, r, n) = tCT(a, b; \alpha, r, n)$ and $CT^{(d)}(at, bt; \alpha, r, n) = tCT^{(d)}(a, b; \alpha, r, n)$.

According to Definition 2.1, the following characteristic properties for $CT(a, b; \alpha, r, n)$ and $CT^{(d)}(a, b; \alpha, r, n)$ are straightforward, this result gives that the various means are particular cases of the generalized α -centroidal mean and its dual.

Proposition 2.4. For $\alpha \in (0, 1)$, then

(1) $CT(a, b; \alpha, 0, 1) = O(a, b; \alpha)$ is the oscillatory mean (2) $CT^{d}(a, b; \alpha, 0, 1) = O^{d}(a, b; \alpha)$ is dual oscillatory mean (3) $CT(a, b; \alpha, r, 1) = O(a, b; \alpha, r)$ is the r^{th} oscillatory mean (4) $CT^{d}(a, b; \alpha, r, 1) = O^{d}(a, b; \alpha, r)$ is the r^{th} dual oscillatory mean (5) $CT(a, b; \alpha, -1, 2) = O(a, b; \alpha)$ is the α -centroidal mean (6) $CT^{d}(a, b; \alpha, -1, 2) = O^{d}(a, b; \alpha)$ is the dual α -centroidal mean (7) $CT(a, b; \frac{1}{3}, 0, 1) = H_{e}(a^{2}, b^{2})$ is the Heron mean. (8) $CT^{d}(a, b; \frac{1}{3}, 0, 1) = G^{\frac{1}{3}}(a, b)A^{\frac{2}{3}}(a, b) = O^{d}(a, b; \frac{1}{3}, 0),$ (9) $min(a, b) \leq CT^{(d)}(a, b; \alpha, r, n) \leq CT(a, b; \alpha, r, n) \leq Max(a, b).$ (10) $M_{r}(a, b) \leq CT^{(d)}(a, b; \alpha, r, n) \leq CT(a, b; \alpha, r, n) \leq C_{n}(a, b).$ (11) $CT(a, b; 1, -1, n) = H(a, b) = CT^{d}(a, b; 1, -1, n).$ (12) $CT(a, b; 1, 0, n) = G(a, b) = CT^{d}(a, b; 1, 0, n).$ (13) $CT(a, b; 1, 1, n) = A(a, b) = CT^{d}(a, b; 0, r, 1).$ (14) $CT(a, b; 0, r, 1) = A(a, b) = CT^{d}(a, b; 0, r, 2).$ (15) $CT(a, b; 0, r, 2) = A(a, b) = CT^{d}(a, b; 0, r, 2).$ (16) $CT(a, b; 0, r, \frac{3}{2}) = \frac{a^{3/2} + b^{3/2}}{a^{1/2} + b^{1/2}} = CT^{d}(a, b; 0, r, \frac{3}{2}).$

From the above results, the following remarks are drawn:

Remark 2.5.

(1)
$$CT(a,b;1,-1,n) + CT(a,b;0,r,2) = 2CT(a,b;1,1,n)$$

- $(2) \ CT^{d}(a,b;1,-1,n) + CT^{d}(a,b;0,r,2) = 2CT^{d}(a,b;1,1,n)$
- (3) $CT(a,b;1,0,n) + CT(a,b;0,r,\frac{3}{2}) = 2CT(a,b;1,1,n)$
- (4) $CT^{d}(a,b;1,0,n) + CT^{d}(a,b;0,r,\frac{3}{2}) = 2CT^{d}(a,b;1,1,n).$ (i.e., $H(a,b) + C_{2}(a,b) = G(a,b) + C_{3/2}(a,b) = 2A(a,b)$)

3. Monotonic Results

In this section, the monotonicity and behavior of the generalized α -centroidal mean and its dual are studied.

Theorem 3.1. For $\alpha \in (0,1)$ and for a, b > 0, then $CT^{(d)}(a,b;\alpha,r,n) \leq CT(a,b;\alpha,r,n)$.

Proof. The proof follows from the well known power mean inequality:

$$M_r(a,b) = \begin{cases} \left(\frac{a^r + b^r}{2}\right)^{\frac{1}{r}}, & r \neq 0; \\ \sqrt{ab}, & r = 0. \end{cases}$$

$$(3.1)$$

Lemma 3.2. For a real number r and a, b > 0, we have $M_{r+1}(a, b) \ge M_r(a, b)$.

Theorem 3.3. The generalized α -centroidal mean $CT(a, b; \alpha, r, n)$ is an increasing function with respect to r, for a, b > 0 and $\alpha \in (0, 1)$. That is,

$$CT(a,b;\alpha,r+1,n) \ge CT(a,b;\alpha,r,n)$$
(3.2)

Proof. From Definition 2.1,

$$CT(a,b;\alpha,r+1,n) = \alpha M_{r+1}(a,b) + (1-\alpha)C_n(a,b)$$

$$\geq \alpha M_r(a,b) + (1-\alpha)C_n(a,b)$$

$$= CT(a,b;\alpha,r,n)$$

This completes the proof.

Theorem 3.4. The generalized α -centroidal mean $CT(a, b; \alpha, r, n)$ is an increasing function with respect to r, for a, b > 0 and $\alpha \in (0, 1)$. That is,

$$CT^{d}(a,b;\alpha,r+1,n) \ge CT^{d}(a,b;\alpha,r,n)$$
(3.3)

Proof. From Definition 2.1,

$$CT^{d}(a, b; \alpha, r+1, n) = M_{r+1}(a, b)^{\alpha} + C_{n}(a, b)^{1-\alpha}$$
$$\geq M_{r}(a, b)^{\alpha} + C_{n}(a, b)^{1-\alpha}$$
$$= CT^{d}(a, b; \alpha, r, n).$$

This completes the proof.

Lemma 3.5. For a real number n and a, b > 0, we have $C_{n+1}(a, b) \ge C_n(a, b)$.

Theorem 3.6. The generalized α -centroidal mean $CT(a, b; \alpha, r, n)$ an increasing function with respect to n, for a, b > 0 and $\alpha \in (0, 1)$. That is,

$$CT(a, b; \alpha, r, n+1) \ge CT(a, b; \alpha, r, n)$$
(3.4)

Proof. From Definition 2.1,

$$CT(a,b;\alpha,r,n+1) = \alpha M_r(a,b) + (1-\alpha)C_{n+1}(a,b)$$

$$\geq \alpha M_r(a,b) + (1-\alpha)C_n(a,b)$$

$$= CT(a,b;\alpha,r,n).$$

This completes the proof.

Theorem 3.7. The generalized α -centroidal mean $CT(a, b; \alpha, r, n)$ an increasing function with respect to n, for a, b > 0 and $\alpha \in (0, 1)$. That is,

$$CT^{d}(a,b;\alpha,r,n+1) \ge CT^{d}(a,b;\alpha,r,n)$$
(3.5)

Proof. From Definition 2.1,

$$CT^{d}(a,b;\alpha,r,n+1) = M_{r}(a,b)^{\alpha} + C_{n+1}(a,b)^{1-\alpha}$$

$$\geq M_{r}(a,b)^{\alpha} + C_{n}(a,b)^{1-\alpha}$$

$$= CT^{d}(a,b;\alpha,r,n).$$

This completes the proof.

Theorem 3.8. The generalized α -centroidal mean and its dual are varies from $C_n(a, b)$ to $M_i(a, b)$ with α varies from 0 and 1.

Proof. The proof is follows from Definition 2.1.

Remark 3.9.

For r = n, the generalized α -centroidal mean and its dual are respectively given by:

$$CT(a,b;\alpha,r,r) = \alpha M_r(a,b) + (1-\alpha)C_r(a,b)$$

and

$$CT^{d}(a, b; 1, r, n) = M_{r}(a, b)^{\alpha} + C_{r}(a, b)^{1-\alpha}.$$

Example 3.10.

When r = n = 0, $CT(a, b; \alpha, 0, 0) = \alpha G(a, b) + (1 - \alpha)H(a, b)$, this mean is increasing with request to α , then

$$H(a,b) \le \frac{3H+G}{4} \le \frac{H+G}{2} \le \frac{H+3G}{4} \le G(a,b).$$

Theorem 3.11. For a, b > 0, $\alpha \in (0, 1)$ and n = r, the generalized α -centroidal mean and its dual are increasing with respect to r.

Proof. The proof is follows from Lemma 3.2 and 3.5.

Example 3.12. From Remark 3.9 and Theorem 3.11, the following holds:

$$H(a,b) \le \frac{M_{1/4}(a,b) + C_{1/4}(a,b)}{4} \le \frac{3G+A}{4} \le \frac{M_{3/4}(a,b) + C_{3/4}(a,b)}{4} \le A(a,b).$$

Lemma 3.13. For a real number r and a > b > 0, we have

(1) $M_r(a,b) > C_r(a,b)$ if r < 1(2) $M_r(a,b) < C_r(a,b)$ if r > 1(3) $M_r(a,b) = C_r(a,b)$ if r = 1.

Proof. Put a = t, b = 1, in $M_r(a, b)$ and $C_r(a, b)$. Consider

$$f(t) = \frac{1}{r} [ln(t^{r}+1) - ln2] - [ln(t^{r}+1) - ln(t^{r-1}+1)]$$

then

$$f^{\dagger}(t) = (1-r)(t-1)\frac{t^{r-2}}{(t^r+1)(t^{r-1}+1)} > 0$$

if r < 1, f(t) is increasing (i.e., $M_r(a, b) > C_r(a, b)$). The result is clear.

Theorem 3.14. If a, b > 0, $\alpha \in (0, 1)$ and n = r, then the generalized α -centroidal mean and its dual monotonically increasing with respect to α if r < 1, monotonically decreasing with respect to α if r > 1 and inequality turns out to be equality if r = 1.

Proof. The proof follows from Theorem 3.11 and Lemma 3.13.

4. Some Inequalities

From the definitions of oscillatory mean, the α -centroidal mean and its dual forms for $\alpha \in (0, 1)$ and a, b > 0, the following inequalities hold:

$$CT^{d}(a, b; 1, r, n) \le \dots \le O^{d}(a, b; 1) \le \dots \le O(a, b; 0) \le \dots \le CT(a, b; 0, r, n).$$

By replacing a = t + 1, b = 1 in Definition 2.1, the Taylor's series expansion of $O(a, b; \alpha, k)$ and $O^{(d)}(a, b; \alpha, k)$ are as follows:

$$L(a,b) = L(t,1) = 1 + \frac{t}{2} - \frac{1}{12}t^2 + \dots$$
(4.1)

$$I(a,b) = I(t,1) = 1 + \frac{t}{2} - \frac{1}{24}t^2 + \dots$$
(4.2)

$$H_p(a,b) = H_p(t,1) = 1 + \frac{t}{2} + \frac{2p-3}{24}t^2 + \dots$$
(4.3)

$$M_r(a,b) = M_r(t,1) = 1 + \frac{t}{2} + \frac{r-1}{8}t^2 + \dots$$
(4.4)

$$CT(t,1;\alpha,n,n) = 1 + \frac{1}{2}t + \frac{3n-3}{8}t^2 + \dots$$
(4.5)

$$CT^{(d)}(t,1;\alpha,n,n) = 1 + \frac{1}{2}t + \frac{3n-3}{8}t^2 + \dots$$
 (4.6)

From the above Taylor's series expansions of various means, we compute the following inequalities.

If $a, b > 0, \alpha \in (0, 1)$ and n_1, n_2 are real numbers, then we have following results.

Proposition 4.1. For $n_1 \leq \frac{7}{9} \leq n_2$, the following double inequality holds

 $CT^{(d)}(a,b;\alpha,n_1,n_1) \leq L(a,b) \leq CT(a,b;\alpha,n_2,n_2).$ (4.7)

Further, $n_1 = \frac{7}{9} = n_2$ is the best possible for (4.7).

Proof. From eqs 4.1 to 4.6, we have

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \le L(a,b) \le CT(a,b;\alpha,n_2,n_2)$$

$$\frac{3n_1-3}{4} \le \frac{-1}{12} \le \frac{3n_2-3}{4}.$$
(4.8)

whenever,

Proposition 4.2. For $n_1 \leq \frac{8}{9} \leq n_2$, the following double inequality holds

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \le I(a,b) \le CT(a,b;\alpha,n_2,n_2).$$
 (4.9)

Further, $n_1 = \frac{8}{9} = n_2$ is the best possible for (4.9).

Proof. From eqs 4.1 to 4.6, we have

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \leq I(a,b) \leq CT(a,b;\alpha,n_2,n_2)$$
(4.10)
$$\frac{3n_1-3}{4} \leq \frac{-1}{24} \leq \frac{3n_2-3}{4}.$$

whenever,

Proposition 4.3. For $n_1 \leq \frac{r+2}{3} \leq n_2$, the following double inequality holds

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \le M_r(a,b) \le CT(a,b;\alpha,n_2,n_2).$$
 (4.11)

Further, $n_1 = \frac{r+2}{3} = n_2$ is the best possible for (4.11).

Proof. From eqs 4.1 to 4.6, we have

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \le I(a,b) \le CT(a,b;\alpha,n_2,n_2)$$
 (4.12)

whenever,
$$\frac{3n_1-3}{4} \le \frac{r-1}{8} \le \frac{3n_2-3}{4}$$
.
Or $n_1 \le \frac{r+2}{3} \le n_2$.

Proposition 4.4. For $n_1 \leq \frac{2p+6}{3} \leq n_2$, the following double inequality holds

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \le M_r(a,b) \le CT(a,b;\alpha,n_2,n_2).$$
(4.13)

Further, $n_1 = \frac{2p+6}{3} = n_2$ is the best possible for (4.13).

Proof. From eqs 4.1 to 4.6, we have

$$CT^{(d)}(a,b;\alpha,n_1,n_1) \le I(a,b) \le CT(a,b;\alpha,n_2,n_2)$$
 (4.14)

whenever, Or $\frac{\frac{3n_1-3}{4}}{n_1 \le \frac{2p+3}{3} \le n_2.} \le \frac{3n_2-3}{4}.$

Acknowledgments. We would like to thank referees for their careful review and constructive comments on our manuscript.

References

- 1. P. S. Bullen, *Handbook of means and their inequalities*, Kluwer Acad. Publ., Dordrecht, 2003.
- G. H. Hardy, J. E. Littlewood and G. Pòlya, *Inequalities*, 2nd edition, Cambridge University Press, Cambridge, 1959.
- V. Lokesha, S. Padmanabhan, K. M. Nagaraja and Y. Simsek, Relation between Greek means and various means, *General Mathematics*, 17(3), (2009), 3-13.
- V. Lokesha, K. M. Nagaraja and Y. Simsek, New inequalities on the homogeneous functions, J. Indones. Math. Soc., 15(1), (2009), 49-59.
- V. Lokesha, K. M. Nagaraja, B. Naveen Kumar and S. Padmanabhan Oscillatory type mean in Greek means, *Int. e-Journal of Engg. Maths Theory and Applications*, 9(3), (2010), 18-26.
- K. M. Nagaraja, V. Lokesha and S. Padmanabhan, A simple proof on strengthening and extension of inequalities, *Advn. Stud. Contemp. Math.*, **17**(1), (2008), 97-103.
- V. Lokesha, Zhi-Hua Zhang and K. M. Nagaraja, rth Oscillatory mean for several positive arguments, *International J. of Phy. Sci.*, 18(3), (2006), 519-522.
- K. M. Nagaraja and P. Siva Kota Reddy, α-Centroidal mean and its dual, Proceedings of the Jangjeon Math. Soc., 15(2), (2012), 163-170.
- B. Naveen Kumar, K. M. Nagaraja, A. Bayad and M. Saraj, New means and its properties, Proceedings of the Jangjeon Math. Soc., 14(3), (2010), 243-254.
- S. Padmanabhan, V. Lokesha, M. Saraj and K. M. Nagaraja, Oscillatory mean for several positive arguments, *Journal of Intelligent System Research*, 2(2), (2008), 137-139.
- M. Z. Sarikaya, A. Saglam and H. Yildirim, On generalization of Čebyš type inequalities, Iranian Journal of Mathematical Sciences and Informatics, 5(1), (2010), 41-48.
- G. Toader and S. Toader, Greek means and Arithmetic mean and Geometric mean, RGMIA Monograph, Australia, 2005.