

Case Report

Massive Retinal Pigment Epithelial Detachment Following Acute Hypokalemic Quadriparesis in Dengue Fever

Neha Goel¹, MS, DNB, FRCS; Vishaal Bhambhwani², MS; Pooja Jain², MS; Basudeb Ghosh², MD, MNAMS

¹ICARE Eye Hospital and Postgraduate Institute, Noida, Uttar Pradesh, India

²Guru Nanak Eye Centre, Maulana Azad Medical College, New Delhi, India

Abstract

Purpose: To describe an unusual retinal manifestation of dengue fever in an endemic region.

Case Report: A 35 year old male presenting with acute onset decreased vision in his right eye, was found to have a massive retinal pigment epithelial detachment (PED) extending up to the vascular arcades. He had been diagnosed with acute hypokalemic quadriparesis in dengue fever in the preceding week, which had resolved following treatment. The patient was managed conservatively. At three months follow up, there was spontaneous flattening of the PEDs with improvement in visual acuity.

Conclusion: Dengue fever complicated by acute hypokalemic quadriparesis can be associated with PED, which can be large. The condition resolves spontaneously and bears a good prognosis.

Keywords: Dengue Fever; Flaccid Quadriparesis; Hypokalemia; Pigment Epithelial Detachment

J Ophthalmic Vis Res 2016; 11 (2): 231-233.

INTRODUCTION

Dengue is a major public health problem in the tropics and subtropics.^[1] Ocular involvement in dengue fever is uncommon.^[2,3] Herein, we describe a young man diagnosed with acute hypokalemic quadriparesis due to dengue infection, who presented with unilateral visual impairment and was found to have massive retinal pigment epithelial detachment (PED).

CASE REPORT

A 35-year-old man presented with sudden-onset decreased vision in his right eye since 5 days before.

Correspondence to:

Neha Goel, MS, DNB, FRCS. 57, Sadar Apartments, Mayur Vihar Phase 1 Extension, New Delhi - 11009, India. E-mail: nehadoc@hotmail.com

Received: 04-08-2014

Accepted: 16-02-2015

One week earlier, he had presented to the medical emergency with high grade fever and progressive weakness of all four limbs for 2 days. Based on blood investigations including a positive dengue NS1 antigen, a diagnosis of dengue fever with thrombocytopenia (platelet count 40,000/mm³) and hypokalemic acute flaccid motor paralysis was made. He was started on intravenous potassium chloride infusion. Within 8 hours, his motor power in all limbs improved significantly and serum potassium levels increased from 1.8 meq/liter at presentation, to 4.2 meq/liter by day 2 of treatment. He was discharged after one week of hospitalization, with no neurological sequelae, and presented to the ophthalmology department a few days later. There was no history of hypertension during this period.

On examination, best corrected visual acuity (BCVA) was 20/200 in the right and 20/20 in the left eye. Anterior segments of both eyes were unremarkable,

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Goel N, Bhambhwani V, Jain P, Ghosh B. Massive retinal pigment epithelial detachment following acute hypokalemic quadriparesis in dengue fever. *J Ophthalmic Vis Res* 2016;11:231-3.

Access this article online

Quick Response Code:



Website:
www.jovr.org

DOI:
10.4103/2008-322X.158899

with intraocular pressures measuring 16 mm Hg bilaterally. The pupillary reactions were normal. Fundus examination of the right eye showed a large PED occupying almost the entire macula with linear streaks of altered pigmentation on the surface [Figure 1a]. The left eye showed two PEDs [Figure 1b]. Fluorescein angiography (FA, VISUPAC and FF450 plus, Carl Zeiss Meditec, Inc., Dublin, California, USA) demonstrated intense hyperfluorescence from pooling of dye in the PEDs [Figure 1c and d]. Hypofluorescence was seen corresponding to the linear streaks on the surface of the PED in the right eye. In addition, another small PED was noted superotemporal to the disc in the right eye. Spectral domain optical coherence tomography (SD-OCT, RTVue, Optovue Inc., Fremont, California, USA) confirmed the serous nature of the PEDs [Figure 1e and f]. The massive PED in the right eye also showed a pocket of subretinal fluid at its apex and retinal pigment epithelium (RPE) heaping in the area of the linear streaks.

The patient was a nonsmoker and had no history of steroid intake in any form. Blood pressure monitoring was within normal range. Repeat hemogram including platelet count, serum electrolytes, renal and liver function tests were within normal limits. A conservative

approach was taken and the patient was advised to attend regular follow-up.

Three months following the acute presentation, BCVA in his right eye improved to 20/40 while that of the left eye was maintained at 20/20. Fundus examination showed the faint outline of the collapsed PED in the right eye with linear hypopigmented streaks in the vicinity of the fovea [Figure 2a]. FA showed window defects in the region previously occupied by the PED, with hypofluorescence due to the streaks. The small PED superotemporal to the disc persisted [Figure 2c]. Fundus examination in the left eye appeared unremarkable, however small window defects in the area of the PEDs were present on FA [Figure 2b and d]. SD-OCT confirmed flattening of the PEDs in both eyes with RPE hyperplasia corresponding to the streaks in the right eye [Figure 2e and f]. The patient was advised to maintain regular follow-up and self-check his vision using an Amsler grid.

DISCUSSION

The four serotypes of the dengue virus can cause a wide spectrum of diseases ranging from a nonspecific febrile illness to classic dengue fever, dengue hemorrhagic fever or dengue shock syndrome. Various atypical




Figure 1. Fundus findings at presentation. Massive pigment epithelial detachment (PED) occupying the right macula (a) and two small PEDs in the left eye (b). Fluorescein angiography (c and d) show pooling with hypofluorescent streaks corresponding to altered pigmentation on the PED surface in the right eye (c). Spectral domain optical coherence tomography demonstrates serous PEDs (e and f); PED in the right eye had subretinal fluid at its apex with retinal pigment epithelium hyperplasia (arrow).



Figure 2. After 3 months, flattening of the pigment epithelial detachments (PEDs) was noted clinically (a and b), angiographically (c and d) and on spectral domain optical coherence tomography (e and f) (except one PED superotemporal to the right disc). In the right eye, linear streaks of altered pigmentation persisted, which blocked fluorescence and were likely retinal pigment epithelium hyperplasia (arrows).

manifestations have also been recognized such as neurological involvement.^[4] Acute neuromuscular weakness may occur in dengue due to myositis, Guillain–Barre syndrome or hypokalemia;^[5,6] this remains an uncommonly described entity. The possible mechanism for development of hypokalaemia could be transient self-limited renal tubular defects secondary to infection leading to urinary potassium wasting or intracellular shift of potassium due to increased catecholamine levels in response to the stress of the infection and secondary insulin release.^[6]

Dengue related ocular complications are uncommon; however, an increasing incidence has been reported over the past few years.^[2,3,7] While several ocular features such as subconjunctival hemorrhages, intraretinal hemorrhages and Roth's spots are the manifestations of the thrombocytopenia and resultant bleeding diathesis; the presence of retinal vasculitis, macular edema, anterior uveitis and optic neuritis suggests that an inflammatory or immune-mediated pathophysiology may also be considered after acute dengue infections.^[3,8] The prevalence of dengue maculopathy (macular swelling, yellow spots in the macula and macular hemorrhages) was found to be 10% among seropositive patients hospitalized for dengue fever during an epidemic by Su et al.^[7] The onset of visual impairment usually occurs at the nadir of thrombocytopenia. Prognosis is generally good as the disease is often self-limited.^[3]

Acute onset, reversible, pure motor quadriparesis complicating dengue fever was attributed to hypokalemia in our patient. Other causes of hypokalemia had been ruled out by clinical examination and appropriate investigations, and the patient demonstrated dramatic recovery with potassium supplementation. Visual complaints ensued on the 5th day of the illness. To the best of our knowledge, this is the first report of a large PED following dengue fever complicated by acute hypokalemic quadriparesis. There was spontaneous flattening of the PEDs by 3 months with improvement in BCVA. Considering the risk of choroidal neovascularization, the patient was kept on regular follow-up.

Although more commonly associated with serous retinal detachment, isolated PEDs can occur in systemic

diseases with inflammatory, renal, hematologic or neoplastic origin.^[9] While it may not be possible to attribute a single definitive pathophysiological basis for this occurrence, the putative mechanism for both the hypokalemia and the PEDs might be increased catecholamine levels due to the stress of the infection. Hyperactivity of the adrenomedullary system and elevated levels of catecholamines have been shown to play a role in the pathogenesis of idiopathic central serous chorioretinopathy.^[10] Clinicians, especially in endemic areas, should be aware of such an association and its potential for spontaneous resolution.

Financial Support and Sponsorship

Nil.

Conflicts of Interest

There are no conflicts of interest.

REFERENCES

- Gubler DJ. Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev* 1998;11:480-496.
- Lim WK, Mathur R, Koh A, Yeoh R, Chee SP. Ocular manifestations of dengue fever. *Ophthalmology* 2004;111:2057-2064.
- Chan DP, Teoh SC, Tan CS, Nah GK, Rajagopalan R, Prabhakaragupta MK, et al. Ophthalmic complications of dengue. *Emerg Infect Dis* 2006;12:285-289.
- Gulati S, Maheshwari A. Atypical manifestations of dengue. *Trop Med Int Health* 2007;12:1087-1095.
- Hira HS, Kaur A, Shukla A. Acute neuromuscular weakness associated with dengue infection. *J Neurosci Rural Pract* 2012;3:36-39.
- Jha S, Ansari MK. Dengue infection causing acute hypokalemic quadriparesis. *Neurol India* 2010;58:592-594.
- Su DH, Bacsal K, Chee SP, Flores JV, Lim WK, Cheng BC, et al. Prevalence of dengue maculopathy in patients hospitalized for dengue fever. *Ophthalmology* 2007;114:1743-1747.
- Wen KH, Sheu MM, Chung CB, Wang HZ, Chen CW. The ocular fundus findings in dengue fever. *Gaoxiong Yi Xue Ke Xue Za Zhi* 1989;5:24-30.
- Wolfensberger TJ, Tufail A. Systemic disorders associated with detachment of the neurosensory retina and retinal pigment epithelium. *Curr Opin Ophthalmol* 2000;11:455-461.
- Sun J, Tan J, Wang Z, Yang H, Zhu X, Li L. Effect of catecholamine on central serous chorioretinopathy. *J Huazhong Univ Sci Technolog Med Sci* 2003;23:313-316.