[Downloaded free from http://www.jovr.org on Friday, September 25, 2020, IP: 5.237.102.59]

Letter

Author's Reply

Georgios D. Panos¹, MD; Zisis Gatzioufas², MD, PhD

¹Department of Ophthalmology, Ipswich Hospital NHS Trust, University of Cambridge, Ipswich, Suffolk, UK ²Cornea and Cataract Service, Moorfields Eye Hospital, London, UK

J Ophthalmic Vis Res 2018; 13 (1): 89

Dear Editor,

We would like to thank Dr. Mohan for his comments and for sharing his interesting case with us.

We fully agree that extended retinal nerve myelination combined with high myopia is associated with treatment refractory amblyopia. However, the literature reports positive treatment outcomes in a number of cases of myopia and amblyopia related to unilateral myelinated nerve fibers.^[1,2] According to Hittner and Antoszyk, the status of the macula is the main prognostic factor for final visual outcomes in these cases.^[1] This observation is also supported by Kodama et al.^[3]

In our case, funduscopy did not reveal any obvious macular or optic nerve pathology. Moreover, the baseline best corrected visual acuity was 20/60, which is significantly higher than in the case reported by Dr. Mohan with the patient only being able to count fingers close to face. We believe that the absence of obvious macular and optic disc anomalies is the reason that our patient recovered partially after treatment with glasses and patching of the fellow eye.

Financial Support and Sponsorship

Nil.

Correspondence to:

Georgios D. Panos, MD. Department of Ophthalmology, Ipswich Hospital NHS Trust, Heath Road, Ipswich, Suffolk IP4 5PD, UK. E-mail: gdpanos@gmail.com

Received: 04-05-2017 Accepted: 15-05-2017

Conflicts of Interest

There are no conflicts of interest.

REFERENCES

- Hittner HM, Antoszyk JH. Unilateral peripapillary myelinated nerve fibers with myopia and/or amblyopia. *Arch Ophthalmol* 1987;105:943-948.
- Summers CG, Romig L, Lavoie JD. Unexpected good results after therapy for anisometropic amblyopia associated with unilateral peripapillary myelinated nerve fibers. J Pediatr Ophthalmol Strabismus 1991;28:134-136.
- Kodama T, Hayasaka S, Setogawa T. Myelinated retinal nerve fibers: Prevalence, location and effect on visual acuity. *Ophthalmologica* 1990;200:77-83.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.jovr.org
	DOI: 10.4103/jovr.jovr_232_17

How to cite this article: Panos GD, Gatzioufas Z. Author's Reply. J Ophthalmic Vis Res 2018;13:89.