

A Photo Essay

Intraoperative Optical Coherence Tomography of a Dislocated Intraocular Lens

Sid A. Schechet, MD; Liliya Golas, MD; Seenu M. Hariprasad, MD

The University of Chicago, Department of Ophthalmology and Visual Sciences, Chicago, IL, USA

J Ophthalmic Vis Res 2019; 14 (1): 116-117

PRESENTATION

An 88-year-old female with pseudoexfoliation syndrome presented with sudden vision loss and was found to have a dislocated one piece intraocular lens (IOL) that was resting on the retina. She underwent 23-gauge pars plana vitrectomy, IOL explantation, and anterior chamber IOL insertion. Intraoperative optical coherence tomography (EnFocus iOCT; Leica, Wetzlar, Germany) was used to assess the position and vault of the IOL on the retinal surface [Figure 1a]. Utilization of iOCT enabled the surgeon to safely grasp the IOL with forceps in the location with the greatest vault in order to minimize any potential trauma to the retina [Figure 1b].

DISCUSSION

Intraoperative optical coherence tomography (iOCT) is an exciting new tool in the ophthalmology surgical realm. In the past few years, its clinical value has been demonstrated in both posterior and anterior segment surgeries.^[1,2] The PIONEER study showed that while iOCT adds 4.9 minutes to a procedure, it alters decision making in 48% of lamellar keratoplasty cases and 43% of membrane peeling cases.^[3] Similarly, the

DISCOVER study showed that iOCT altered surgeon decision making in 38% of lamellar keratoplasty cases and 19% of membrane peeling cases.^[4] Our case is the first reported case of using iOCT to successfully explant a dislocated IOL without causing any retinal breaks.

Declaration of Patient Consent

The authors certify that all appropriate patient consent forms were obtained.




Figure 1. (a) Still image from iOCT showing the position of the IOL (yellow arrow) on the retinal surface (green arrow). (b) Still image from iOCT showing the haptic of the IOL being safely grasped by intraocular forceps IOL (yellow arrow).

Correspondence to:

Sid A. Schechet, MD. 4309 Davis Street, Skokie, IL 60076, USA.

E-mail: Schechets@gmail.com

Received: 06-12-2017

Accepted: 05-02-2018

Access this article online

Quick Response Code:



Website:
www.jovr.org

DOI:
10.4103/jovr.jovr_261_17

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Schechet SA, Golas L, Hariprasad SM. Intraoperative optical coherence tomography of a dislocated intraocular lens. *J Ophthalmic Vis Res* 2019;14:116-7.

Archive of SID

Financial Support and Sponsorship

Nil.

Conflicts of Interest

There are no conflicts of interest.

REFERENCES

1. Ehlers JP. Intraoperative optical coherence tomography: past, present, and future. *Eye (Lond)* 2016;30:193-201.
2. Carrasco-Zevallos OM, Viehland C, Keller B, Draelos M, Kuo AN,

Cynthia A. et al. Review of intraoperative optical coherence tomography: technology and applications [Invited]. *Biomed Opt Express* 2017;8:1607-1637.

3. Ehlers JP, Goshe J, Dupps WJ, Kaiser PK, Singh RP, Gans R, et al. Determination of feasibility and utility of microscope-integrated optical coherence tomography during ophthalmic surgery: the DISCOVER Study RESCAN Results. *JAMA Ophthalmol* 2015 Oct;133:1124-1132.
4. Ehlers JP, Dupps WJ, Kaiser PK, Goshe J, Singh RP, Petkovsek D, Srivastava SK. The Prospective Intraoperative and Perioperative Ophthalmic Imaging with Optical Coherence Tomography (PIONEER) Study: 2-year results. *Am J Ophthalmol* 2014 Nov;158:999-1007.

