

Correlates of Mental Health Service Utilization in Married Women in Tehran 2011

Masuod Ahmadzad-Asl MD* , Farnoush Davoudi MD** , Homa Mohammad Sadeghi MD*
Noshin Khademolreza MD*** , Noushin Zarei MD* , Morteza Naserbakht MD, MPH****
Marziyeh Nojomi MD***** , Maryam Rasoulia MD*****

(Received: 16/06/2012; Revised: 6/9/2012; Accepted: 26/2/2013)

Objective: There are disparities in mental health services (MHS) utilization within and between populations and several factors are studied as its potential correlates. Identifying those correlates would help health policy makers to adjust service provision with characteristics of their community. To evaluate demographic, socioeconomic and system correlates of MHS utilization among married women from Tehran, Iran.

Methods: A household survey of 615 married women residents of 22 municipal districts of Tehran selected via a cluster sampling method. All subjects were asked about health services utilization during last one and 12 months as well as need and access for MHS, demographic and socioeconomic factors. Independent correlates of MHS use were determined with logistic regression analysis.

Results: Total 615 women, mean±SE age and duration of marital life of 42.6±0.9 and 22±0.8 years, respectively were selected, rate of MHS utilization during last one and 12 months were 5.2% and 10.1% respectively. 23.6% of women reported having mental illness and 19.3% and 17.9% had need for MHS and access to outpatient health services, respectively. Logistic regression models showed that need for MHS (OR:5.25, 95%CI:2.7-10.1), access to outpatient services (OR:2.17, 95%CI:1.04-4.52), smoking (OR:3.4, 95%CI:1.16-10.2) and crowding index (OR:0.69, 95%CI:0.48-0.99).

Conclusions: Rate of MHS utilization in women are low considering the near to estimated rate of perceived illness. Bridging the gap between perceived illness and need for services, then providing better access to services in areas with higher crowding index and higher rates of smoking in residents should considered in any mental health promotion programs.

Declaration of interest: None.

Citation: Ahmadzad-Asl M, Davoudi F, MohammadSadeghi H, Khademolreza N, Zarei N, Naserbakht M, et al. Correlates of mental health service utilization in married women in Tehran 2011. *Iran J Psychiatry Behav Sci* 2013; 7(1): 51-60.

Keywords: • Access to Services • Health Services Utilization • Mental Health • Perceived Need

Introduction

Mental health services utilization could be an indicator of mental health status in community and met mental health needs. Addressing correlates of utilization helps reduce unmet need for

mental health care (MHC), which might improve health and reduce costs for both physical and mental health care (1-3).

Perceived or documented history of mental and physical illness, as well as severity of current mental illness, are reported to be associated with higher rates of perceived need for care (4).

An understanding of how citizens evaluate their mental health status and need for mental health services may help health care policy makers and managers evaluate and plan more effectively for MHC demographic ones and socioeconomic factors (5).

Increased access may not mean better health outcomes. Assessment of correlates of service utilization in health sector may be useful To ensure resource efficiency and avoid wastage, To assist the establishment of priorities in limited resource environments, To increase the possibilities for reducing health inequities

Affiliations author:*Resident of psychiatry, Mental Health Research Center, Tehran psychiatry institute, Tehran University of Medical Sciences, Tehran, Iran. **Resident of Community Medicine, Department of Community Medicine, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.***Psychiatrist, Mental Health Research Center, Tehran psychiatry institute, Tehran University of Medical Sciences, Tehran, Iran.****Specialist of Community Medicine, Mental Health Research Center, Tehran psychiatry institute, Tehran University of Medical Sciences, Tehran, Iran.*****Professor of Community Medicine. Department of Community Medicine, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran. *****Associate professor of psychiatry. Mental Health Research Center, Tehran psychiatry institute, Tehran University of Medical Sciences, Tehran, Iran.

Corresponding author: Maryam Rasoulia M.D., Mental Health Research Center, Tehran psychiatry institute, Tehran University of Medical Sciences, Sattarkhan Ave., Niayesh St., Mansuri St., No.1, Tehran Psychiatry Institute, Tehran, Iran.
Tel: +98-21-66506862
Fax: +98-21-66506862,
Email:Rasoulia.m@gmail.com

and In general – to provide an appropriate evidence base for health planning (6-8).

Health demands are influenced by demographics; socio-economic status; health services supply; environmental situation – e.g. water, sanitation; nutrition; illness behaviors; life style; perceptions of illness & health expectations and trade-offs (9-10).

Health services utilization conceptualized among proximate factors that lead to health outcomes such as mortality, morbidity and disability (11-12). Several reports delineate the role of demographic factors such as age, sex, education, occupation, and household income/wealth as well as perceived need to mental health services (2) in health services utilization. All of these factors in aforementioned levels are among determinants of health in community (13-15).

There are reports of disparities in mental health utilization among different community members and study of factors leading to these disparities and planning to eliminate disparities in mental health utilization was noted as to priority in MHC system management (16). Identification of correlates of mental health services utilization would help health policy makers and managers to have better view of MHC indices and do more evidence based planning for MHC system (17-19). The aim of this study is to evaluate demographic and socioeconomic correlates of MHC utilization along with other correlates such as need for MHC and access to MHC among married women from Tehran, Iran.

Materials and Methods

This manuscript shows findings from “study of service utilization, depression and anxiety with domestic violence among married women in Tehran, Iran” project as a part of two psychiatric specialty dissertations of M.A.A and N.Z that is also granted project no. 89-04-121-12426 from MHRC, TUMS.

Design and setting

This study is a cross sectional household survey among married women from all 22 districts of Tehran metropolitan. Tehran, the capital city of Iran has about 8.5 million

populations in 2011 that contains about 11 percent of population of Iran. Tehran has 22 municipal districts each with different socio-cultural status. There are 370 neighborhoods in these 22 districts.

Sampling

We used 42 neighborhoods as sampling clusters each with 15 samples and then based on proportional population in each district we assigned between 1-3 clusters to each districts and then based on random number table we selected sampling units. In each cluster with clockwise turn and systematic random sampling method 15 residential units were selected and then interview with married women (the youngest one if there were more) of the unit was done about study measurements. If there were no response to ring or refusal to participate the next nearest unit were replaced. Among 630 initial samples finally we enrolled 615 completed samples in the study.

Data collection

Study data were collected by direct interview with married women in selected residential units. Interview was done at morning, noon, and afternoon hours proportionally to be assured about participation of all women with different status of presence at home. If there were no response on first ring the interviewer would come back during next three hours and if no response again, the next consecutive residential unit would replace. In case of refusal to participate, two efforts would take to encourage participation by explaining the valence of research in women’s health promotion and exploring possible barrier of participation and try to overcome it by giving information and assurance. If refusal continues, the next residential unit would be replaced.

Five interviewers were selected that were graduate bachelors and master’s degree of psychology and sociology. They trained in a two-hour workshop about concept and method of project, sampling and data collection to assure consistency and reliability. Five percent of samples were re-

evaluated by an independent supervisor to confirm validity of sampling and interview process.

Main measurements

Participants asked about Health service needs; i.e. need for outpatient medical and consultation services during last month and need for inpatient services during last 12 months and the frequency that those needs occur; Service utilization; i.e. frequency of visiting a general practitioner (GP), psychiatrist, consultant/psychologists and receiving non-prescribed medications from pharmacy during last month and last 12 months. Mental health services (MHS) were defined as utilizing either psychiatrists or consultant/psychologists services.

Demographic and socioeconomic variables also were asked from participants: age, education, job, health insurance status, household income, crowding index (number of households per room), smoking and substance use in past year (in subject and her spouse) and perceived mental and/or physical illness status (in subject and her spouse).

Statistics and data analysis

The analysis was done by SPSS for Windows 16.0 (SPSS Inc, Chicago, ILL). After initial data entry, an independent operator qualified data entry. Descriptive analysis revealed mean, frequency and standard error (SE), the correlated of mental health services used were defined by logistic regression analysis including demographic and socioeconomic factors as independent variable in the logistic regression model. In all analyses $p < 0.05$ was considered significant.

We graphed distribution of MHC utilization indices in Tehran area by Arc view 3.2 (ESRI Inc.) to show geographic distribution.

Results

In a household survey, we enrolled 615 married women, residing in 22 municipal districts of Tehran with cluster sampling method. Mean (\pm SE) age of women and their spouses were 42.6 ± 0.9 and 47.8 ± 0.6 years old, respectively. Mean marital life duration

in current relationship was 22 ± 0.8 years. Age at marriage was 19.7 ± 0.2 years and 96.1% of participants reported that they are living their first marriage (and first marriage of their spouse in 92.5%). Education level high school and lower, high school diploma and college and higher were reported in 35.2%, 42.3% and 22.4% respectively and the corresponding educational level in spouses were 36.5%, 35.8% and 27.7%, respectively. About job status 82.4% were house workers for their own, 10.3% employed, 2.5% unemployed and 3.1% retired. The rate of employment in spouses was 89.2%, while 8.0% were unemployed, and 2.0% retired.

Table 1 has summarized demographic and socioeconomic factors of subjects.

Crowding index was 1.87 ± 0.09 and mean household members were 3.97 ± 0.05 person.

Subjects claimed themselves having mental and physical illness in 23.4% and 23.6% respectively and overall perceived mental/physical illness was reported in 38.5% of subjects. While reported mental, physical and either illnesses in spouses were 12.8%, 20.3% and 27.8% respectively. Only 3.7% of subjects reported having disabled/chronic diseased child in home.

Smoking, alcohol and substance use during last 12 months was found in 3.4%, 0.2% and 0.7% of women, respectively. Smoking, alcohol use and substance use in spouses during last 12 months was reported in 25.5%, 0.8% and 4.2%, respectively.

Health insurance and access factors

Among study subjects, 82.4% had health insurance coverage (52.2% social security insurance, 24.6% health services insurance) and 33.7% had complementary health insurance coverage. Perceived need for medical outpatient services and mental health services and inpatient services reported in 42.1%, 19.3% and 14.5% of subjects, respectively. Access to outpatient services in different degrees reported in 17.9% of subjects.

Mental Health services utilization and its correlates

Table 2 delineates services utilization during last month and last 12 months in study subjects. Rate of MHS utilization during last one and 12 months were 5.2% and 10.1% respectively. Women utilized MHS, were about 14.4 ± 3.8

years older than others (mean age 56.2 ± 13 vs. 41.8 ± 0.5 years old, $p<0.01$) and age of spouse of women visiting consultants were significantly younger than others (44.3 ± 2.2 vs. 48.1 ± 0.5 years old, $p<0.05$).

Table 1. Demographic, socioeconomic and health correlates of services utilization during last 12 months in women, Tehran, Iran, 2011-12

	Total [§]	Selected health services				
		Mental health			GP	Any
		Psychiatrist	Consultant	Any		
Total, n (%)	615 (100)	37 (6.0%)	43 (7.0%)	62 (10.1%)	465 (73.8%)	555 (90.2%)
Age; years old, mean \pm SE	42.6 \pm 0.9	56.2 \pm 13 [†]	39.1 \pm 1.9	47.8 \pm 7.8 [‡]	42.8 \pm 1.2	42.7 \pm 1.0
Education level,						
High school and lower	35.2%	6.0%	4.2%	8.3%	75.5%	88.9%
High school diploma	42.3%	5.4%	6.5%	9.2%	72.7%	89.2%
College and higher	22.4%	7.3%	12.4% [†]	14.6%	73.7%	94.9%
Job,						
House-worker employed	82.4%	6.1%	6.7%	10.1%	74.5%	81.7%
unemployed	10.3%	4.3%	11.4% [†]	11.4%	72.9%	91.4%
retired	2.5%	13.3% [†]	0%	13.3%	66.7%	93.3%
Household's monthly income,						
Less than 4*1,000,000 RIs	3.1%	4.5%	4.5%	4.5%	68.2%	100%
4-7*1,000,000 RIs	23.7%	5.5%	5.5%	8.9%	72.6%	86.3%
More than 7*1,000,000 RIs	38.4%	7.6%	5.9%	10.2%	77.5%	92.4%
Not declared	24.9%	5.2%	9.8%	11.8%	67.3%	90.8%
First marriage,	13.0%	3.8%	7.5%	8.8%	77.5%	90.0%
Marriage duration, years, mean \pm SE	96.1%	5.9%	6.9%	10.2%	74.3%	90.5%
No. of rooms in house, mean \pm SE	22 \pm 0.8	23.2 \pm 2.2	17.8 \pm 2.1	20.8 \pm 1.9	21.6 \pm 0.7	21.5 \pm 0.6
Crowding index, mean \pm SE	2.44 \pm 0.04	2.4 \pm 0.15	2.4 \pm 0.04	2.4 \pm 0.1	2.4 \pm 0.04	2.4 \pm 0.04
No. of children, mean \pm SE	1.87 \pm 0.09	1.8 \pm 0.2	1.6 \pm 0.1	1.7 \pm 0.1	1.9 \pm 0.05	1.9 \pm 0.05
Personal living place	3.3 \pm 0.08	3.3 \pm 0.3	2.9 \pm 0.3	3.3 \pm 0.1	3.3 \pm 0.1	2.35 \pm 0.07
Having motor vehicle	58.2%	5.9%	6.4%	9.8%	73.2%	90.2%
Physical illness,	66.2%	5.7%	6.6%	9.6%	72.2%	89.7%
Mental illness,	23.4%	4.8%	6.2%	9.0%	79.3 [†]	90.3%
Illness (mental/physical),	23.6%	15.3% [‡]	11.8% [†]	21.5% [‡]	82.6% [‡]	92.4%
Smoking [‡] ,	38.%	10.5% [‡]	9.3%	15.6% [‡]	78.5% [†]	90.7%
Substance use [‡] ,	3.4%	19.0% [†]	19.0% [†]	28.6% [‡]	71.4%	85.7%
Need for medical services	0.7%	0%	0%	0%	100%	100%
Need for mental health services	42.7%	7.3%	10.0% [†]	13.1% [†]	78.0%	91.9%
Access to outpatient services	19.3%	14.3% [‡]	17.6% [‡]	23.5% [‡]	79.8%	89.1%
Spouse	17.9%	8.2%	9.1%	16.4% [†]	75.5%	90.9%
Age; years old, mean \pm SE	47.8 \pm 0.6	50.2 \pm 2.5	44.3 \pm 2.2 [*]	46.9 \pm 1.8	47.6 \pm 0.6	47.8 \pm 0.6
Physical illness,	20.3%	8.8%	6.4%	10.4%	75.2%	92.8%
Mental illness,	12.8%	12.7% [†]	16.5% [‡]	17.7% [†]	77.2%	94.9%
Illness (mental/physical),	27.8%	9.4% [†]	9.9%	12.9%	76.0%	93.6%
Smoking [‡] ,	25.5%	8.9%	10.2%	14.6% [†]	72.6%	91.7%
Substance use [‡] ,	4.2%	3.8%	7.7%	11.5%	73.1%	84.6%

[†] Values in this column denote value in each subgroup among all study subjects. [‡]GP. General practitioner. [§] "Any" in selected health services denotes all health services (more than mentioned list) utilization in the table during last 12 months. ^{*}Significant at 0.05 level, lower in test group; [†]Significant at 0.05 level, higher in test group; [‡]Significant at 0.01 level, higher in test group; [¶] during last 12 months.

Table 2. Frequency of service utilization among married women in Tehran, Iran, 2011-12

	Service Type	Mean \pm SE	No. of Visits	% of all health services	No. Visits per 1000 population (95%CI [†])
Last month	General Practitioner	0.65 \pm 0.05	397	24.6%	646 (627-664)
	Psychiatrist	0.05 \pm 0.01	32	2.0%	52 (48-56)
	Consultant/psychologist/psychoanalyst	0.04 \pm 0.01	27	1.7%	44 (41-47)
	Pharmacy, non-prescribed medication use	0.82 \pm 0.05	504	31.2%	820 (808-831)
	Mental health services	0.10 \pm 0.02	59	3.7%	96 (89-103)
Last 12 months	General Practitioner	2.7 \pm 0.1	1655	38.0%	2691 (2331-3051)
	Psychiatrist	0.19 \pm 0.04	114	2.6%	185 (173-197)
	Consultant/psychologist/psychoanalyst	0.21 \pm 0.05	131	3.0%	213 (200-226)
	Mental health services	0.40 \pm 0.07	245	5.6%	395 (376-414)

[†] CI: Confidence Interval

Smoking women had more visits with psychiatrists (19% and 5.6% of smoking and non-smoking women, respectively) and consultants (19% and 6.6% of smoking and non-smoking women, respectively) during last 12 months (but not last months); $P < 0.05$. Smoking women utilized more mental health services (28.6%) compared to non-smoking women (9.5%); $p < 0.01$.

Spouse smoking and substance use was correlated to more pharmacy use by women (62.5% vs. 39.3% in women with substance use and substance null spouse and 48.1% and 37.6% in women with smoking and non-smoking spouse, and respective, $p < 0.05$). Other services utilization did not differ based on smoking and substance use in women and their spouses.

During last month subjects with mental illness utilized 14.6% more GP services (50.7% vs. 36.1% in last month, in subject with and without mental illness, $p < 0.01$), 10.2% more psychiatrist services (11.1% vs. 0.9% in subject with and without mental illness, $p < 0.001$), 3.5% more consultant services (5.6% vs. 2.1% in subject with and without mental illness, $p < 0.05$) but not pharmacy services, (44.8% vs. 38.8% in subject with and without mental illness, $p < 0.01$). MHS utilized in 14.6% of subjects that reported having mental illness.

During last month also subjects with reported physical illness compared to subjects without physical illness also utilized more GP services (47.7% vs. 34.4%, respectively; $P < 0.001$), Psychiatrist services (7.2% vs. 1.1%, respectively; $p < 0.001$) but not consultant services (4.2% vs. 2.1%, respectively; $p > 0.1$). Among subjects with physical illness reported more utilization of MHS during last month (9.7% vs. 2.4%, respectively; $p < 0.01$).

Services utilization during last 12 months also showed significant differences between subjects with and without mental/physical illness (Table 1) subject with mental illness utilized 11.6% more GP services

(82.6% vs. 71.0%, respectively; $p < 0.01$), 12.3% more psychiatry services (15.3% vs. 3.0%, respectively; $p < 0.001$), 6.2% more consultant services (11.8% vs. 5.6%, respectively; $p < 0.05$) and 15.1% more MHS (21.5% vs. 6.4%, respectively; $p < 0.001$).

Considering service utilization in last 12 months, having physical illness showed no significant correlation with different group of services utilization (in all cases $p > 0.05$). Subjects with physical illness utilized 8.1% more GP services ($p > 0.05$), 1.4% less psychiatry services ($p > 0.05$), 1.0% less consultant services ($p > 0.05$) and 1.6% more all services ($p > 0.05$).

Subjects with college and higher education level reported more consultant service use (12.4%, $p < 0.05$) but not psychiatry or other groups of service use. Employment status, marital life duration, being in first marriage, showed no correlation with service use in different groups. Household's income level and crowding index showed no significant correlation with different group of services utilization.

Independent correlates of mental health services utilization

Logistic Regression models revealed that perceived need for mental health services (Odds ratio; OR: 5.25), perceived having mental illness (OR: 2.29), smoking (OR: 3.44), access to outpatient health services (OR: 2.17) and crowding index (OR: 0.69) are independent determinants of mental health services utilization in married women in Tehran. Age, age of spouse, physical illness, education level, job status, household's income, spouse's mental/physical illness and smoking/substance use showed no Independent correlation with mental health services utilization (Table 3).

Geographic distribution of mental health services use in Tehran municipal districts and distribution of perceived having illness and smoking/substance use status in Tehran have plotted in figures 1 and 2.

Table 3. Logistic regression model of correlates of mental health services use among married women in Tehran, 2011-12

	Total	OR [†] (95% CI [‡])	P. value
Need for MH [§] services, Perceived	19.3%	5.25 (2.72-10.11)	<0.001
Age	42.6±0.9	1.01 (0.99-1.03)	0.190
Mental illness, Perceived	23.6%	2.29 (1.16-4.53)	0.017
Smoking	3.4%	3.44 (1.16-10.20)	0.026
Spouse's Age	47.8±0.6	0.98 (0.93-1.03)	0.375
Crowding index	1.87±0.09	0.69 (0.48-0.99)	0.043
Spouse's Smoking	25.5%	1.46 (0.67-3.19)	0.340
Access to outpatient services, Perceived	17.9%	2.17 (1.04-4.52)	0.040
Constant	-	0.154	0.002

†. Odds Ratio; ‡. CI: Confidence Interval; §. Mental Health; ||. Crowding index: ratio of household's members to rooms.




Figure 1. Distribution of mental health services utilization and perceived mental/physical illness in women and their spouses in Tehran municipal district, 2011-12




Figure 2. Distribution of mental health services utilization and smoking/substance use during last 12 months in women and their spouses in Tehran municipal district, 2011-12

Discussion

Our study showed several determinants of mental health utilization and disparity and frequency of MHS utilization in married women in Tehran. Study of health service utilization could help policy makers, managers and directors in health sector. These data are useful in determining outreach populations, correlates of services use, over/under use of services based on global standards and unmet needs, and estimation of resources need for service provision (20-21).

Service utilization depends on several factors: demographic factors (age, marital status, family width, education level, employment, and socioeconomic status), insurance and social support systems, access to services, acceptability of services, and need for service (perceived/real) (22-23).

Our findings showed relatively low rate of mental health services utilization among women in Tehran that would reflect also an optimistic picture of MHS utilization in Iran as well. Only 10.1% of women have utilized psychiatric and/or psychological services in last year, considering perceived mental illness in 23.4% and need for MHS in 19.3%, it is clear that there is large gap of unmet need for services in MH system. Rate of MHS use have reported in wide range between 6.5% (in elderly) and 34.5% (in college students) during last year (20, 24-25). Subjects with mental illnesses (26-27), living in stressful conditions such as immigrants (28-29) and victims of crimes such as rape or violence reported higher rates (between 52%-72.5% in victims of rape (30-31), 37% in victims of intimate partner violence (32-33). Need for MHS in our study were similar to study in California in 2005 (34).

About 76.5% of women who declared need for MHS did not use any MH facility in last year, on the other hand need for MHS was the most potent predictor of MHS use in women (OR: 5.25). This finding emphasizes both planning to need management programs and bridging gap of unmet need in MH system promotion. Note that only 32.6% of women who had mental illness had perceived need of MHS (15.2% in women without mental

illness, $p < 0.001$) and the gap between perceived illness and need for services should be considered as a point of intervention in MH system promotion.

Need for MHS in women estimated to be more than 1.5 times than men (34), so it could be estimated about 13% need for MHS in men. Noorbala et al. reported the rate of mental illness in women and men in Tehran in 2009, 37.9% and 28.6%, respectively (35). So we may have about 15-18% gap between illness and need for service in Tehran and considering that about 67.4% of women with perceived illness did not have any perceived need for MHS, this gap between health condition and need for services would be even more. So it seems that health system should plan to overcome this gap. Need for service is one of primary steps in the way of reaching fulfilled needs and mental health promotion programs. Comparing to Noorbala et al. (35), that used the same geographic area for sampling with our study and similar sampling method, the perceived having mental illness (23.4% of women) and real mental health condition (37.9% of women with mental illness) also shows a significant wide gap that needs attention by mental health authorities [36]. This may be highlighted by the reports that the trend of mental illness seems to have an incremental pattern in Tehran during the last decade (37).

The rate of need fulfillment and service use in MH care varies in different countries and among different populations [9, 38-39]. In our study, the disparity of MHS utilization in Tehran geographic areas varied between about zero to 26% (average 10.1%) and this reflects inequities in MHS provision and marketing programs that have been pointed as first priority of MH system to overcome (3, 16, 25-26, 38, 40).

In this study, we found perceived need for MHS, access to outpatient health services, smoking, perceived mental illness, and a socioeconomic determinant i.e. crowding index to be correlated with mental health services utilization and these items could be used in planning mental health promotion programs in Iran. Socioeconomic factors have already

been reported as one of major correlates of MHS use in literature (20, 31, 41-44).

Providing better access to services in areas with higher crowding index and higher rates of smoking in residents as well as need management programs to overcome gaps between perceived illness and need for services as well as unmet need for MHS should be considered in any mental health promotion programs.

Limitations

In this study, we did not assess some predisposing factors (e.g. attitude to MH), and need factors (e.g. mental health condition, severity of illness) on service utilization. Method of study that gathered data from community while conforming several advantages (such as better sampling coverage, community based design, and more real view of health condition of community) simultaneously suffers from some disadvantages such as sampling biases and issues on accuracy of reports.

Authors' Contribution

MAA, MR and NZ conceived and designed the evaluation and helped to draft the manuscript. MAA, MR, FD and HM participated in executing the evaluation, collected the clinical data, interpreted them and performed the statistical analysis. FD, HM, NK, MNb and MN re-evaluated the clinical data and revised the manuscript. All authors read and approved the final manuscript.

This paper is based on results from two dissertations for psychiatry specialty degree of MAA and NZ and grant no. 89-04-121-12426 of MR from MHRC, TUMS.

References

1. Meadows, G.N. and I. Bobevski, Changes in met perceived need for mental healthcare in Australia from 1997 to 2007. *Br J Psychiatry* 2011; 199(6): 479-84.
2. Flisher AJ, Kramer RA, Grosser RC, Alegria M, Bird HR, Bourdon KH, et al. Correlates of unmet need for mental health services by children and adolescents. *Psychol Med* 1997; 27(5): 1145-54.
3. Archer KR, Castillo RC, MacKenzie EJ, Bosse MJ; LEAP Study Group. Perceived need and unmet need for vocational, mental health, and other support services after severe lower-extremity trauma. *Arch Phys Med Rehabil* 2010; 91(5): 774-80.
4. Berren MR, Santiago JM, Zent MR, Carbone CP. Health care utilization by persons with severe and persistent mental illness. *Psychiatr Serv* 1999; 50(4): 559-61.
5. Jang Y, Chiriboga DA, Herrera JR, Tyson DM, Schonfeld L. Attitudes toward mental health services in Hispanic older adults: the role of misconceptions and personal beliefs. *Community Ment Health J* 2011; 47(2): 164-70.
6. Browne GB, Arpin K, Corey P, Fitch M, Gafni A. Individual correlates of health service utilization and the cost of poor adjustment to chronic illness. *Medical Care* 1990; 28(1): 43-58.
7. Sevilla-Dedieu C, Kovess-Masfety V, Gilbert F, Vilagut G, Koenig HH, Bruffaerts R, et al. Mental health care and out-of-pocket expenditures in Europe: results from the ESEMeD project. *J Ment Health Policy Econ* 2011; 14(2): 95-105.
8. Grimes KE, Schulz MF, Cohen SA, Mullin BO, Lehar SE, Tien S. Pursuing cost-effectiveness in mental health service delivery for youth with complex needs. *J Ment Health Policy Econ* 2011; 14(2): 73-83.
9. Zuvekas SH, Fleishman JA. Self-rated mental health and racial/ethnic disparities in mental health service use. *Med Care* 2008; 46(9): 915-23.
10. Soller M, Kharrazi N, Prentiss D, Cummings S, Balmas G, Koopman C et al. Utilization of psychiatric services among low-income HIV-infected patients with psychiatric comorbidity. *AIDS Care* 2011; 23(11): 1351-9.
11. MC Donald A, et al. Effect of QUEBEC Medicine on physician consultation for selected symptoms. *New England J Med* 1974; 290: 649-52.

12. Anderson GM, Brook R, Williams A. A comparison of cost-sharing versus free care in children: effects on the demand for office-based medical care. *Med Care* 1991; 29(9): 890-8.
13. Andersen R, Newman JF. Societal and individual determinants of medical care utilization in the United States. *Milbank Mem Fund Q Health Soc* 1973; 51: 95-124.
14. Donabedian A. Aspect of healthcare care administration. Specify recruitments for health care. Cambridge: Harvard University press; 1973.
15. Andy LA, Sellers C, Andersen RM. Potential for local health survey Astute -of-the art summery.american. *Am J Public Health* 1981; 71(8): 835-40.
16. Cook BL, McGuire T, Miranda J. Measuring trends in mental health care disparities, 2000–2004. *Psychiatr Serv* 2007; 58(12): 1533-40.
17. Elhai JD, Ford JD. Correlates of mental health service use intensity in the National Comorbidity Survey and National Comorbidity Survey Replication. *Psychiatric Serv* 2007; 58(8): 1108-15.
18. Rabinowitz J, Gross R, Feldman D. Correlates of a perceived need for mental health assistance and differences between those who do and do not seek help. *Soc Psychiatry Psychiatr Epidemiol.* 1999; 34(3): 141-6.
19. Alegría M, Mulvaney-Day N, Woo M, Torres M, Gao Sh, Oddo V. Correlates of past-year mental health service use among Latinos: Results from the National Latino and Asian American Study. *Am J Public Health* 2007; 97(1): 76–83.
20. Garrido MM, Kane RL, Kaas M, Kane RA. Use of mental health care by community-dwelling older adults. *J Am Geriatr Soc* 2011; 59(1): 50-6.
21. Girma E, Tesfaye M. Patterns of treatment seeking behavior for mental illnesses in Southwest Ethiopia: a hospital based study. *BMC Psychiatry* 2011; 11: 138.
22. Sareen J, Jagdeo A, Cox BJ, Clara I, ten Have M, Belik SL, et al., Perceived barriers to mental health service utilization in the United States, Ontario, and the Netherlands. *Psychiatric Services* 2007; 58(3): 357-64.
23. Golberstein E, Eisenberg D, Gollust SE. Perceived stigma and mental health care seeking. *Psychiatr Serv* 2008; 59(4): 392-9.
24. Eisenberg D, Hunt J, Speer N, Zivin K. Mental health service utilization among college students in the United States. *J Nerv Ment Dis* 2011; 199(5): 301-8.
25. Cairney J, Corna LM, Streiner DL. Mental health care use in later life: results from a national survey of Canadians. *Can J Psychiatry* 2010; 55(3): 157-64.
26. Mendenhall AN, Demeter C, Findling RL, Frazier TW, Fristad MA, Youngstrom EA, et al. Mental health service use by children with serious emotional and behavioral disturbance: results from the LAMS study. *Psychiatr Serv* 2011; 62(6): 650-8.
27. Evans-Lacko SE, dosReis S, Kastelic E, Riley AW. Mental health service use before and after diagnosis of early-onset bipolar disorder. *J Behav Health Serv Res* 2011; 38(3): 398-413.
28. Simning A, Richardson TM, Friedman B, Boyle LL, Podgorski C, Conwell Y. Mental distress and service utilization among help-seeking, community-dwelling older adults. *Int Psychogeriatr* 2010; 22(5): 739-49.
29. Chen J, Vargas-Bustamante A. Estimating the effects of immigration status on mental health care utilizations in the United States. *J Immigr Minor Health* 2011; 13(4): 671-80.
30. Amstadter AB, Zinzow HM, McCauley JL, Strachan M, Ruggiero KJ, Resnick HS, et al. Prevalence and correlates of service utilization and help seeking in a national college sample of female rape victims. *J Anxiety Disor* 2010; 24(8): 900-2.
31. Alonzo DM, Harkavy-Friedman JM, Stanley B, Burke A, Mann JJ, Oquendo MA. Predictors of treatment utilization in major depression. *Arch Suicide Res* 2011; 15(2): 160-71.
32. Ahmed AT, BR McCaw. Mental health services utilization among women

- experiencing intimate partner violence. *Am J Manag Care* 2010; 16(10): 731-8.
33. Elklit A, M Shevlin. General practice utilization after sexual victimization: a case control study. *VAW* 2010. 16(3): 280-90.
34. Grant D, Kravitz-Wirtz N, Aguilar-Gaxiola S, Sribney WM, Aydin M, Brown ER. Mental health status and use of mental health services by California adults. *Policy Brief UCLA Cent Health Policy Res* 2010 ; 6: 1-8.
35. Noorbala A, Bagheri Yazdi S, Asadi Lari M, Vaez Mahdavi M. [Mental Health Status of Individuals Fifteen Years and Older in Tehran-Iran (2009)]. *Iran J Psychiatry Clin Psychol* 2011. 16(4): 479-83. Persian.
36. WHO, C.O.i.t.I.R.o.I., WHO-AIMSREPORT ON MENTAL HEALTH SYSTEM IN THE ISLAMIC REPUBLIC OF IRAN. 2006, World Health Organization: Tehran, the Islamic Republic of Iran.
37. Noorbala AA, Bagheri Yazdi SA, Hafezi M. Trends in Change of Mental Health Status in the Population between 1998 and 2007. *Arch Iran Med* 2012;15(4): 201-4.
38. Sturm, R., J.S. Ringel, and T. Andreyeva, Geographic disparities in children's mental health care. *Pediatrics* 2003; 112(4): 308.
39. Merikangas KR, He JP, Burstein M, Swendsen J, Avenevoli S, Case B, et al. Service utilization for lifetime mental disorders in U.S. adolescents: results of the National Comorbidity Survey-Adolescent Supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry*, 2011. 50(1): 32-45.
40. Cummings JR, Druss BG. Racial/ethnic differences in mental health service use among adolescents with major depression. *J Am Acad Child Adolesc Psychiatry* 2011. 50(2): 160-70.
41. Wang Y, Li B, Pan J, Sengupta S, Boland Emrick C, Cohen MS, et al., Factors associated with utilization of a free HIV VCT clinic by female sex workers in Jinan City, Northern China. *AIDS Behav* 2011. 15(4): 702-10.
42. Low LF, Anstey KJ, Lackersteen SM, Camit M. Help-seeking and service use for dementia in Italian, Greek and Chinese Australians. *Aging Ment Health* 2011; 15(3): 397-404.
43. Lu MW, Duckart JP, O'Malley JP, Dobscha SK. Correlates of utilization of PTSD specialty treatment among recently diagnosed veterans at the VA. *Psychiatr Serv*, 2011. 62(8): 943-9.
44. Eisenberg D, Golberstein E, Gollust SE. Gollust, Help-seeking and access to mental health care in a university student population. *Med Care*, 2007. 45(7): 594-601.