

Male Sexual Dysfunction Before and After Lower Limb Trauma

Abbasali Ebrahimian,¹ Hossein Davari,^{1,*} and Maedeh Tourdeh¹

¹Nursing Care Research Center, Semnan University of Medical Sciences, Semnan, IR Iran

*Corresponding author: Hossein Davari, Nursing Care Research Center, Basij Street, Semnan, Iran. Tel: +98-2333437825, E-mail: hosseindavari30@yahoo.com

Received 2016 June 02; Revised 2016 November 20; Accepted 2017 February 09.

Abstract

Background: Sexual dysfunction, which may occur after lower extremity trauma, is a main physical problem with negative mental health impacts.

Objectives: This study aimed at comparing male sexual dysfunction before and after lower extremity trauma.

Methods: In this cross sectional study, all patients with lower extremity trauma who were admitted to Kowsar hospital in Semnan, Iran, were recruited and assessed. Two main questionnaires were administered: a questionnaire including demographic and incident variables and the international index of erectile dysfunction (IIEF). Sampling was done from June 2014 to August 2015. Statistical analyses were performed using Friedman and Wilcoxon's test in SPSS Version 19.

Results: Overall, 153 male patients participated in the study. The changes of sexual dysfunctions were significantly different before and after the incident in 1 and 3 months after the incident in all the IIEF subscales ($P = 0 < 0001$). Moreover, within 1 to 3 months after the trauma, a significant difference was found only in intercourse satisfaction subscale ($P < 0.05$).

Conclusions: Males with lower extremity trauma are at risk of sexual dysfunction, which may continue for months after the trauma. Therefore, health care providers should evaluate and treat sexual dysfunction in patients after the lower extremity trauma.

Keywords: lower Extremity, Sexual Dysfunction, Trauma

1. Background

Trauma is considered as one of the main causes of death and disability worldwide (1, 2) and the leading cause of death (3). Trauma is the second leading cause of death in Iran after cardiovascular diseases (4).

According to the national trauma data bank in 2014, lower extremity trauma with a prevalence of 36.99% was the most common type of trauma. The death and mortality arising from this trauma has been 15.39% (5). A large number of individuals with lower extremity trauma survive, but a significant number of them are affected by physical disabilities and mental problems during the treatment process and even after it (6, 7). The most common mental disorders after organs trauma are posttraumatic stress syndrome, depression, anxiety, mental distress, and alcohol abuse (8). Mental disorders causing such a trauma usually diminish over time. A longitudinal study indicated that the prevalence of mental disorders in individuals with lower limb trauma was reduced from 48% to 42% during 3 to 12 months after the incident (9). Seaborg et al. (2012) found that the severity of physical and mental disorders in patients with trauma is reduced after 5 years of trauma, but the prevalence of these disorders is still significantly more than the other individuals in the society (10). Soberg (2015) found that 10 years after the trauma, the prevalence of mental disorders in injured individuals was similar to others in the society, but their physical health was worse

(11). Therefore, paying sufficient attention to physical and mental disorders is necessary even for several years after the trauma.

Sexual function is one of the main physical and mental concerns of each individual (12) that may be impaired after lower extremity trauma (13). Physical and mental disorders are also considered the major causes of sexual dysfunction (14, 15). Shulman et al. (2015) reported the prevalence of sexual dysfunction in lower extremity trauma to be 11% to 47% in the first 3 months after the incident and about 20% in the first year (16). Harvey-Kelly (2014) also reported the prevalence of sexual dysfunction to be 52.1% in males with pelvic trauma. They also showed that all aspects of these individuals' sexual function were reduced significantly (17).

The patients' physical problems and disabilities are usually most considered during the treatment of lower extremity trauma and the treatment of their mental disorders, but their sexual problems are not much considered. However, the prevalence of sexual dysfunction is high in Iran and in the world; various studies indicated the prevalence of 20% to 40% for sexual dysfunction in Iran and 25% to 65% in Europe and the United States (18-21). Nonetheless, the types of sexual dysfunction caused by lower extremity trauma have not been well-defined.

2. Objectives

This study aimed at comparing male sexual dysfunction before and after lower extremity trauma.

3. Materials and Methods

3.1. Study Design and Settings

This research is a prospective cross sectional study. The study population was patients with lower extremity trauma who were admitted to the surgical unit of Kowsar hospital in Semnan from June 2014 to August 2015.

3.2. Participants

Available sampling method was used in the present study. The sample consisted of 150 individuals. The inclusion criteria were as follow: full consciousness, the ability to respond to the questionnaire, absence of pain at the time of completing the questionnaire, lack of risk of chronic diseases, no history of mental and sexual dysfunction, being married, and having just one lower extremity trauma. Exclusion criteria included infection of surgical area, bed sore, staying in the hospital for more than 2 weeks, bereaved patients, and the incidence of new trauma during the study.

3.3. Measures

3.3.1. A Demographic and Incident Questionnaire

The researcher made a 2-part questionnaire. In the demographic part, the questions inquired about age, number of children, duration of marriage, education level, and occupation. In the incident part, the questions inquired about the reason of trauma, the place of trauma, the time, and the reason of the incident.

3.3.2. International Index of Erectile function (IIEF)

Rosen et al. (1997) introduced IIEF. This questionnaire has 15 questions in Likert scale, which assigns 0 to 5 scores to each question, allocating a total of 75 scores. The subscales of the questionnaire are sexual desire (0 - 10 score), erectile function (0 - 30 score), satisfaction with sexual intercourse (0 - 15 score), orgasm (0 - 10 score), and overall satisfaction with sexual intercourse (0 - 10 score), respectively (22). Mehrban et al. approved the validity and reliability of the questionnaire for use in Iranian population in 2006 (23).

3.4. Data Analysis

The data were analyzed with SPSS software, edition 19. Then, demographic characteristics and the features of the incidents were presented using descriptive statistics (frequency, frequency percentage, mean, and standard deviation). Pearson correlation coefficients and Chi-square test were conducted to demonstrate the correlations between sexual dysfunction with demographic variables. Then, the changes of sexual dysfunctions of the participants were determined in the 3 stages of sampling using Friedman statistic test. The changes of sexual dysfunctions were investigated using the Wilcoxon test as follow: before the incident with 1 month after the incident; before the incident with 3 months after the incident; and 1 month before the incident with 3 months after the incident.

3.5. Ethical Considerations

The present study was approved by the ethics committee of Semnan University of Medical Sciences (No: 92.299762). Participation was voluntary and confidential.

3.6. Study Procedure

The researcher introduced himself to the patients and presented them with the approval of ethics committee of Semnan University of Medical Sciences. Then, he explained the study procedure to the patients. After the announcement of the patients' willingness to participate in the study, they filled the informed consent forms. Then, demographic and incident information questionnaire and international index of erectile function were given to the patients and they were asked to complete the demographic and incident information questionnaires based on their current condition and to fill the IIEF questionnaires based on their own sexual function status 1 month before the incident. Two stamped postal envelopes with 2 series of IIEF questionnaires were provided to the patient to complete in 1 and 3 months after the trauma; the patients were asked to mail the questionnaires to the address written at the back of the envelopes. The researcher also contacted the patients 1 and 3 months after the trauma to check the eligibility criteria and remind them to complete the questionnaire.

4. Results

Overall, 153 male patients participated in the study. The mean age of the participants was 38.55 ± 12.84 , and the average length of their marriage was 13.33 ± 12.5 years. The participants had the average number of 1.82 ± 1.56 children. Most participants (32%) had a diploma and others

held other academic degrees. In addition, most of the participants (52.9%) were workers and others were working in other occupations (Table 1). The results of the Pearson correlation coefficients revealed a statistically significant correlation between age, years of marriage, and number of children with sexual dysfunction score ($P < 0.05$). This means that older age, longer marriage duration, and more number of children, increased the severity of the disorder. The results of the Chi-square test also revealed a statistically significant correlation between the level of education and occupation with the severity of sexual dysfunction ($P < 0.05$), meaning that the intensity of sexual disorders was lower in those participants with higher-level education and better occupations.

The cause of the incident in 54.2% of the cases was traffic accidents. Most types of lower extremity trauma (31.4%) were related to the injuries of the left and right legs and the lowest type was related to hip trauma (9.2%). The event occurred in the morning in 43.1%, and in 3.3% it occurred at midnight; 23.5% of the lower extremity trauma were caused by car accidents (Table 1). The curves of the trends of the sexual dysfunction changes showed that all the subsets of the disorders had a sharp increase by the end of the first month. Nevertheless, their progress has fallen until the end of the third month after the accident. Furthermore, the curves indicated that the severity of any of the following classes of sexual dysfunction did not return to pretrauma period (Figure 1).




Figure 1. The Mean Scores of IIEF Subscales Changes Before and After Lower Extremity Trauma

The results of the Friedman test revealed a significant difference in 3 stages among all the subscale IIEF (Table 2). The results of the Wilcoxon test indicated that the changes of sexual dysfunctions were significantly different before the incident with 1 month and 3 months after the incident in all the IIEF subscales ($P = 0.000$). However, the changes of sexual dysfunction was not significantly different 1 month after the incident with 3 months after the incident in all cases, except the intercourse satisfaction ($P >$

Table 1. Demographic and Lower Limb Trauma Variables^a

Variables	Frequency
Age, y	38.55 ± 12.48
Child number, No	1.82 ± 1.56
Marriage history, y	13.33 ± 12.54
Educational status	
Primary education	31 (20.3)
Primary and junior high school	35 (22.9)
Diploma	49 (32)
Higher education	38 (24.9)
Type of job	
Government's employee	16 (10.4)
Proletarian	81 (52.9)
Student	1 (0.7)
Driver	3 (2)
Tradesman	9 (5.9)
Retired	17 (11.1)
Others job	26 (17)
Cause of trauma	
Traffic accidents	54.2
Non-traffic accidents	45.8
Trauma spot	
Hip	14 (9.2)
Right or left thigh	31 (20.3)
Right or left knee	27 (17.6)
Right or left tibia	48 (31.4)
Right or left ankle	18 (11.8)
Right or left toes	15 (9.8)
Time of incidents	
6 - 12	66 (43.1)
12 - 18	60 (39.2)
18 - 24	22 (14.4)
24 - 6	5 (3.3)
Mode of trauma	
Motorcycle	34 (22.2)
Car	36 (23.5)
Truck	8 (5.3)
Fall	26 (17.0)
Downfall of Things	31 (20.3)
Other classifiable	18 (11.7)

^aValues are expressed as No. (%) or mean ± SD.

0.05).

5. Discussion

Trauma, diseases, and injuries can cause sexual dysfunction in males and even in their spouses (24, 25). However, a few studies have been conducted in Iran about the effects of trauma, especially lower extremity trauma, on sexual health. Therefore, male sexual dysfunctions were assessed before and after lower limb trauma in this study.

Table 2. Sexual Dysfunction Before, One and Three Months After Lower Limb Trauma

Times Domain and Clinical Interpretation	Before the Incident	One Month Later	Three Months Later	P Value	
Erectile Function	Severe dysfunction	10 (6.5)	15 (9.8)	18 (11.8)	0.000 ^a
	Moderate dysfunction	0	2 (1.3)	15 (9.8)	
	Mild to moderate dysfunction	3 (2)	28 (18.3)	26 (17)	
	Mild dysfunction	36 (23.5)	83 (54.2)	49 (32)	
	No dysfunction	104 (68)	25 (16.3)	45 (29.4)	
Orgasmic Function	Severe dysfunction	10 (6.5)	15 (9.8)	14 (9.2)	0.000 ^a
	Moderate dysfunction	6 (3.9)	14 (9.2)	22 (14.4)	
	Mild to moderate dysfunction	11 (7.2)	38 (24.8)	33 (21.6)	
	Mild dysfunction	41 (26.8)	52 (34)	58 (37.9)	
	No dysfunction	85 (55.6)	34 (22.2)	26 (17)	
Sexual Desire	Severe dysfunction	11 (7.2)	14 (9.2)	10 (6.5)	0.000 ^a
	Moderate dysfunction	2 (1.3)	17 (11.1)	26 (17)	
	Mild to moderate dysfunction	12 (7.8)	57 (37.3)	38 (24.8)	
	Mild dysfunction	60 (39.2)	53 (34.6)	62 (40.5)	
	No dysfunction	68 (44.4)	12 (7.8)	17 (11.1)	
Intercourse Satisfaction	Severe dysfunction	10 (6.5)	15 (9.8)	11 (7.2)	0.000 ^a
	Moderate dysfunction	3 (2)	13 (8.5)	16 (10.5)	
	Mild to moderate dysfunction	13 (8.5)	63 (41.2)	45 (29.4)	
	Mild dysfunction	81 (52.9)	43 (28.1)	53 (34.6)	
	No dysfunction	46 (30.1)	19 (12.4)	28 (18.3)	
Overall Satisfaction	Severe dysfunction	9 (5.9)	15 (9.8)	13 (8.5)	0.000 ^a
	Moderate dysfunction	4 (2.6)	16 (10.5)	13 (8.5)	
	Mild to moderate dysfunction	10 (6.5)	46 (30.1)	53 (34.6)	
	Mild dysfunction	78 (51)	53 (34.6)	56 (36.6)	
	No dysfunction	52 (34)	23 (15)	18 (11.8)	

^aThe results of the Friedman test revealed a significant difference in 3 stages among all the subscale of IIEF.

The incidence of sexual dysfunction showed a significant increase in the participants after lower extremity trauma. Mild to severe sexual dysfunction were between 8.5% and 17.6% in all subcategories of the IIEF before the incidence of trauma. The incidence was almost the same as sexual dysfunctions in other studies. In Papagiannopoulos (2015), the incidence of erectile dysfunction has been reported to be 15% to 20% (26), and it was 21.1% in Foruzania study (27).

The results of the present study revealed a significant increase in the incidence of sexual dysfunction 1 month after the lower extremity trauma compared to before the event. In the first posttraumatic month, compared to before the event, the rate of moderate to severe erectile function disorder showed an increase from 8.5% to 29.4%, orgasmic function disorder from 17.6% to 67%, sexual desire disorder from 16.4% to 57.6%, intercourse satisfaction disorders from 17% to 59.5%, and overall satisfaction dysfunction from 14% to 50.4%. Other studies have reported the incidence of high sexual dysfunction after pelvic trauma and lower extremity trauma. For example, in a review of studies, Blaschko (2014) found that the incidence of erectile dysfunction was 25% to 45% after pelvic fractures in different studies (28). Also, Harvey-Kelly (2011) reported this quantity in males to be 35.9% in the other review studies (17). In another review study, Koraitim (2013) reported the incidence of sexual dysfunction to be 44% in patients with hip fractures (29). In the present study, the prevalence of disorders has increased to more than 50% in the other subgroups of the IIEF, except in erectile function disorders. The normal condition of erectile function in more than 70% of the patients showed that most participants in the study had not been physiologically banned from sexual activities and other reasons were involved in the incidence of these disorders. It seems that, in our study, the causes of sexual dysfunction in the first posttraumatic month were pains from trauma, sedative drugs, performing surgical procedures, being away from the family, fear and worry about the future, particularly the future job, disruption of order of the work and daily activities, the failure of the lower part of the body as an effective organ in sexual activity, and most importantly, not receiving the necessary training on how to perform sexual activities appropriate to the disease. The results revealed a significant increase in the incidence of

function was 25% to 45% after pelvic fractures in different studies (28). Also, Harvey-Kelly (2011) reported this quantity in males to be 35.9% in the other review studies (17). In another review study, Koraitim (2013) reported the incidence of sexual dysfunction to be 44% in patients with hip fractures (29). In the present study, the prevalence of disorders has increased to more than 50% in the other subgroups of the IIEF, except in erectile function disorders. The normal condition of erectile function in more than 70% of the patients showed that most participants in the study had not been physiologically banned from sexual activities and other reasons were involved in the incidence of these disorders. It seems that, in our study, the causes of sexual dysfunction in the first posttraumatic month were pains from trauma, sedative drugs, performing surgical procedures, being away from the family, fear and worry about the future, particularly the future job, disruption of order of the work and daily activities, the failure of the lower part of the body as an effective organ in sexual activity, and most importantly, not receiving the necessary training on how to perform sexual activities appropriate to the disease. The results revealed a significant increase in the incidence of

sexual dysfunctions in the third posttraumatic month of the lower limb trauma compared to before the event. The rate of mild to severe erectile function disorder increased from 8.5% to 38.6%, orgasmic function disorder from 17.6% to 45.2%, sexual desire disorder from 16.4% to 48.3%, intercourse satisfaction disorder from 17% to 47.1%, and overall satisfaction disorder from 14% to 51.6% in the third posttraumatic month compared to before it. However, the results showed a decline in the severity of sexual dysfunction in the third month compared to the first posttraumatic month. It seems that the reduction of stress arising from trauma and physical improvement of patients have been very effective in the reduction of an average of sexual dysfunctions severity in the third month; and the results of McCarthy (2003) and Soberg (2015) has confirmed it. In their studies, they found that the prevalence of mental disorders in individuals with lower extremity trauma was reduced in the months after the event (9, 11). The results also revealed no significant difference in the incidence of sexual dysfunction in all subcategories of IIEF, except intercourse satisfaction in the first posttraumatic month of lower limb extremity trauma compared to the third posttraumatic month. This finding is worrying considering the average age of 38.5% of the participants in the study. Because most of the participants were at the peak age of sexual functioning, and a sharp drop in their long-term sexual functions could cause adverse consequences such as mental disorders, disheartening relationship with their spouse, the desire to divorce, tendency to be unfaithful, tendency to drugs and alcohol, and other irritating substances of sexual activity (18).

It seems that one of the causes of such findings in the present study was the low education level of the participants in the study, as more than 75% of the participants had no college education. Perhaps, after the trauma, these individuals would not be able to learn about their needs through the study of literature and the electronic resources. They probably won't refer to the counselors. In addition, more than half of the participants were workers. Workers are usually active. They have regular patterns of sleeping, waking, working, and resting. Therefore, it was likely that they suffered from some depression, which had a negative impact on their sexual function.

As one of the first reports from Western Asia, especially the Persian Gulf region, the current study added something new to the literature. However, the study had a cross sectional design and lacked female participants, thus, conducting further studies on this topic is suggested.

5.1. Conclusions

Many males with lower extremity trauma are at risk of sexual dysfunction. These dysfunctions may continue for

months after the trauma. Lower extremity trauma occurs in young and middle age individuals and these ages are the climax of sexual activity of each person. Lack of attention to these dysfunctions can have adverse consequences for individuals, families, and the society. Thus, health care providers should evaluate, support, and if needed, treat sexual dysfunction in patients from the first days after the lower extremity trauma. Also, other members of the treatment team and family members of these patients should learn the required trainings to help these patients.

Acknowledgments

We would like to thank the nursing care research center and clinical research development unit of Kowsar educational, research and therapeutic center of Semnan University of Medical Sciences for providing facilities for this work and we also thank all the patients for their sincere cooperation.

Footnotes

Authors' Contribution: Abbasali Ebrahimiyan designed the study and developed the project, developed the data collection, and wrote the first draft of the manuscript. Hossein Davari and Maedeh Tourdeh undertook recruitment and data collection. All authors contributed to review and revision of the manuscript and all take responsibility for the final version.

Declaration of Interests: The authors declare that they have no competing interests.

Funding/Support: Semnan University of Medical Sciences has financially supported this study (grant No: 500).

References

1. Puvanachandra P, Hoe C, Ozkan T, Lajunen T. Burden of road traffic injuries in Turkey. *Traffic Inj Prev*. 2012;**13** Suppl 1:64-75. doi: [10.1080/15389588.2011.633135](https://doi.org/10.1080/15389588.2011.633135). [PubMed: [22414130](https://pubmed.ncbi.nlm.nih.gov/22414130/)].
2. Rasouli MR, Saadat S, Haddadi M, Gooya MM, Afsari M, Rahimi-Movaghar V. Epidemiology of injuries and poisonings in emergency departments in Iran. *Public Health*. 2011;**125**(10):727-33. doi: [10.1016/j.puhe.2011.07.006](https://doi.org/10.1016/j.puhe.2011.07.006). [PubMed: [21906762](https://pubmed.ncbi.nlm.nih.gov/21906762/)].
3. Gosselin RA, Spiegel DA, Coughlin R, Zirkle LG. Injuries: the neglected burden in developing countries. *Bull World Health Organ*. 2009;**87**(4):246-246a. [PubMed: [19551225](https://pubmed.ncbi.nlm.nih.gov/19551225/)].
4. Yadollahi M, Shamsedini N, Shayan L, Rezaianzadeh A, Bolandparvaz S. The Performance of Trauma Research Centers of Iran during the Past 10 Years; A Science Monitor Survey. *Bull Emerg Trauma*. 2014;**2**(1):44-51. [PubMed: [27162863](https://pubmed.ncbi.nlm.nih.gov/27162863/)].
5. American College of Surgeons Committee on Trauma Leadership. Annual Report of the National Trauma Data Bank. ;2014.
6. Morris S, Lenihan B, Duddy L, O'Sullivan M. Outcome after musculoskeletal trauma treated in a regional hospital. *J Trauma*. 2000;**49**(3):461-9. [PubMed: [11003324](https://pubmed.ncbi.nlm.nih.gov/11003324/)].

7. Ebrahimian A, Saberi-Zafarghandi MB, Alaei S, Tourdeh M, Davari H. The trend of changes in depression, anxiety and stress in men with lower limb trauma: a prospective study [In Persian]. *Koomesh*. 2016;**17**(4).
8. Stevelink SAM, Malcolm EM, Mason C, Jenkins S, Sundin J, Fear NT. The prevalence of mental health disorders in (ex-)military personnel with a physical impairment: a systematic review. *Occup Environ Med*. 2015;**72**(4):243-51. doi: [10.1136/oemed-2014-102207](https://doi.org/10.1136/oemed-2014-102207).
9. McCarthy ML, MacKenzie EJ, Edwin D, Bosse MJ, Castillo RC, Starr A, et al. Psychological distress associated with severe lower-limb injury. *J Bone Joint Surg Am*. 2003;**85-A**(9):1689-97. [PubMed: [12954826](https://pubmed.ncbi.nlm.nih.gov/12954826/)].
10. Soberg HL, Finset A, Roise O, Bautz-Holter E. The trajectory of physical and mental health from injury to 5 years after multiple trauma: a prospective, longitudinal cohort study. *Arch Phys Med Rehabil*. 2012;**93**(5):765-74. doi: [10.1016/j.apmr.2011.08.050](https://doi.org/10.1016/j.apmr.2011.08.050). [PubMed: [22459698](https://pubmed.ncbi.nlm.nih.gov/22459698/)].
11. Soberg HL, Bautz-Holter E, Finset A, Roise O, Andelic N. Physical and mental health 10 years after multiple trauma: A prospective cohort study. *J Trauma Acute Care Surg*. 2015;**78**(3):628-33. doi: [10.1097/TA.0000000000000541](https://doi.org/10.1097/TA.0000000000000541). [PubMed: [25710437](https://pubmed.ncbi.nlm.nih.gov/25710437/)].
12. Pouretamad HR, Naghavi HR, Malekafzali H, Noorbala AA, Davidian H, Ghanizadeh A, et al. Prevalence of mood disorders in Iran. *Iran J Psychiatry*. 2006;**1**(2):59-64.
13. Verschuren JE, Geertzen Jan H, Enzlin P, Dijkstra Pieter U, Rienk D. Individuals with lower limb amputation and their sexual functioning and sexual well-being. *Disabil Rehabil*. 2015;**37**(3):187-93.
14. Cobb AR, Tedeschi RG, Calhoun LG, Cann A. Correlates of posttraumatic growth in survivors of intimate partner violence. *J Trauma Stress*. 2006;**19**(6):895-903. doi: [10.1002/jts.20171](https://doi.org/10.1002/jts.20171). [PubMed: [17195973](https://pubmed.ncbi.nlm.nih.gov/17195973/)].
15. Belen RA, Garcia Salord J, Fiallos L. [Prevalence of male sexual dysfunctions]. *Rev Fac Cien Med Univ Nac Cordoba*. 2000;**57**(2):239-47. [PubMed: [12934239](https://pubmed.ncbi.nlm.nih.gov/12934239/)].
16. Shulman BS, Taormina DP, Patsalos-Fox B, Davidovitch RI, Karia RJ, Egol KA. Sexual Function Is Impaired After Common Orthopaedic Nonpelvic Trauma. *J Orthopaed Trauma*. 2015;**29**(12):e487-92. doi: [10.1097/bot.0000000000000397](https://doi.org/10.1097/bot.0000000000000397).
17. Harvey-Kelly KF, Kanakaris NK, Obakponwwe O, West RM, Giannoudis PV. Quality of life and sexual function after traumatic pelvic fracture. *J Orthop Trauma*. 2014;**28**(1):28-35. doi: [10.1097/BOT.0b013e31828fc063](https://doi.org/10.1097/BOT.0b013e31828fc063). [PubMed: [23481925](https://pubmed.ncbi.nlm.nih.gov/23481925/)].
18. Ebrahimian A, Heydari M, Saberi Zafar Ghandi MB, Delavari S. Comparing sexual dysfunctions in men before and during their wives' pregnancy. *Iran J Obstetr Gynecol Infertil*. 2013;**19**:25.
19. Ebrahimian A, Heydari M, Zafarghandi S. Comparison of female sexual dysfunctions before and during pregnancy. *Iran J Obstetr Gynecol Infertil*. 2010:30-6.
20. Sandfort TG, Ehrhardt AA. Sexual health: a useful public health paradigm or a moral imperative?. *Arch Sex Behav*. 2004;**33**(3):181-7. doi: [10.1023/B:ASEB.0000026618.16408.e0](https://doi.org/10.1023/B:ASEB.0000026618.16408.e0). [PubMed: [15129037](https://pubmed.ncbi.nlm.nih.gov/15129037/)].
21. Heiman JR. Sexual dysfunction: overview of prevalence, etiological factors, and treatments. *J Sex Res*. 2002;**39**(1):73-8. doi: [10.1080/00224490209552124](https://doi.org/10.1080/00224490209552124). [PubMed: [12476261](https://pubmed.ncbi.nlm.nih.gov/12476261/)].
22. Rosen RC, Riley A, Wagner G, Osterloh IH, Kirkpatrick J, Mishra A. The international index of erectile function (IIEF): a multidimensional scale for assessment of erectile dysfunction. *Urology*. 1997;**49**(6):822-30. [PubMed: [9187685](https://pubmed.ncbi.nlm.nih.gov/9187685/)].
23. Mehraban D, Shabaninia SH, Naderi GHH, Esfahani F. Farsi international index of erectile dysfunction and doppler ultrasonography in the evaluation of male impotence [In Persian]. *Iran J Surg*. 2006;**14**(1):9-15.
24. Sander AM, Maestas KL, Pappadis MR, Sherer M, Hammond FM, Hanks R, et al. Sexual functioning 1 year after traumatic brain injury: findings from a prospective traumatic brain injury model systems collaborative study. *Arch Phys Med Rehabil*. 2012;**93**(8):1331-7. doi: [10.1016/j.apmr.2012.03.037](https://doi.org/10.1016/j.apmr.2012.03.037). [PubMed: [22840831](https://pubmed.ncbi.nlm.nih.gov/22840831/)].
25. Nasir AR, Zehri AA, Abbas F, Ather MH. The correlation between international prostate symptoms score and sexual health inventory in men with lower urinary tract symptoms. *Int Urol Nephrol*. 2011;**43**(3):625-9. doi: [10.1007/s11255-010-9888-3](https://doi.org/10.1007/s11255-010-9888-3). [PubMed: [21229391](https://pubmed.ncbi.nlm.nih.gov/21229391/)].
26. Papagiannopoulos D, Khare N, Nehra A. Evaluation of young men with organic erectile dysfunction. *Asian J Androl*. 2015;**17**(1):11-6. doi: [10.4103/1008-682X.139253](https://doi.org/10.4103/1008-682X.139253). [PubMed: [25370205](https://pubmed.ncbi.nlm.nih.gov/25370205/)].
27. Foruzan-Nia SK, Abdollahi MH, Hekmatimoghaddam SH, Namayandeh SM, Mortazavi MH. Incidence of sexual dysfunction in men after cardiac surgery in Afshar hospital, Yazd. *Int J Reprod Bio Med*. 2012;**9**(2):85-94.
28. Blaschko SD, Sanford MT, Schlomer BJ, Alwaal A, Yang G, Villalta JD, et al. The incidence of erectile dysfunction after pelvic fracture urethral injury: A systematic review and meta-analysis. *Arab J Urol*. 2015;**13**(1):68-74. doi: [10.1016/j.aju.2014.09.004](https://doi.org/10.1016/j.aju.2014.09.004). [PubMed: [26019983](https://pubmed.ncbi.nlm.nih.gov/26019983/)].
29. Koraitim MM. Predictors of erectile dysfunction post pelvic fracture urethral injuries: a multivariate analysis. *Urology*. 2013;**81**(5):1081-5. doi: [10.1016/j.urology.2012.12.040](https://doi.org/10.1016/j.urology.2012.12.040). [PubMed: [23465164](https://pubmed.ncbi.nlm.nih.gov/23465164/)].