



Intensity and Frequency of Moral Distress among Iranian Nurses: A Systematic Review and Meta-Analysis

Mohammad Hossein YektaKooshali,¹ Mohammad Esmailpour-Bandboni,^{2,*} and Mohsen Andacheh¹

¹Student Research Committee, School of Nursing, Midwifery and Paramedicine, Guilan University of Medical Sciences, Rasht, Iran

²School of Nursing, Midwifery and Paramedicine, Guilan University of Medical Sciences, Rasht, Iran

*Corresponding author: Mohammad Esmailpour-Bandboni, School of Nursing, Midwifery and Paramedicine, Guilan University of Medical Sciences, Rasht, Iran. Tel: +98-1342333553, Fax: +98-1342565051, E-mail: esmaeilmmm@yahoo.com

Received 2017 January 20; Revised 2017 March 25; Accepted 2017 July 10.

Abstract

Context: Moral distress is a common phenomenon in the nursing profession. This study was aimed to determine the mean intensity and frequency of moral distress among Iranian Nurses.

Evidence Acquisition: This systematic review and meta-analysis was based on PRISMA guideline and searching in National databases as SID, Magiran, Irandoc, Noormag, IranMedex, and Medlib as well as International databases as Madlib and Web of Sciences, Scopus, PubMed, Direct-Science, Cochrane, Embase, Springer, Online Library Wiley, Ebsco, CEBM, CINHAL, SAGE, and Google Scholar were published by 2016. Researched keywords were in Persian and their English equivalents in accordance with their Mesh: "Moral", "Morale", "Distress", "Iran", "Frequency", "Intensity", "Ethics", "Nurse", and all possible combinations. The obtained documents were analyzed based on appropriate checklist through a random effects model by using Review Manager Software ver: 5.3.5.

Results: According to 12 (3.08%) selected documents out of 389, 2655 Iranian nurses with a mean age of 32.3 ± 15.265 years entered to meta-analysis. Mean intensity of moral distress was calculated 2.2329 [CI95%: 1.7651, 2.5230] out of 4, 2.9566 [CI95%: 2.4362, 3.4769] out of 5, and 3.4328 [CI95%: 3.3200, 3.5455] out of 6. Furthermore, its frequency was calculated 1.9237 [CI95%: 1.5533, 2.2941] out of 4, 3.0278 [CI95%: 2.4533, 3.6024] out of 5, and 2.6261 [CI95%: 2.5210, 2.7312] out of 6. Which the range of both has been reported in moderate to intense

Conclusions: According to the results, the intensity and frequency of moral distress have been reported in the range of moderate to intense. Due to the most harmful effects of moral distress in nursing, management and prevention protocols of moral distress should be developed and done by nurses.

Keywords: Ethics, Frequency, Intensity, Iran, Morale, Morals, Moral Distress, Nurse

1. Context

Moral distress is a common phenomenon in the nursing profession (1). It is the emotional and mental pain and discomfort, which is morally inevitable while having the required knowledge and ability to make moral judgments, due to actual or mental limitations (2). Today, since nurses are correlated with many moral issues in their working shifts and patient care is always associated with ethics and ethical considerations, moral distress is of great importance (3, 4).

Insult, mistrust, and conflicting demands of the patient and his family from the therapy team (5, 6), shortage of manpower in clinical environments, negligence and medical errors, lack of time, poor communication of doctors and nurses (7), opposition to superiors, medical limitations, institutional policies and personal or ideological

ethical considerations, and incompetence of some medical personnel can be the reasons for the emergence of moral distress (8, 9). It causes a negative impact on personal and professional lives of nurses (10), physical and psychological stress, low efficiency, quality of care, and increase in the cost of nursing care (11-15).

As Corley, a pioneer in the field of nurses' moral distress, stated moral distress can cause nurses to experience conflict in caring and avoid facing with the patient and providing high-quality care. It involves absence from work and leaving the profession that leads to a defect in improvement and increasing the duration of hospitalization and it also affects other medical professionals' services. It can also act as a deterrent and destructive factor for nurses to provide caring (11, 16-18). Effects of moral distress in nurses can lead to wrath, depression, disability and frustration, anger and physiological or psychological problems

(7).

Moral distress is examined by two intensity and frequency dimensions: intensity means the perceived amount of moral distress but the frequency is the number of times nurses face with it in a limited interval. As Hamric Said "questionnaire contains 21 question and each of the 21 items scored by participants in terms of the how often the situation arises (frequency) and how disturbing the situation is when it arises ("intensity)". The scale ranges never to very frequently, and none to great extent for frequency and intensity, respectively (11, 19-21).

1.1. Objectives

Increasing sample size due to the increasing number of relevant studies, reducing difference in existing parameters and confidence interval are the main goals of meta-analysis studies. In fact, these studies are a vital link between research studies and decision making at the bedside (22). Due to the controversy in reporting the moral distress there is a lack of access to the global community to this factor in Iranian nurses, on the other hand, review of several studies cannot lead to final conclusions for policy-making to planning the proper preventive management at the national level. Therefore, a systematic review of all documents and their combination through meta-analysis can create a complete viewpoint (22) of the dimensions of moral distress in Iranian nurses. This first international study was done aiming to investigate the "Intensity and Frequency of Moral Distress in Iranian Nurses".

2. Evidence Acquisition

This is the first systematic and meta-analysis review study aiming to evaluate the intensity and frequency of moral distress in Iranian nurses by 2016. The present meta-analysis study was conducted in accordance with the PRISMA guideline (23). To avoid publication bias, the entire processes of research including search, selection of studies, qualitative evaluation of studies, and data extracting were independently done by 2 researchers. Any additional encounters was assessed by a third researcher.

2.1. Study Selection

2.1.1. Inclusion and Exclusion Criteria

The main criteria for inclusion in the study were an evaluation of the frequency and intensity of moral distress in Iranian nurses. The exclusion criteria included: 1. Non-Iranian sample, 2. non-nurse sample, 3. poor information like failing to report just one of the items "intensity and frequency", 4. unrelated to the research subject, 5. non-random sample, 6. letters to the editor, and 7. duplicate articles.

2.1.2. Search Strategy and Study Selection

Results of this study are based on all papers published in national and international journals, dissertations, and reference websites. Related English and Persian literature were reviewed in national databases like SID, Magiran, Irandoc, IranMedex, and international banks such as Madlib and Web of Sciences, Scopus, PubMed, Direct-Science, Cochrane, Embase, Springer, Online Library Wiley, Ebsco, CEBM, CINHALL, SAGE, and Google Scholar search engine. Searching articles was done using Persian keywords and their English equivalents in accordance with their Mesh: "Moral", "Morale", "Distress", "Iran", "Frequency", "intensity", "Ethics", "Nurse", and all possible combinations of words using "Boolean operators" in combination for English databases. The manual search to find more articles was also done through reviewing references of identified articles.

What is noticeable in searching databases, is "High Sensitive Searching". Searching was performed by a researcher and master familiar with searching databases. To avoid bias, the search was separately done by 2 researchers ("M.H. YK" and "M. A") any encounters was assessed by a third researcher. As soon as the search ended, titles of the collected articles were entered into EndNote™, resource management software, to find similar articles. Manual searching was also done through reviewing the reference list of relevant articles.

2.1.3. Quality Assessment

In the next step, using STROBE (24) checklist, which has 22 sections, researchers evaluated the selected articles from methodological aspects, including sampling method, measurement of variables, statistical analysis, and study objectives, independently and any encounters were assessed by a third researcher. Articles, which obtained a minimum score of 16 based on the checklist, were selected for meta-analysis (Figure 1).

2.2. Data Extraction

At first, a checklist was designed based on objectives and by studying other available resources. The designed checklist included items such as authors, year, place, sample size, questionnaire, score of scale, mean intensity (\pm SD), mean frequency (\pm SD), mean age (\pm SD), and Job Experience, which was extracted independently by 2 researchers (Table 1). All documents were assessed and data extracted by 2 researchers independently and all extracted data were rechecked after they were finished by the third researcher who is an expert in this field (M.E-B).




Figure 1. A flow Diagram Following the PRISMA Template (Depicted by RevMan Software)

Table 1. Specifications Studies Entered Into Meta-Analysis (Sorted by Year)

Authors	Year	Place	Sample Size	Questionnaire	Score Scale	Mean Intensity	Mean Frequency	Mean Age	Job Experience (Lowest-Highest)
Borhani et al. (25)	2015	Tehran	153	Corely995	4	1.8667 ± 0.6950	1.9417 ± 0.6854	32.9 ± 6.52	9.23 ± 6.43(1-35)
Shafei et al. (15)	2015	Borazjan	91	Corely995	4	2.2329 ± 0.4742	1.9829 ± 0.5921	33.91 ± 7.08	10.5 ± 6.78(1-26)
Borhani et al. (26)	2015	Kerman	300	Corely2001	5	3.54 ± 0.3	3.11 ± 0.6	32 ± 3.1	14 ± 4(2-25)
Mohammadi et al. (5)	2014	Kerman	260	Corely995	5	3.5 ± 0.8	3.9 ± 0.55	37 ± 2.46	14(2-25)
Mohammadi et al. (27)	2014	Kerman	330	Corely995	5	3.5 ± 0.9	3.7 ± 1.2	-	16(1-28)
Borhani et al. (28)	2014	Birjand	220	Corely2001	5	2.25 ± 0.6	2.11 ± 0.56	31.12 ± 5.13	6.54 ± 4.4(1-24)
Borhani et al. (14)	2013	Kerman	110	Corely995	6	3.5 ± 1.2	2.7 ± 1.1	-	(1-27)
AbbasZadeh et al. (29)	2013	Sirjan	184	Corely2002	4	1.92 ± 0.77	1.5 ± 0.7	32.7 ± 5.49	10
AbbasZadeh et al. (30)	2013	Birjand	220	Corely2001	5	2.25 ± 0.6	2.11 ± 0.56	-	(1-24)
Mohammadi-Nafchi et al. (31)	2013	Kerman	300	Corely2001	5	2.7 ± 0.57	3.24 ± 0.43	28 ± -	16(1-28)
AbbasZadeh et al. (32)	2012	Shiraz	117	Corely2005	4	2.5475 ± 0.0558	2.2642 ± 0.0475	31.38 ± 5.65	5.59 ± 4.64

2.3. Statistical Analysis

In any study, after taking into account the intensity and frequency of moral distress as a binomial probability distribution, its variance was calculated through bino-

mial distribution and Cochran test (Q) and I2 index were used to evaluate the heterogeneity of the studies. Due to the heterogeneity of studies, random effects model was used to combine the results of both studies. Data was ana-

lyzed using the Meta-analysis specialized software of “The Cochrane Collaboration Co.”, called Review Manager Ver. 5.3.5 and comprehensive Meta-Analysis V.2 a significance level less than 0.05 was considered.

3. Results

In this systematic study, based on performed searches, 389 article were identified and after review and final assessment in accordance with checklist, 12 (3.08%) titles were included in the list with a total of 2655 nurses with a mean age of 32.3 ± 15.265 and work experience (1 - 25 years) (Table 1 and Figure 2). The results of the systematic review and meta-analysis, are divided into 2 parts: Intensity and frequency.

Results of the meta-analysis of the mean intensity of moral distress in Iranian nurses are 2.2329 [CI95%: 1.7651, 2.5230] out of 4, 2.9566 [CI95%: 2.4362, 3.4769] out of 5, and 3.4328 [CI95%: 3.3200, 3.5455] out of 6 (Figure 3).

Results of the meta-analysis of mean frequency of moral distress in Iranian nurses are 1.9237 [CI95%: 1.5533, 2.2941] out of 4, 3.0278 [CI95%: 2.4533, 3.6024] out of 5, and 2.6261 [CI95%: 2.5210, 2.7312] out of 6 (Figure 4).

The heterogeneity rate in the study was 66.33%, which is in the midrange (I2 index less than 25% is low heterogeneity, between 25% and 75% is midrange, and more than 75% is high heterogeneity).

Also, the probability of publication bias was not statistically significant by using Begg and Egger tests and was determined as $P = 0.6457$ and $P = 0.8632$, respectively.

4. Discussion

This systematic review and meta-analysis study was carried on 12 academic papers, and has provided basic and enough information regarding morale distress among Iranian nurses. This study revealed the frequency and intensity of morale distress, which can be used by policy makers and nursing managers.

Results of this study can give a more accurate understanding of moral distress in the clinical environments to researchers and nurses. What is understood from Iranian studies is that, nurses as the key individuals in the health system the have most direct contact with patients, thus, 50% - 80% of health care is done by nurses (33). There is a consensus on the concept of moral distress by Andrew Jameton (1984) that the nurse knows what is correct, however, the organizational restrictions makes it impossible to do the right thing (26, 32, 34). In Iranian studies, no tool is generated to assess the distress specific to Iranian society and culture, and studies are mainly conducted based on

Jameson tools 1992 (35), Corley edited the version of 1995 (19, 25), 2001 (21, 26), 2002 (20, 29), and 2005 (11, 32). These tools were adjusted by Iranian researchers for Iranian culture in years (1, 3, 5, 11, 14, 15, 25-32, 34-41). Studies in Iran have manipulated the number of Likert scales and in Likert 3, 4, 5, 6, and 7, they have reported that this makes it difficult to compare the results (Table 1). According to the aim of this study, intensity and frequency of moral distress have been reported in the range of moderate to intense.

Iranian studies suggest that most nurses are in moral stressful situations and this can jeopardize their health or have a negative impact on their professional performance (42). Iranian studies expressed the most common side effects of moral distress as qualm, pain, and discomfort, lack of job satisfaction, turnover (26), and it seems that each of the side effects is worse than the moral distress.

Iranian studies suggest that moral distress is a serious challenge in nursing and requires special attention (18, 34). Nafechi (31) and Abbaszadeh (32) also expressed distress was high in nursing staff and moral distress reduction strategies should be planned and implemented. It is always stressed that organizational and nursing managers should develop mechanisms to reduce moral distress and support nurses (43, 44). However, since care in Iran is not based on evidence, the results of these studies have never been used. Everyone believes that further studies should be performed in this regard, and basically it is unclear when and how the nursing community will deal with this moral issue in practice. These structures must be developed properly, impartially and, naturally by nurses.

Trisha states that more studies need to be done by all members of the treatment team on why we cannot apply more effective interventions against the phenomenon of moral distress (44). Before studying the definitions of moral distress, we should focus more on the issue of what we should do to reduce the effects of moral distress, especially in nurses.

The heterogeneity rate (I2) has been calculated to 66%. Observed differences are due to different sampling. In addition, differences in measured parameters in different populations can be the reasons of heterogeneity in this study (45-47).

Overall, to avoid personal, organizational, and professional effects of moral distress in the clinical setting, a standard tool based on Iranian health care culture should be developed. The frequency and intensity of moral distress and related factors on nurses across the country should be evaluated. In the next step, the solutions of dealing with each of these factors should be provided according to Iranian health care background. Results of this study, our knowledge, and experiences show that because of terrible disruptive effects of moral distress on nurses, man-



Figure 2. Iranian Selected Studies on Intensity and Frequency of Moral Distress Among Nurses Based on Continent for Random Effects Meta-Analyses




Figure 3. Mean Intensity of Moral Distress from Score 4, 5 and 6 Among Iranian Nurses (Sorted by Year)

agement, and preventative strategies should only be developed and run by nurses.

Limitation: The number of men was very low so it was not possible to express the distressing rate based on gender. In addition, lack of focus on wards made it impossi-

ble to detect the rate of distress in each ward and comparison between the ward and other wards. It is recommended to conduct further studies to assess distressing rate based on gender and on wards (e.g. Emergency, Internal, ICU/CCU/NICU).




Figure 4. Mean Frequency of Moral Distress from Score 4, 5 and 6 Among Iranian Nurses (Sorted by Year)

Acknowledgments

Hereby, we would like to thank Guilan University of Medical Sciences and student research committee with IR.GUMS.REC.1395.361 (ethical committee code) for supporting us in publishing the first systematic review and meta-analysis of moral distress among Iranian nurses. Furthermore, we would like to thank Dr. Aboozar Ramezani for his scientific advising thorough searching in different databases.

Footnotes

Authors' Contribution: Mohammad Hossein YektaKooshali conceived and designed the study proposal, drafted the manuscript, performed the searching, data extraction according to the check list, and Meta-analysis. Mohammad Esmaeilpour-Bandboni performed scientific advising, searching, data extraction (as an expert), and Proof reading. Mohsen Andacheh performed searching, data extraction and English editing. All authors read and approved the final manuscript.

Declaration of Interests: None declared.

Funding/ Support: This study was founded and supported by Guilan University of Medical Sciences and Stu-

dent Research Committee with IR.GUMS.REC.1395.361 (ethical committee) code.

References

- Shafipour V, Esmaeili R, Heidari MR, Aghaei N, Saadatmehr SR, Sanagoo A. Investigating the level of moral distress and its related factors among nurses in mazandaran burn center. *J Mazandaran Univ Med Sci*. 2015;25(126):58-67. Persian.
- Nathaniel AK. Moral reckoning in nursing. *West J Nurs Res*. 2006;28(4):419-38. discussion 439-48. doi: [10.1177/0193945905284727](https://doi.org/10.1177/0193945905284727). [PubMed: [16672631](https://pubmed.ncbi.nlm.nih.gov/16672631/)].
- Beikmoradi A, Rabiee S, Khatiban M, Cheraghi MA. Nurses distress in intensive care unit: a survey in teaching hospitals. *Iran J Med Ethics History Med*. 2012. Persian.
- Range LM, Rotherham AL. Moral distress among nursing and non-nursing students. *Nurs Ethics*. 2010;17(2):225-32. doi: [10.1177/0969733009352071](https://doi.org/10.1177/0969733009352071). [PubMed: [20185446](https://pubmed.ncbi.nlm.nih.gov/20185446/)].
- Mohammadi S, Borhani F, Roshanzadeh L, Roshanzadeh M. Moral distress and compassion fatigue in patient care: a correlational study on nurses. *Iran J Med Ethics History Med*. 2014;7(2):69-79. Persian.
- Janvier A, Nadeau S, Deschenes M, Couture E, Barrington KJ. Moral distress in the neonatal intensive care unit: caregiver's experience. *J Perinatol*. 2007;27(4):203-8. doi: [10.1038/sj.jp.7211658](https://doi.org/10.1038/sj.jp.7211658). [PubMed: [17304208](https://pubmed.ncbi.nlm.nih.gov/17304208/)].
- Jones A. Moral distress among nurses. *J Emerg Nurs*. 2014;40(1):4. doi: [10.1016/j.jen.2013.03.022](https://doi.org/10.1016/j.jen.2013.03.022). [PubMed: [24384480](https://pubmed.ncbi.nlm.nih.gov/24384480/)].
- Cronqvist A, Theorell T, Burns T, Lutzen K. Caring about-caring for: moral obligations and work responsibilities in intensive care nursing. *Nurs Ethics*. 2004;11(1):63-76. doi: [10.1191/0969733004ne6670a](https://doi.org/10.1191/0969733004ne6670a). [PubMed: [14763651](https://pubmed.ncbi.nlm.nih.gov/14763651/)].

9. Harrowing JN, Mill J. Moral distress among Ugandan nurses providing HIV care: a critical ethnography. *Int J Nurs Stud*. 2010;**47**(6):723-31. doi: [10.1016/j.ijnurstu.2009.11.010](https://doi.org/10.1016/j.ijnurstu.2009.11.010). [PubMed: [20004395](https://pubmed.ncbi.nlm.nih.gov/20004395/)].
10. Hart TJ. *Moral distress in a non-acute continuing care setting: The experience of registered nurses. [dissertation]*. Canada: Queen's Univ., Kingston, Ont; 2009.
11. Corley MC, Minick P, Elswick RK, Jacobs M. Nurse moral distress and ethical work environment. *Nurs Ethics*. 2005;**12**(4):381-90. doi: [10.1191/0969733005ne8090a](https://doi.org/10.1191/0969733005ne8090a). [PubMed: [16045246](https://pubmed.ncbi.nlm.nih.gov/16045246/)].
12. Redman BK, Fry ST. Nurses' ethical conflicts: what is really known about them? *Nurs Ethics*. 2000;**7**(4):360-6. doi: [10.1177/096973300000700409](https://doi.org/10.1177/096973300000700409). [PubMed: [11221412](https://pubmed.ncbi.nlm.nih.gov/11221412/)].
13. Russell AC. Moral distress in neuroscience nursing: an evolutionary concept analysis. *J Neurosci Nurs*. 2012;**44**(1):15-24. quiz 25-6. doi: [10.1097/JNN.0b013e31823ae4cb](https://doi.org/10.1097/JNN.0b013e31823ae4cb). [PubMed: [22210300](https://pubmed.ncbi.nlm.nih.gov/22210300/)].
14. Borhani F, Abbaszadeh A, Arab M, Faramarzpour M. Moral Distress in Nurses. *Med Ethics J*. 2013;**7**(26):35-57. Persian.
15. Shafiea A, Ravanipour M, Yazdankhah-Fard MR, Mirzaei K. [The relationship between moral distress and burnout among nurses of Shahid Ganji Hospital, Borazjan in 1394]. *nursing of the vulnerable journal*. **2**(5). 2016. p. 15-26. Persian.
16. Jameton A. Dilemmas of moral distress: moral responsibility and nursing practice. *AWHONN's Clin Issues Perinatal Women's Health Nurs*. 1993;**4**(4):542-51. [PubMed: [8220368](https://pubmed.ncbi.nlm.nih.gov/8220368/)].
17. Hamric AB, Blackhall LJ. Nurse-physician perspectives on the care of dying patients in intensive care units: collaboration, moral distress, and ethical climate. *Crit Care Med*. 2007;**35**(2):422-9. doi: [10.1097/01.CCM.0000254722.50608.2D](https://doi.org/10.1097/01.CCM.0000254722.50608.2D). [PubMed: [17205001](https://pubmed.ncbi.nlm.nih.gov/17205001/)].
18. Elpern EH, Covert B, Kleinpell R. Moral distress of staff nurses in a medical intensive care unit. *Am J Crit Care*. 2005;**14**(6):523-30. [PubMed: [16249589](https://pubmed.ncbi.nlm.nih.gov/16249589/)].
19. Corley MC. Moral distress of critical care nurses. *Am J Crit Care*. 1995;**4**(4):280-5. [PubMed: [7663591](https://pubmed.ncbi.nlm.nih.gov/7663591/)].
20. Corley MC. Nurse moral distress: a proposed theory and research agenda. *Nurs Ethics*. 2002;**9**(6):636-50. doi: [10.1191/0969733002ne5570a](https://doi.org/10.1191/0969733002ne5570a). [PubMed: [12450000](https://pubmed.ncbi.nlm.nih.gov/12450000/)].
21. Corley MC, Elswick RK, Gorman M, Clor T. Development and evaluation of a moral distress scale. *J Adv Nurs*. 2001;**33**(2):250-6. [PubMed: [11168709](https://pubmed.ncbi.nlm.nih.gov/11168709/)].
22. YektaKooshali MH, Esmailpour-Bandboni M, Sharemi SH, Alipour Z. Survival Rate and Average Age of the Patients with Breast Cancer in Iran: Systematic Review and Meta-Analysis. *J Babol Univ Med Sci*. 2016;**18**(8):29-40. Persian.
23. Moher D, Liberati A, Tetzlaff J, Altman DG, Prisma Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol*. 2009;**62**(10):1006-12. doi: [10.1016/j.jclinepi.2009.06.005](https://doi.org/10.1016/j.jclinepi.2009.06.005). [PubMed: [19631508](https://pubmed.ncbi.nlm.nih.gov/19631508/)].
24. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;**370**(9596):1453-7. doi: [10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X).
25. Borhani F, Abbaszadeh A, Mohamadi E, Ghasemi E, Hoseinabad-Farahani MJ. Moral sensitivity and moral distress in Iranian critical care nurses. *Nurs Ethics*. 2017;**24**(4):474-82. doi: [10.1177/0969733015604700](https://doi.org/10.1177/0969733015604700).
26. Borhani F, Mohammadi S, Roshanzadeh M. Moral distress and perception of futile care in intensive care nurses. *J Med Ethics Hist Med*. 2015;**8**:2. [PubMed: [26839676](https://pubmed.ncbi.nlm.nih.gov/26839676/)].
27. Mohammadi S, Talebi F, Borhani F, Roshanzadeh M. Moral distress and attitude to euthanasia: A correlation study in nurses. *Med Ethics J*. 2014;**8**(28). Persian.
28. Borhani F, Abbaszadeh A, Nakhaee N, Roshanzadeh M. The relationship between moral distress, professional stress, and intent to stay in the nursing profession. *J Med Ethics Hist Med*. 2014;**7**:3. [PubMed: [25512824](https://pubmed.ncbi.nlm.nih.gov/25512824/)].
29. Zadeh AA, Borhani F, Ameri GF, Dolat-Abadi SN. Relation between moral distress and Work engagement in nurse's cross-sectional-correlation study. *Med Ethics J*. 2015;**7**(24):33-53. Persian.
30. Abbaszadeh A, Nakhaei N, Borhani F, Roshanzadeh M. The relationship between moral distress and retention in nurses in Birjand teaching hospitals. *Iran J Med Ethics History Med*. 2013;**6**(2):57-66. Persian.
31. Nafchi SM, Borhani F, Roshanzadeh M. Relationship between moral distress and environmental empowerment for nurses in education hospitals in Kerman. *Med Ethics J*. 2015;**8**(26):91-113. Persian.
32. Abbaszadeh A, Borhani F, Sharif F, Toghray FAS. The relationship between nurse's moral distress and end of life decision making in ICU. *Med Ethics J*. 2016;**7**(22):117-33. Persian.
33. Heidari MR, Norouzadeh R. Supporting families of dying patients in the intensive care units. *Holist Nurs Pract*. 2014;**28**(5):316-22. doi: [10.1097/HNP.0000000000000045](https://doi.org/10.1097/HNP.0000000000000045). [PubMed: [25099985](https://pubmed.ncbi.nlm.nih.gov/25099985/)].
34. Vaziri MH, Merghati-Khoei E, Tabatabaei S. Moral Distress among Iranian Nurses. *Iran J Psychiatry*. 2015;**10**(1):32-6. [PubMed: [26005478](https://pubmed.ncbi.nlm.nih.gov/26005478/)].
35. Keighobadi F, Sadeghi H, Keighobadi F, Tabaraei Y. The relationship between moral distress and emotional exhaustion in nurses. *Iran J Med Ethics History Med*. 2014;**7**(3):36-47. Persian.
36. Fazljoor E, Borhani F, Abbaszadeh A, Razban F. The relationship between nurses' perceptions of moral distress and the ethical climate in Shahid Sadoughi University of Medical Sciences of Yazd. *Iran J Med Ethics History Med*. 2014;**7**(2):80-90. Persian.
37. Ameri M, Safavibayatneed Z, Kavousi A. Moral distress of oncology nurses and morally distressing situations in oncology units. *Aust J Adv Nurs*. 2016;**33**(3):6.
38. Shoorideh FA, Ashktorab T, Yaghmaei F, Alavi Majd H. Relationship between ICU nurses' moral distress with burnout and anticipated turnover. *Nurs Ethics*. 2015;**22**(1):64-76. doi: [10.1177/0969733014534874](https://doi.org/10.1177/0969733014534874). [PubMed: [24948793](https://pubmed.ncbi.nlm.nih.gov/24948793/)].
39. Atashzadeh Shorideh F, Ashktorab T, Yaghmaei F. Iranian intensive care unit nurses' moral distress: a content analysis. *Nurs Ethics*. 2012;**19**(4):464-78. doi: [10.1177/0969733012437988](https://doi.org/10.1177/0969733012437988). [PubMed: [22691602](https://pubmed.ncbi.nlm.nih.gov/22691602/)].
40. Dargahi H, Shaham G. Life Change Units (LCU) rating as stressors in Iranian hospitals' nurses. *Acta Med Iran*. 2012;**50**(2):138-46. [PubMed: [22359085](https://pubmed.ncbi.nlm.nih.gov/22359085/)].
41. Soleimani MA, Sharif SP, Yaghoobzadeh A, Sheikhi MR, Panarello B, Win MT. Spiritual well-being and moral distress among Iranian nurses. *Nurs Ethics*. 2016. doi: [10.1177/0969733016650993](https://doi.org/10.1177/0969733016650993). [PubMed: [27312198](https://pubmed.ncbi.nlm.nih.gov/27312198/)].
42. Abbasi M, Hajatmand F, Khaghanizadeh M, Gashtroudkhani AA. Moral distress in nurses employed in selected hospitals of Shahid Beheshti University of Medical Sciences. *Med Ethics J*. 2015;**9**(32):121-40. Persian.
43. Rathert C, May DR, Chung HS. Nurse moral distress: A survey identifying predictors and potential interventions. *Int J Nurs Stud*. 2016;**53**:39-49. doi: [10.1016/j.ijnurstu.2015.10.007](https://doi.org/10.1016/j.ijnurstu.2015.10.007). [PubMed: [26515025](https://pubmed.ncbi.nlm.nih.gov/26515025/)].
44. Prentice T, Janvier A, Gillam L, Davis PG. Moral distress within neonatal and paediatric intensive care units: a systematic review. *Arch Dis Childhood*. 2016;**101**(8):701-8. doi: [10.1136/archdischild-2015-309410](https://doi.org/10.1136/archdischild-2015-309410).
45. Mansouri A, Norouzi S, Sharifi A, YektaKooshali MH, Azami M. The relationship of maternal subclinical hypothyroidism during pregnancy and preterm birth: a systematic review and meta-analysis of cohort studies. *Iran J Obstetr Gynecol Infertil*. 2017;**19**(40):69-78. Persian.
46. Azami M, Shamloo MBB, Nasirkandy MP, Veisani Y, Rahmati S, YektaKooshali MH, et al. Prevalence of vitamin D deficiency among pregnant women in Iran: a systematic review and meta-analysis. *Koomesh*. 2017;**19**(3). Persian.
47. Mansouri A, Adhami Mojarad MR, Badfar G, Abasian L, Rahmati S, Kooti W, et al. Epidemiology of *Toxoplasma gondii* among blood donors in Iran: A systematic review and meta-analysis. *Transfus Apher Sci*. 2017;**56**(3):404-9. doi: [10.1016/j.transci.2017.03.011](https://doi.org/10.1016/j.transci.2017.03.011). [PubMed: [28433405](https://pubmed.ncbi.nlm.nih.gov/28433405/)].