

Human Papilloma Virus in Head and Neck Squamous Cell Cancer

Asvadi Kermani I¹, Seifi SH¹, Dolatkah R¹, Sakhinia E², Dastgiri S³, Ebrahimi A¹, Lotfy A⁴, Esmaili HA⁵, Mohammadi G⁴, Naderpour M⁴, Hajalipour SH⁴, Asghari Haggi A¹, Nadri M¹

Abstract

Background: Epidemiologic and molecular evidences have established a strong link between high risk types of Human Papilloma Virus and a subgroup of Head and Neck Squamous Cell Carcinomas (HNSCC). We evaluated the frequency of HPV positivity in HNSCC and its relationship to demographic and some risk factor variables in an open case-control study.

Methods: Fourteen recently diagnosed patients with squamous cell cancer of oropharynx, hypopharynx and larynx aged 18-50 years were examined from 2008-2010 in Tabriz, Iran. HPV DNA was extracted from paraffin-embedded blocks of each patient's sample for PCR evaluation. Saliva samples of 94 control cancer-free subjects were collected for DNA analysis. Multivariable logistic regression method was used to calculate odds ratio for case-control comparisons.

Results: High risk HPV was detected in 6(42.8%) patients, and 6(5.3%) control subjects which was statistically significant ($p < 0.0001$). HPV-18 was the most frequent type both in the cases and controls. HPV-16 DNA was detected in two patients of the case group, but it was not detected in any of the controls. The relation between demographic and risk factor variables was not statistically significant.

Conclusion: HPV infection has a significant impact on HNSCC. Despite HPV-16 stronger impact, HPV-18 is more likely to cause malignant degeneration in such cancers amongst some communities. It is vital to introduce and conduct immunization schedules in health care systems to protect communities to some extent.

Keywords: Head and Neck Neoplasms; Squamous Cell Carcinoma; Human Papilloma Virus 16; Human Papilloma Virus 18; Sex

Please cite this article as: Asvadi Kermani I, Seifi SH, Dolatkah R, Asvadi Kermani A, Sakhinia E, Dastgiri S, Ebrahimi A, Lotfy A, Esmaili HA, Mohammadi G, Naderpour M, Hajalipour SH, Asghari Haggi A, Nadri M. Human Papilloma Virus in Head and Neck Squamous Cell Cancer. Iran J Cancer Prev. 2012; Vol5, No1, P21-26.

1. Haematology and Oncology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran
 2. Dept. of Biochemistry, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran
 3. Dept. of Social Medicine, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran
 4. Dept. of ENT, Tabriz University of Medical Sciences, Tabriz, Iran
 5. Dept. of Pathology, Tabriz University of Medical Sciences, Tabriz, Iran

Corresponding Author:
 Sharareh Seifi, MD
 Haematologist and Medical Oncologist
 Tel: (+98)41 13 34 38 11
 Email: sh_seifi@yahoo.com

Received: 2 Nov. 2011
 Accepted: 3 Dec. 2011
 Iran J Cancer Prev 2012; 1:21-26

Introduction

Head and neck cancer is the fifth [1] most common cancer in the world. Squamous cell carcinoma includes 90% of head and neck cancers [2]. Among known risk factors, smoking and alcohol consumption are considered as two major risk factors [2-5]. However, some patients do not have any obvious risk factors. In recent years, both epidemiologic and molecular evidences have established a strong link between HPV, a highly transmitted sexually virus, and the upper aerodigestive tract cancers [2, 5-8]. This association is stronger with a subset of Head and Neck Squamous Cell Carcinoma (HNSCC),

particularly oropharyngeal squamous cell carcinoma [2, 9-11]. Similar to cervical cancer in women, among more than 100 types of HPV, only high risk ones play a role in developing HNSCC [5, 12-15]. HPV-16 and HPV-18 are the two most common high risk types in this group of HNSCC respectively [5, 7, 8]. Patients with HPV-related HNSCC are usually younger, nonsmoker and non-alcoholic consumers [3, 9-13]. Therefore, it seems these tumors are different in several aspects including: molecular implication, pathological characteristics, clinical course, and response to chemoradiotherapy and outcome [6, 17, 18]. Studies conducted on molecular pathogenesis in HPV related tumors have postulated down regulation

of p53 wild type and Rb protein and up regulation of P16 that is in contrary to HPV negative tumors [16, 20, 21]. E6 and E7 oncogenes of HPV inactivate P53 and PRb respectively [1, 6, 10-15]. All of these evidences suggest that HPV positive head and neck squamous cell cancers are distinct entities compared to non HPV -related HNSCC [5, 6, 17]. Epidemiologic studies have indicated a decreasing prevalence of head and neck cancers while the incidence of oropharyngeal SCC is increasing [3, 20, 22]. As sexual contact and oral sex play roles in transmission of oral HPV infection, investigators believe that possible cause of increase in the frequency of HPV positive tumors is having either oral or vaginal sex partners [1-3, 23]. Considering the role of HPV infection in developing and increasing incidence of HPV-related HNSCC, it is crucial to design preventive methods through vaccination against infection, similar to cervical cancer. We conducted a current open case-control study to determine prevalence of HPV positive oropharyngeal SCC and other subsite of head and neck. Meanwhile, we evaluated the relation between demographic and risk factor variables and HPV positivity.

Materials and Methods

This open case-control study was carried out in Haematology and Oncology Research Center of Tabriz University of Medical Sciences with cooperation of Otolaryngology Department. Eligibility criteria included recently diagnosed 18-50 year old patients with histologically confirmed squamous cell cancer of oral cavity, oropharynx, hypopharynx and larynx. Informed consent was obtained from all patients. Exclusion criteria were the ones with the history of other malignancies including nasopharynx SCC. The control group included 94 people. Age and sex were matched with cancer-free subjects who were referred to the hospital for a routine hematologic check up. The study protocol was approved by ethics committee of the University.

Data collection

After informed consent was obtained from the participants, risk factor questionnaires containing information on demographic characteristics, oral hygiene, sexual behaviour, smoking and alcohol consumption were completed by the research team for all patients. Site and stage of tumor for each patient were determined and registered in the questionnaires.

The control group completed a self-administered risk factor questionnaire which contained information similar to the case group. Then, saliva samples of the

controls were collected and sent to the laboratory for DNA analysis by PCR (Polymerase Chain Reaction).

For the case group, formalin-fixed, paraffin-embedded tumor samples were obtained. After revising the confirmed histology report, three to four 4-micron-thick sections of paraffin-embedded blocked of each tumor sample were sent to the molecular laboratory for DNA extraction.

Laboratory Method

DNA was extracted using salting out method; and to control DNA quality, the GAPDH gene was amplified in all samples. PCR was performed in a final reaction volume of 25 μ L, containing 10 μ L of template DNA, 2.5 μ L 10X PCR buffer, 2 μ L MgCl₂ (50 mM), 1 μ L dNTPs [100 μ M], 3 μ L of My09 Primer (10 μ M), 3 μ L My11 Primer (10 μ M) and 0.5 μ L of Taq DNA Polymerase (5 U/ μ L). The PCR conditions were as follows: preheating for 5 min at 94°C followed by 40 cycles of 1 min at 94°C, 1 min at 55°C and 1 min at 72°C and a final extension of 7 min at 72°C. HPV detection by PCR was carried out in a nested-PCR system using the primers MY09/11 and GP5+/6+. Nested-PCR was performed in a final volume of 60 μ L, containing 10 μ L of the first reaction, 6 μ L 10X PCR buffer, 3 μ L MgCl₂ (50 mM), 1 μ L dNTPs [100 μ M], 6 μ L Primer GP5+ (10 μ M), 6 μ L Primer GP6+ (10 μ M) and 0.5 μ L of Taq DNA Polymerase (5 U/ μ L). The PCR conditions were as follows: preheating for 4 min at 94°C, 2 min at 40 °C and 2 min at 72 °C followed by 43 cycles of 1 min at 94°C, 2 min at 40°C and 1 min at 72°C which was followed by 1 min at 94°C, 2 min at 40 °C and 4 min at 72 °C. Amplification products obtained from HPV- positive cases were subjected to direct sequencing. Finally for genotyping of HPV, a complementary analysis of sequences obtained from blast was performed (<http://www.ncbi.nlm.nih.gov/blast>).

Statistical Analysis

Total tobacco consumption was measured as a pack-year, smoking 20 cigarettes in a year. All the participants smoked tobacco, and did not use any other substances. Classified levels for smoking were considered as: never, <1-pack year, and >1-pack year smoker. Alcohol consumption was defined as the average number of drinks per week for more than a year. We used descriptive statistics, chi-square test and odds-ratio with confidence interval of 95% for data analysis. Association was considered to be statistically significant for a two sided P<0.05.

Results

Fourteen newly diagnosed patients with squamous cell carcinoma of oral cavity, oropharynx, hypopharynx and larynx were enrolled in an open case-control study from 2008-2010 in the Haematology and Oncology Research Center of Tabriz University of Medical Sciences. Control group included 94 cancer free subjects who were sex and age matched with the patients. Mean age of patients was 39.71 ± 6.7 years with a range between 27-50 years. The mean for control group was 32.91 ± 9.14 , with age range of 18-50 years. Female patients were 57.1% and male patients were 42.9%. The control group consisted of 43(45.7%) males and 49(52.1%) females. HPV DNA was detected in paraffin-embedded block of tumor tissue of 8(57.1%) patients; of which, 6(42.9%) were high risk types, HPV-18 and HPV-16. HPV-18 was the most frequent type in 4(28.6%) patients and HPV-16 DNA was detected in two (14.3%) cases. HPV-6 (low risk type) was identified in other two patients. The frequency of HR-HPV in saliva samples of the control group was 4.3% in four persons which were all HPV-18. HPV-16 was not detected in any control persons. HPV-6 was detected in one (1.1%) subject. The relation of HPV positivity in case and control groups was statistically significant ($p < 0.0001$, OR 19.55 CI 95%, 5.1-74.92).

The mean age of patients with positive HR-HPV was 42.17 ± 5.03 years ranging from 35 to 50.

The frequency of tumor site in relation to HPV positivity is presented in Table 1.

Nine of fourteen patients were in stage IV of the disease, three in stage III and two in stage II.

HPV DNA was detected in four patients with tumor stage of IV, one in stage III, and one in stage II. For statistical analysis, tumor stages were classified in two groups: low (stages I and II), and high (stages III and IV). Association of tumor stage with HPV positivity was not statistically significant ($p = 0.95$).

The relation of all risk factors (cigarette smoking, alcohol consumption, high risk sexual behaviour, oral health) with HPV positivity was not statistically significant. Detailed data are presented in Table 2.

Discussion

The HPV prevalence in HNSCC vary in different studies based on ethnogeographic area, sample size, method of detecting HPV DNA in tissue, and classification of head and neck subsites [1,21,24]. In a meta-analysis, it has been reported that the prevalence of HPV related HNSCC has a wide range between 0-100% in small studies and 1.4-

48.8% in large ones[1]. For instance, in South Africa its prevalence is low but in Japan is high. However, the result of another study conducted in Japan showed a low prevalence of oral HPV related to Oral Squamous Cell Carcinoma (OSCC) in normal population [25]. In another recent meta-analysis carried out by Dayanni et al. the frequency of HPV related tumor in 5681 patients was 21.95% and in oropharyngeal SCC was 41%. These results are nearly similar to previous studies [21]. As a whole, the prevalence of HPV-related HNSCC is approximately 25.9% [2, 4]. In addition, HPV prevalence in OSCC including base of tongue and tonsils is higher compared to other subsites of head and neck [2-6]. In our study, HR-HPV DNA was detected in 43.5% of patients with age range of 27-50 years, considering the fact that we excluded older patients and nasopharynx SCC. In another case-control study conducted by Sahebamee et al. Iran, Tehran, the HPV prevalence in 20 patients was reported 40.9% [26]. However, in another study by Faleki et al. in Mashhad, HPV DNA was not detected in 21 young patients with oral squamous cell carcinoma [27]. Haratian et al. reported a 62.5% HPV frequency for 15 patients with HNSCC [28]. Based on the results obtained from studies carried out in Iran, prevalence of HPV related HNSCC is higher compared with developed countries. However, further studies are required to consolidate above-mentioned conclusion. Our study indicates a strong link between HPV and HNSCC in light of low prevalence (4.3%) of HR-HPV infection in control group.

According to many studies, HPV-16 is the most common high risk HPV in HPV related HNSCC [2, 4-7, 29]. However, the results of few studies have indicated that other HR-HPV types can be the commonest. For instance, Rensburg et al. showed HPV-18 was the only detected type among 66 patients with HPV positive tumors [30]. In addition, in another study in South Africa, HPV-18 was the only HPV type detected in 59 patients [24]. Kojima et al. found that HPV-38 is more common than the other types in Japan [31]. Consistent with those studies, we found that HPV-18 is the most prevalent type of HR-HPV both in the tumor tissue and saliva samples of controls. The results, however, are inconsistent with other studies carried out in Iran on the prevalence of high risk type HPV in HNSCC and cervical cancers in women where HPV-16 had been the most frequent type [22, 26, 28]. Kreimer et al. reported the frequency of HPV-18 to be low in oropharyngeal SCC compared to oral cavity and laryngeal SCC [4]. In our study, we had only two

Table 1. Tumor site frequency in relation to HPV positivity

Tumor site	HR ¹ -HPV ² positive n (%)	HR-HPV negative n(%)	Total 14
Oral cavity	2(40)	3(60)	5
Hypopharynx	2(66.7)	1(33.3)	3
Oropharynx	0(0)	2(100)	2
Larynx	2(50)	2(50)	4

1. HR: High Risk
2. HPV: Human Papilloma Virus

Table 2. Demographic and risk factor variables relation to HPV positivity

Demographic characteristics		Patients N=14	Controls N=94	P
		n (%)		
Sex	Female	8(57.1)	49(52.1)	-
	Male	6(42.9)	43(45.7)	
	Missing	-	2(2.1)	
Smoking	Smoker	3(21.4)	14(14.9)	0.460
	Nonsmoker	11(78.6)	80(85.1)	
Alcohol	User	1(7.1)	10(10.6)	0.99
	Nonuser	13(92.9)	84(89.4)	
Oral Sex	Never	10(71.4)	67(71.3)	0.99
	Ever	4(28.6)	27(28.7)	
Oral hygiene	Good	5(35.7)	50(53.2)	0.262
	Poor	9(64.3))	44(46.8)	
Number of sexual partner	1	12	91(96.8)	0.065
	>1	2(14.3)	3(3.2)	

patients with oropharyngeal SCC and this could be the reason for the higher prevalence of HPV-18 in our study. On the other hand, HPV-18 was the most frequent type in the control group as well. Other contributory factors affecting our results such as different nature of North West Iran, using different techniques for PCR and DNA sequencing should be considered. To achieve a concrete conclusion conducting further studies with larger sample size are recommended.

Some studies have shown the relevance of HPV positive HNSCC and more advanced disease and nodal metastases [2, 3, 9]. In our study, although the majority of patients had advanced stage of the disease (stage III and IV), this relation with HPV positivity was not significant due to small sample size.

Although tobacco smoking and alcohol consumption are main risk factors for HNSCC, numerous studies have demonstrated that HPV-related HNSCC patients are less likely to smoke and consume alcohol [2, 3, 16]. There is some controversy over the impact of smoking and alcohol consumption on HPV infection in HNSCC [34-36]. This is contrary to cervical cancer in women where smoking has an

obvious synergistical effect with HPV. Some studies have suggested additive effects of smoking on HPV positive oral cavity and oropharyngeal SCC. Other investigators have shown that HPV seropositive smokers have a higher risk for developing HNSCC [34, 37, 38]. Smith et al. reported that tobacco and alcohol have synergistical effect on positive HR-HPV [11]. Dr. Park's showed that tobacco smoking and drinking alcohol help promote HPV invasion [39]. Considering the small sample size; we did not find any link between cigarette smoking and alcohol consumption and HPV positive tumor. Overall, further studies are needed to determine the impact of smoking and alcohol on HPV related HNSCC.

Although poor oral hygiene may be considered a risk factor for HNSCC [14, 17, 40], no association between HPV related HNSCC has been defined in the literature [17]. In our study, we did not find this relation either.

Accumulating data have revealed high risk sexual behaviour including oral sex and multiple sexual partners to be the risk factors for HPV related HNSCC [20, 23, 41]. In our work, this relation was not shown properly due to few patients having oral sex

or more than one sexual partner. It should be mentioned that some patients had social limitations to talk about their sexual practices. Another limitation in our study included small sample size. As HNSCC, except for nasopharynx SCC, within the age range of 18-50 years was too rare in our hospital, it is difficult to comment on demographic and risk factor variables in association with HPV DNA detection.

Conclusion

HR-HPV has a definite impact on HNSCC. Given that HPV-18 is the most prevalent type in this study, the need to design and introduce a preventive program through vaccination seems more logical. Further studies are required to identify risk factors related to HPV infection.

Acknowledgement

We would like to thank the Research Deputy of Tabriz University of Medical Sciences for financial support of this project.

Conflict of Interest

There is no conflict of interest in this study.

Authors' Contribution

IAK designed the study, ShS collected the data and wrote the article, RD and AAK collected the data and revised the manuscript, ES, AE, AAH and MN contributed to the laboratory work, SD contributed to the analysis and interpretation of the data, AL, GM, MN and ShH contributed to patients diagnosis, HAE contributed to pathologic diagnosis. All authors read and approved the final revision.

References

1. Termine N, Panzarella V, Falaschini S. HPV in oral squamous cell carcinoma vs head and neck squamous cell carcinoma biopsies: a meta-analysis (1988–2007). *Annals of Oncology*. 2008; 19:1681–90.
2. Syrjänen S. Human Papillomaviruses in Head and Neck Carcinomas. *N Engl J Med*. 2007; 356:1993–5.
3. Chaturvedi AK, Engels EA, Anderson WF. Incidence trends for human papillomavirus-related and -unrelated oral squamous cell carcinomas in the United States. *J Clin Oncol*. 2008; 26:612–19.
4. Kreimer AR, Clifford GM, Boyle P, Franceschi S. Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. *Cancer Epidemiol Biomarkers Prev*. 2005 Feb; 14(2):467–75.
5. DeVita VT, Lawrence TS, Rosenberg SA, DePinho RA, Weinberg RA. DeVita, Hellman, and Rosenberg's *Cancer: Principles and Practice of Oncology*, 8th ed. Lippincott Williams and Wilkins, USA; 2008:799–876.
6. Hobbs CG, Sterne JA, Bailey M, Heyderman RS, Birchall MA, Thomas SJ. Human papillomavirus and head

and neck cancer: a systematic review and meta-analysis. *Clin Otolaryngol*. 2006 Aug; 31(4):259–66.

7. WSOC Charlotte-Human Papillomavirus (HPV). [<http://www.wsocvtv.com/print/14978959/detail.html>]. Accessed January 4, 2008.

8. Mork J, Lie AK, Glatte E, Hallmans G, Jellum E, Koskela P, et al. Human papillomavirus infection as a risk factor for squamous-cell carcinoma of the head and neck. *N Engl J Med*. 2001 Apr 12; 344(15):1125.

9. Anderson CE, McLaren KM, Rae F, Sanderson RJ, Cuschieri KS. Human papillomavirus in squamous carcinoma of the head and neck: a study of cases in southeast Scotland. *J Clin Pathol*. 2007; 60:439–41.

10. Ramqvist T, Dalianis T. Oropharyngeal Cancer Epidemic and Human Papillomavirus. *Emerg Infect Dis*. 2010 Nov; 16(11):1671–7.

11. Smith EM, Ritchie JM, Summersgill KF, Hoffman HT, Wang DH, Haugen TH, et al. Human papillomavirus in oral exfoliated cells and risk of head and neck cancer. *J Natl Cancer Inst*. 2004; 96(6):449–55.

12. Shukla S, Bharti AC, Mahata S, Hussain S, Kumar R, Hedau S, et al. Infection of human papillomaviruses in cancers of different human organ sites. *Indian J Med Res*. 2009 Sep; 130(3):222–33.

13. Kreimer AR, Bhatia RK, Messegue AL, González P, Herrero R, Giuliano AR. Oral human papillomavirus in healthy individuals: a systematic review of the literature. *Sex Transm Dis*. 2010 Jun; 37(6):386–91.

14. D'Souza G, Kreimer AR, Viscidi R. Case-control study of human papillomavirus and oropharyngeal cancer. *N Engl J Med*. 2007; 356:1944–56.

15. Ragin CC, Modugno F, Gollin SM. The epidemiology and risk factors of head and neck cancer: a focus on human papillomavirus. *J Dent Res*. 2007 Feb; 86(2):104–14.

16. Haraf DJ, Nodzinski E, Brachman D, Mick R, Montag A, Graves D, et al. Human papilloma virus and p53 in head and neck cancer: clinical correlates and survival. *Clin Cancer Res*. 1996 Apr; 2(4):755–62.

17. Gillison ML, D'Souza G, Westra W, Sugar E, Xiao W, Begum S, et al. Distinct risk factor profiles for human papillomavirus type 16-positive and human papillomavirus type 16-negative head and neck cancers. *J Natl Cancer Inst*. 2008 Mar 19; 100(6):407–20.

18. Syrjänen S. The role of human papillomavirus infection in head and neck cancers. *Ann Oncol*. 2010 Oct; 21 Suppl 7:vii243–5.

19. Gillison ML. Human papillomavirus and prognosis of oropharyngeal squamous cell carcinoma: implications for clinical research in head and neck cancers. *J Clin Oncol*. 2006 Dec 20; 24(36):5623–5.

20. Psyrris A, Gouveris P, Vermorken JB. Human papillomavirus related head and neck tumor: clinical and research implication. *Curr Opin Oncol*. 2009 May; 21(3):201–5.

21. Dayyani F, Etzel CJ, Liu M, Ho CH, Lippman SM, Tsao AS. Meta-analysis of the impact of human papillomavirus (HPV) on cancer risk and overall survival in head and neck squamous cell carcinomas (HNSCC). *Head Neck Oncol*. 2010 Jun 29; 2:15.

22. Lassen P, Eriksen JG, Hamilton-Dutoit S, Tramm T, Alsner J, Overgaard J. Effect of HPV-associated p16INK4A expression on response to radiotherapy and survival in squamous cell carcinoma of the head and neck. *J Clin Oncol.* 2009 Apr 20; 27(12):1992-8. Epub 2009 Mar 16.
23. Chaudhary A K, Singh M, Sundaram Sh. Role of human papillomavirus and its detection in potentially malignant and malignant head and neck lesions: updated review. *Head Neck Oncol.* 2009; 1: 22.
24. Boy S, Van Rensburg EJ, Engelbrecht S, Dreyer L, van Heerden M, van Heerden W. HPV detection in primary intra-oral squamous cell carcinomas commensal, aetiological agent or contamination? *J Oral Pathol Med.* 2006 Feb; 35(2):86-90.
25. Kurose K, Terai M, Soedarsono N, Rabello D, Nakajima Y, Burk RD, et al. Low prevalence of HPV infection and its natural history in normal oral mucosa among volunteers on Miyako Island, Japan. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004 Jul; 98(1):91-6.
26. SahebJamee M, Boorghani M, Ghaffari SR. Human papillomavirus in saliva of patients with oral squamous cell carcinoma. *Med Oral Patol Oral Cir Bucal.* 2009 Oct 1; 14 (10):e525-8.
27. Falaki F, Dalirsani Z, Pakfetrat A, Falaki A, Saghraevanian N, Nosratzahi T, et al. Clinical and Histopathological analysis of oral squamous cell carcinoma of young patients in Mashhad, Iran: A retrospective study and review of literature. *Med Oral Patol Oral Cir Bucal.* 2011 Jul 1; 16(4):e473-7.
28. Haratian K, Mohseni Meybodi A, Zari Moradi S, Vosough P. Detection of High Risk Human Papillomavirus DNA Sequences in Head and Neck Squamous Cell Carcinoma in Iranian Fanconi Anemia Patients. *Yakhteh Medical Journal.* Spring 2010; 12(1): 43-50.
29. Mannarini I, Kratochvil V, Calabrese L. Human Papilloma Virus (HPV) in head and neck region: review of literature. *Acta Otorhinolaryngol Ital.* 2009 June; 29(3): 119-26.
30. Van Rensburg EJ, van Heerden WF, Venter EH, Raubenheimer EJ. Detection of human papillomavirus DNA with in situ hybridisation in oral squamous carcinoma in a rural black population. *S Afr Med J.* 1995 Sep; 85(9):894-6.
31. Kojima A, Maeda H, Sugita Y, Tanaka S, Kameyama Y. Human papillomavirus type 38 infection in oral squamous cell carcinomas. *Oral Oncol.* 2002 Sep; 38(6):591-6.
32. Jabbarpour Bonyadi M, Esmaeili M, Dastranj A. Oncogene HPV types determined by PCR in multiple lesions of cervical cancer in the North West of Iran. *Journal of Infectious Diseases and Tropical Medicine.* 2008; 13(41):29-34.
33. Mortazavi S, Zali M, Raoufi M, Nadji M, Kowsarian P, Nowroozi A. The Prevalence of Human Papillomavirus in Cervical Cancer in Iran. *Asian Pac J Cancer Prev.* 2002; 3(1):69-72.
34. Esquenazi D, Bussoloti Filho I, Carvalho Mda G, Barros FS. The frequency of human papillomavirus findings in normal oral mucosa of healthy people by PCR. *Braz J Otorhinolaryngol.* 2010 Feb; 76(1):78-84.
35. Xi LF, Koutsky LA, Castle PE, Edelstein ZR, Meyers C, Ho J, et al. Relationship between cigarette smoking and human papillomavirus type 16 and 18 DNA load, Cancer Epidemiol Biomarkers Prev. 2009; 18(12):3490-6.
36. Simen-Kapeu A, Kataja V, Yliskoski M, Syrjänen K, Dillner J, Koskela P, et al. Smoking impairs human papillomavirus (HPV) type 16 and 18 capsids antibody response following natural HPV infection. *Scand J Infect Dis.* 2008; 40(9):745-51.
37. Silva KC, Rosa ML, Moyse N, Afonso LA, Oliveira LH, Cavalcanti SM. Risk factors associated with human papillomavirus infection in two populations from Rio de Janeiro, Brazil. *Mem Inst Oswaldo Cruz.* 2009 Sep; 104(6):885-91.
38. Alam S, Conway MJ, Chen HS, Meyers C. The cigarette smoke carcinogen benzo[a]pyrene enhances human papillomavirus synthesis. *J Virol.* 2008 Jan; 82(2):1053-8. Epub 2007 Nov 7.
39. Mouth Cancer and the Human Papillomavirus. [<http://www.rdoc.org.uk/hpv.html>]
40. Guha N, Boffetta P, Wunsch Filho V. Oral Health and Risk of Squamous Cell Carcinoma of the Head and Neck and Esophagus: Results of Two Multicentric Case-Control Studies. *Am J Epidemiol.* 2007; 166 (10): 1159-73.
41. Kreimer AR, Alberg AJ, Daniel R, Gravitt PE, Viscidi R, Garrett ES, et al. Oral human papillomavirus infection in adults is associated with sexual behavior and HIV serostatus. *J Infect Dis.* 2004 Feb 15; 189(4):686-98.