

The Impact of Laughter Yoga on the Stress of Cancer Patients before Chemotherapy

Shadi Farifteh¹, Alireza Mohammadi-Aria², Alireza Kiamanesh¹, Bahram Mofid³

Abstract

Background: Cancer is usually accompanied by considerable stress for the sufferer, and the stress has destructive effects on Chemotherapy treatment process. Therefore, the current research deals with the effect of yoga laughter on the cancer patients' stress before chemotherapy.

Methods: In this research, as the first step, 37 cancer sufferers, who had been hospitalized in Shohada Tajrish Hospital (Behnam Daneshpoor Charity Organization) and had the requirements necessary for being taken as research samples, were selected for data collection. The mentioned patients were classified randomly in experimental and control groups. Collected data were analyzed by the multi-variable covariance analysis test.

Results: The results show there is a meaningful difference in the stress average before and after interference in the test group ($p < 0.05$).

Conclusion: Laughter yoga can decrease the stress in cancer sufferers before chemotherapy.

Keywords: Laughter yoga; Stress; Cancer; Chemotherapy

Please cite this article as: Farifteh SH, Mohammadi-Aria AR, Kiamanesh AR, Mofid B. The Impact of Laughter Yoga on the Stress of Cancer Patients before Chemotherapy. *Iran J Cancer Prev.* 2014;7(4):179-83.

Introduction

In terms of social aspects, cancer disturbs daily functions and social activities of the sufferer, and alters his or her abilities to fulfill their normal roles and responsibilities, up to an extent that creates some new roles. The nature of the new-born roles may cause a feeling of inefficiency and social isolation in the sufferer [1]. In fact, after cancer is diagnosed, the patient will be hit by a crisis, his self-confidence will be at stake, his individual relationships will be disturbed due to uncertainty about future, the previous adaptation mechanisms will seem insufficient, and being hospitalized repeatedly, the patient will be flooded with solitude [2]. A set of factors mentioned above lead to psychological stress in the patient [3]. Cancer is a prevailing, chronic, non-infectious disease [4]. The exact reason is unknown, but genetic and exterior factors like viruses and carcinogen chemicals are likely to be influential [5]. Taking anti-cancer medicines and chemotherapy are some treatments for tumors, though they can be mixed with operation and radiotherapy. Chemotherapy is very effective in treating tumors, and the effect is increasing nowadays by implementing newly-discovered anti-

1. Dept. of Psychology, Islamic Azad University Science and Research Branch, Tehran, Iran

2. Dept. Educational Psychology, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

3. Radiotherapy Group, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Corresponding Author:

Shadi Farifteh, MA;

Tel: (+98) 936 3892600

Email: shadi.farifteh@gmail.com

Received: 28 Nov. 2013

Accepted: 15 Sep. 2014

Iran J Cancer Prev. 2014; 4:179-83

tumor medicines. Conducted studies indicate that cancer is the second cause of Americans' death after cardiovascular diseases. As an effective treatment for increasing life-expectancy in the patients, chemotherapy has a significant role [6]. Research shows that the patients get stressed before chemotherapy, and it causes body resistance against treatment and disturbs the process. Having done a set of tests about breast cancerous cells, researchers have found out that the protein created after stress helps the cancerous cells continue living, getting adapted and resisting against the treatment [7].

Research has shown that immune system is effective in controlling and healing cancer, in other words, a weakening immune system raises the potentials to develop various types of cancer. Stress has been discussed as a suppressing factor of the immune system, so there is a possibility of cancerous cell growth after stress. Studies have indicated that stressed people are more likely to develop breast cancer [1]. Therefore, looking for suitable methods of stress reduction seems necessary. One of non-medical ways of stress reduction is laughing and being in delightful environments. Laughter is a positive sensation, and

seems to be a useful and healthy way to overcome stress. Decreasing stress-making hormones floating in blood, laughter removes the effects of stress. Nowadays, there is evidence indicating that being humorous and exposed to funny stimuli can have a positive influence on the side-effects of some specific diseases. Laughter is an emotional reaction influencing human's individual and social life. It enjoys features that distinguish it from other humane emotional reactions. Happiness increases Cortisol hormone and boosts body immunity against diseases. Laughter creates a balance in the chemicals and hormones of the body [8]. Lidi maintains that fun and laughter decrease red globule sedimentation level and increase natural killer-cells' activity, reduce cortisol and blastogene lymphocytes and boosts monoglobin A as well [9]. Fri (1971) declares that laughter strengthens the immune system, facilitates phagocytosis process in the immune system and also helps the body to fight against infections. Dogan believes that biological functions of laughing and crying cause biochemical changes in the body and reduces the tension made by painful emotions like fear and anger. Tears and nasal discharge during laughing and crying contain humorous, steroids and toxins accumulated in the body during tension. Laughing and crying both increases coolamine production in the blood [10].

A way to make people laugh is laughter yoga, innovated by Kataria (1995) in India. Laughter yoga is the only way enabling people to laugh heartily without getting involved in logical thinking. This method puts wisdom and logic aside as natural laughter obstacles. The reason the Cataria named his innovative method as laughter yoga was that he had mixed the respiratory exercises of yoga –called paranayama- into laughter exercises. The former exercises are very strong and effective, and they have been applied for thousands of years to influence human body, psyche and emotions in a positive manner. Based on yoga philosophy, our life is subject to a flow of universal energy into our bodies via breathing. This is called Yaprana vital energy. As a result of stress and negative tensions, our respiration gets irregular and shallow, and this disturbs the flow of energy in our bodies [11].

It should be noted that the studies on this domain have examined laughter effects using the mentioned methods and other variables reducing stress, and all of these studies have regarded the long-term effects of laughter, while the short-term effects of laughter on the stress in patients with

cancer is being dealt with in the current research using laughter yoga. Therefore, in this study, the question is if laughter yoga is effective on cancer-sufferers stress level before Chemotherapy.

Materials and Methods

Statistic population of the research was patients hospitalized in Shohada Tajrish hospital for chemotherapy in the period of May to August, 2013. Sample population data is presented in this part separately for control and experimental groups. Table 1 shows distribution of samples in terms of groups, education and genders for sampling, those qualified patients were selected for trial, which were randomly placed in experimental and control groups. The minimum population of sample for each of control and experimental groups is 15 in semi-experimental studies [12]. Accordingly, researchers had initially selected 25 samples to avoid sample drop, 2 patients in experimental group and 11 in control group were omitted from the sample list in pre-test due to leaving the hospital, moving to other clinics or lacking interest for taking part in the research. There ultimately remained 23 patients in experimental group and 14 in control group.

This study examined the effects of laughter yoga on the stress before chemotherapy.

Instruments

Stress measuring questionnaire: Questionnaire QSC-R23 (Questionnaire on Stress in Cancer Patients) was used in this research to measure stress (the questionnaire is attached). This questionnaire is specially related to specific diseases and evaluates psychological stress in cancer sufferers. Questionnaire articles were divided into 5 equal scales, including psych-physical complaints, fears, information defects, daily-life limitations and social conflicts. QSC was first formed in Germany after passing different phases, namely exact interviews, examining the initial versions of the questionnaire and measuring their validity and constancy by Herchbach. Cronbach's Alpha method of analysis was implemented to estimate tools constancy in the research. It includes 23 articles, showing daily-life potential stress in all perspectives. The answers to the questions included two parts: first, the article did not apply to the respondent (zero degree), second, respondent's conditions conformed to the specifications in the article. If the second part was selected, the answer should have determined the degree to which the subject matter of that article was

Table 1. It shows distribution of samples in terms of groups, education and genders.

Group	Education	Abundance	percentage	Gender	Abundance	percentage
Experiment	Elementary	4	17.4	Male	9	39
	Mid-school	5	21.7	Female	14	61
	High-school	7	30.4			
	University	7	30.4			
	Total	23	100	total	23	100
Control	Elementary	2	14.3	Male	5	35.7
	Mid-school	2	14.3	Female	9	64.3
	High-school	4	28.6			
	University	6	42.9			
	Total	14	100	total	14	100
Total	Elementary	6	16.2	Male	14	37.8
	Mid-school	7	18.9	Female	23	62.2
	High-school	11	29.7			
	University	13	35.1			
	Total	37	100	total	37	100

affecting him/her. This degree can be positioned on a scale of 1 (very small problem) to 5 (very serious problem).

Procedure

First, QSC-R23 questionnaire was administered to the patients accompanied by exact and complete explanations about it, and they were asked to fill it out in the researcher's presence. In case patients were unable to fill out the forms, the researcher took responsibility herself. Having the forms filled out in control and experimental groups, the experimental group was treated by laughter yoga by a trainer for 20 to 30 minutes under researcher's observation. After that, the tastes in both groups were asked to fill out the questionnaire again. Having conducted the research, laughter yoga was performed in control group as well to conform to the ethics. The data

obtained was then analyzed by descriptive statistic methods such as mean and deviation, and the multi-variable covariance analysis test was applied to examine the research hypothesis.

Results

According to table 2, there is a meaningful difference between the pre-test and the post-test average; however, in control group there is no difference between them. Finally, the results of multivariate covariance analysis between variables in both experimental and control groups are shown in table 3, in which there is no meaningful difference in Psych-physical complaints, Information defects and Total stress score.

Table 2. It shows Mean and Standard Deviation of studied variables for the pre-test and the post-test in each group.

Variables	Pre-test / post-test	experiment group		Control group	
		Mean	SD	Mean	SD
Psych-physical complaints	Pre- test	1.7	1.3	1.5	0.9
	Post-test	1.1	1.1	1.5	1.1
Fear	Pre-test	2.4	1.2	2.8	1.5
	Post-test	1.7	1.6	2.7	1.8
Information defects	Pre-test	2.3	1.3	1.9	1.5
	Post-test	1.5	1.1	2	1.6
Daily-life limitations	Pre-test	2.5	1.2	2.5	1.2
	Post-test	2	1.3	2.4	1.4
Social conflicts	Pre-test	1	1.2	0.7	0.7
	Post-test	0.5	0.8	0.6	0.8
Total stress score	Pre-test	8.9	4.4	8.5	3.8
	Post-test	6.1	4.2	8.4	4.4

Table 3. It shows the results of multivariate covariance analysis between variables in both experimental and control groups.

Dependent variable	square	Freedom degree	Mean square	F	Significant
Psych-physical complaints	1.4	1	1.4	4.3	0.05
Fear	2.2	1	2.2	2.4	0.13
Information defects	3.7	1	3.7	6.9	0.01
Daily-life limitations	0.8	1	0.8	1.2	0.28
Social conflicts	0.4	1	0.4	1.5	0.23
Stress total score	29.8	1	29.8	4.9	0.03

Discussion

Findings show that the intervention was effective in making a meaningful difference in three sub-scales of psych-physical complaints, fear, information defect and total score of stress in the two groups. It indicates that laughter yoga reduces stress in cancer sufferers before chemotherapy.

The result of this research is in line with silva [13], Chaya and crew [14], Marcon [15] and Beke [16], and contrary to Omrani [17].

According to the research, stress increases cancerous cells activity to a meaningful extent, and causes the involved cells to resist against chemotherapy. Using a method to decrease the patient's stress is of a huge prominence.

Conclusion

The findings of the research have shown that as a natural gift, laughter can reduce the stress in cancer sufferers before chemotherapy. Laughter plays an important role in reducing stress, which is done by Endorphin Secretion resulting in mental and physical relaxation. Chronic Stress impacts Limbic system and Hypothalamus continuously and leads to Adrenaline secretion and causes disruption in immune system. Indeed, Laughter balances sympathetic and parasympathetic system; moreover, it functions as an anti-stress. Furthermore, laughter increases Endorphin in brain. Endorphin seems to be the most easily linked structure to morphine, both function in the exhilaration and lessening of pain.

To illustrate this according to the study background, it can be noted that laughter not only creates good spirits in patients and the hospital staff, but also it can reduce the stress in

patients and improve the treatment process without any harmful side-effects by creating a happy environment at hospitals.

Examining the data brings us to conclusion that laughter has a significant effect on the spirits of patients and their stress.

Suggestion for Further Research

In this study, the questionnaire QSC-R23 was applied as the measuring tool. It is strongly suggested that the study under the same title should be conducted using other tools, such as pulse and heart beat measuring in addition to self-reporting to undermine the effect of the samples on the result. Moreover, according to the role of laughter in reducing stress, hospitals had better be equipped with laughter rooms. Doctors and nurses must be trained to be on duty smiling, and finally, comedy shows should be played for the sufferers.

Acknowledgment

We would like to thank the members of the Behnam Daneshpoor Charity Organization in Shohada Tajrish Hospital, Tehran, Iran. Special thanks to Alireza Mohammadi Aria and advisor, Alireza Kiamanesh.

Conflict of Interest

The authors have no conflict of interest in this study.

Authors' Contribution

Shadi Farifteh designed and wrote this article and collected and analyzed the data, with the help of supervisor, Alireza Mohammadi Aria and advisor, Alireza Kiamanesh. All authors read and approved the final manuscript.

References

1. Hann D, Baker F, Denniston M, Gesme D, Reding D, Flynn T, et al. The influence of social support on depression symptoms in cancer patients: Age and Gender Differences. *J Psychosom Res.* 2002;52(5):279-83.
2. Tan M. Social support and coping in Turkish patients with cancer. *Cancer Nurs.* 2007;30(6):498-504.
3. Hall S, Chochinov H, Harding R, Murray S, Richardson A, Higginson IJ. A phase II randomised controlled trial assessing the feasibility, Acceptability and potential effectiveness of dignity therapy for older people in care homes: Study Protocol. *BMC Geriatrics.* 2009;24(9):9.
4. Farzanpei R. Cognition about cancer: prevention and treatment. Tehran: KetabeMad Publications; 1994.
5. Zolala F, Ayat Elahi S.A.R., Shahriary M, Ayat Elahi S.M.T. Determination of Incidence rate of acute lymphoblastic leukemia in children under the age of 15 in Fars province in 2001. *Armaghan Danesh.* 2004;9(35):64-71.
6. Goldman L, Benett G. Text book of medicine. Philadelphia: WB Saunders Co; 2000.
7. Stress May Help Cancer Cells Resist Treatment [Internet] 2011 [updated 2007 April 11; cited 20 July 2012]. Available from: <http://www.sciencedaily.com/releases/2007/04/070410103023.htm>
8. Erfani V. Miracle of Laughter. Pain J. 2004;5(43):19. [Full Text in Persian].
9. Saper B. The therapeutic use of humor for psychiatric disturbance of adolescents and adults. *Psychiatric Quarterly.* 1990;61(4):261-72.
10. Martin RA, Dobbin JP. Sense of humor, hassles and Immunoglobulin A: Evidence for a stress-moderation effect of humor. *International Journal of Psychiatry in Medicine.* 1988;18(2):93-105.
11. Kataria M. Certified laughter yoga leader training (CLYL) [internet]. 2011 [updated 2011 August 1; cited 6 July 2012]. Available from: <http://laughteryoga.org/downloads/Leader%20Manual.pdf>.
12. Delavar A, Naghshbandi S. Statistical Analysis In Psychology and Education. Tehran: Arasbaran; 2009.
13. Silva E. Nursing concepts of stress adaptation. *Rehabilitation Nursing.* 2001;18:25-8.
14. Chaya MS. What are the effects of laughter yoga on stress in the workplace? The Research Institute SVYASA [internet]. 2007 [updated 20 May 2012; cited 12 June 2012]. Available from: <http://www.yogalaff.com/docs/6-INDIA-STUDY-SUMMARY.pdf>.
15. Marcon M. Humour for good health; The Northern Hospital; 2005. The emergency department and child and adolescent health unit (The Clown Doctor Project) Final report, June 2005.
16. Bacqué, M F. Association rire clowns pour enfants hospitalisés. *Psycho-Oncol.* 2010;4(1):65-6.
17. Omrani R. Investigation of music therapy and laughter therapy on stress reduction for women before surgery [M.A. thesis]. Tehran: Faculty of psychology and education. Payame Noor University; 2010.