

Original Article**Fauna and Larval Habitat Characteristics of Mosquitoes in Neka County, Northern Iran**

Seyed Hassan Nikookar ¹, *Seyed Hassan Moosa-Kazemi ², Mohammad Reza Yaghoobi-Ershadi ², Hassan Vatandoost ^{2,3}, Mohammad Ali Oshaghi ², Abolfazl Ataei ², Mehdi Anjamrooz ²

¹Health Sciences Research Center, Department of Medical Entomology and Vector Control, Faculty of Health, Mazandaran University of Medical Sciences, Sari, Iran

²Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³Institute for Environmental Research, Tehran University of Medical Sciences, Tehran, Iran

(Received 29 May 2012; accepted 12 Oct 2013)

Abstract

Background: Ecological studies on mosquitoes are very important in vector control programs. There are a few studies about the ecology of mosquitoes in northern Iran. This study was carried out to detect fauna and larval habitat characteristics of mosquitoes.

Methods: This study aimed to determine fauna and the ecology of mosquitoes in Neka County, Mazandaran Province, northern Iran from April to December, 2009. The larval collection was conducted using standard dipper, and the characteristics of larval habitat were investigated based on degree of transparency of water, type of water (stagnant or running), plant vegetation, sunny or shady, temperature and altitude of the natural or artificial breeding places.

Results: The mosquito larvae were collected from 72 habitats and identified using systematic keys. Nine species of mosquitoes were identified: *Anopheles claviger* (0.31%), *An. maculipennis* (0.54%), *An. plumbeus* (10.28%), *An. superpictus* (0.01%), *Culiseta annulata* (1.07%), *Cs. longiareolata* (8.91%), *Culex mimeticus* (0.03%), *Cx. pipiens* (63.99%), and *Ochlerotatus geniculatus* (14.85%). The range of temperature in the larval habitats was 19.6–22.5 °C. Significant difference was observed in the rate of temperature among the species in the larval habitats ($P < 0.05$). A checklist of mosquitoes including seven genera and 32 species has been provided for Mazandaran Province.

Conclusion: The most dominant species were *Cx. pipiens*. They were collected from the larval habitats like Border Rivers, ponds, rain water pools, discarded tires and tree holes. *Culiseta annulata* was included to the checklist of mosquitoes in Mazandaran Province.

Keywords: Larval habitat, Culicidae, Ecology, Iran

Introduction

According to the latest classification of Culicidae, the family includes two subfamilies, 11 tribes, 112 genera and 3537 species. The subfamily Anophelinae has three genera and Culicinae has 109 genera (Harbach 2007).

Many of the world's major diseases such as Malaria, West Nile, Sindbis, Dirofilariasis, Dengue fever, Yellow fever, Rift valley fever are transmitted by mosquitoes. West Nile and Sindbis viruses have been reported in northern

parts of Iran (Naficy and Saidi 1970, Saidi et al. 1976). The mosquito borne filarial worms, *Dirofilaria immitis* and *D. repens*, *Setaria* (*S. labiatopapillosa*, *S. digitata* and *S. equina*) and *Diptaltonema evansi* (camel filariasis) have been reported in Iran (Siavashi and Massoud 1995, Eslami 1997, Azari-Hamidian et al. 2007, Oryan et al. 2008). *Anopheles maculipennis* and *Culex theileri* were reported as vectors of *Setaria labiatopapillosa* and

Dirofilaria immitis respectively, in Ardebil Province (Azari-Hamidian et al. 2009). Dengue fever cases were reported in Tehran, imported from Malaysia, India and Thailand. In addition, six DF cases were reported from Sistan and Baluchistan Province in southeast Iran (Chinikar et al. 2010, 2012).

The systematic key of *Anopheles* species in Iran was reported by Shahgudian (1960). Many studies have been carried out about the genus of *Culex* and its medical importance (Ghaffary 1956, Lotfi 1970, Lotfi 1973, Lotfi 1976, Zaim et al. 1984a, 1984b, 1985, 1986). The results of these studies have led to the publishing of the Iranian Culicinae checklist (Zaim and Cranston 1986). The role of mosquitoes in malaria transmission and its geographical distribution have been studied by Saebi (1987). The fauna and ecological characteristics of Culicidae with emphasis on their medical importance were conducted in Isfahan and Hormozgan Provinces as well as Chabahar County (Mousakazemi et al. 2000, Moosa-Kazemi et al. 2005, 2009) and Guilan Province (Azari-Hamidian et al. 2002a). Moreover, the study on the genus of *Aedes* is very scattered in the northern parts of the Iran. Azari-Hamidian et al. (2002a) stressed the occurrence of *Oc. caspius*, *Oc. echinus*, *Ae. vexans* and *Oc. geniculatus* in Guilan Province, northern Iran. More than 30 culicidae species reported from Guilan province (Azari-Hamidian et al. 2003a).

Distribution of larval habitat of *Cx. pipiens* was reported by Dehghan et al. (2011) in central Iran. Among the 467 species of *Anopheles* identified in the world (Harbach 2007), about 70 *Anopheles* species are malaria vectors in which about 40 species are important vectors (Service 1993). *Anopheles maculipennis* complex, one of the most important vectors of malaria, was reported in northern Iran (Eslami 1997, Azari-Hamidian 2007a). *Anopheles maculipennis*, and *Cx. pipiens* were reported as dominant species in

Isfahan Province, central Iran (Mousakazemi et al. 2000a) and in East Azarbaijan and Ardebil Provinces (Abai et al. 2007, Azari-Hamidian et al. 2009).

By now, seven species of malaria vectors have been reported in Iran (Vatandoost et al. 2009a). They include: *An. culicifacies*, *An. dthali*, *An. fluviatilis*, *An. maculipennis*, *An. sacharovi*, *An. stephensi*, and *An. superpictus* (Edrissian 2006, Vatandoost et al. 2008, Vatandoost et al. 2009a, Moosa-Kazemi et al. 2010). *Anopheles pulcherrimus* has been identified as potential vector in Iran using serological method (Zaim et al. 1993). Although oocyte of *Plasmodium* was found in the stomach of *An. multicolor* but sporozoite never found in its salivary gland and not considered as the malaria vector (Eshghy 1977). *Anopheles hyrcanus* was found as a malaria vector using the PCR method in northern Iran (Dinparast Djadid et al. 2009). At present, 64 species, three subspecies and seven genera of Culicidae have been reported from Iran (Azari-Hamidian 2007a).

The first study about the mosquitoes in Mazandaran Province was carried out by Gutsevich (1943) and Zolotarev (1945). The occurrence of the species of *An. maculipennis*, *An. melanoon* (*An. subalpinus*), *An. Superpictus*, *An. sacharovi* and *An. messeae* have been reported by Zolotarev (1945). However, Gutsevich (1943) reported the species of *An. pulcherimus*, *Ae. vexans*, *Oc. geniculatus* and *Cq. richiardii* in Mazandaran Province. Subsequently, Garrett Jones (1951) reported the species of *An. claviger*, *An. maculipennis*, *An. plumbeus* and *An. hyrcanus* in Klardasht. Institute of Parasitology and Malariology (1953) reported the species of *An. marteri* in Ramsar. Dow (1953) reported the species of *An. maculipennis*, *An. melanoon*, *An. subalpinus* (synonym of melanoon), *An. hyrcanus*, *An. pseudopictus*, *An. superpictus*, *An. sacharovi*, *Cx. pipiens*, *Cx. mimeticus* and *Cx. tritaeniorhynchus*. However, Faghih (1969) stressed the occurrence of the species

of *An. melanoon*, and *An. messeae*. Minar (1974) reported the species of *An. Maculipennis*, *An. pseudopictus*, *Oc. pulcritarsis*. However, Zaim et al. (1984a) cited the species of *Ae. vexans*, *Oc. echinus*, *Oc. geniculatus* and *Oc. pulcritarsis*. Zaim (1987) cited the species of *Ae. vexans*, *Oc. echinus*, *Oc. geniculatus*, *Oc. pulcritarsis*, *Cx. perexiguss*, *Cx. theileri*, *Cx. mimeticus*, *Cx. Tritaeniorhynchus*, *Cx. hortensis*, *Cx. pipiens*, *Cs. longiareolata* and *Cs. subochrea*. Saebi (1987) stated the species of *An. plumbeus*, *An. hyrchanus*, *An. pseudopictus*, *An. maculipennis* and *An. claviger*. Momeni (1991) reported the species of *An. messeae*, *An. melanoon*, and *An. maculipennis*. Glick (1992) reported the species of *An. pseudopictus*. Karami (2000) cited the presence of *Cx. bitaeniorhynchus*, *Ur. unguiculata*, *Oc. caspius*. Sedaghat et al. (2003) reported the species of *An. Maculipennis*, *An. sacharovi*, *An. persiensis* in Mazandaran Province. Sedaghat and Harbach (2005) stated the occurrence of the species of *An. melanoon*, *An. persiensis*, *An. pseudopictus*. Moreover, Dinparast et al. (2007) cited the species of *An. atroparvovus* and *An. persiensis*. Dinparast et al. (2009) stressed the presence of the species of *An. hyrchanus*. Nikookar et al. (2010) reported the species of *An. plumbeus*, *Oc. echinus*, *Oc. geniculatus*, *Oc. pulcritarsis*, *Cx. pipiens* and *Cs. annulata*.

Iran is in the malaria pre elimination stage (WHO 2008), thus the bionomic study of mosquitoes in this country is very important. There are scattered studies on the fauna and ecological properties of mosquitoes in Mazandaran Province. Many travelers may travel to Mazandaran Province and Neka County during visiting Iran. This part of Iran holds specific location due to access to the Caspian Sea and to other aspects such as trade, commerce and navigation.

This study aimed to determine fauna and the ecology of mosquitoes in Neka County, Mazandaran Province, northern Iran from

April to December, 2009. The data obtain in this study will be valuable to develop programs for future planning of mosquito control in this area.

Materials and Methods

The study area

Mazandaran Province is located in the northern Iran between the Caspian Sea and the Alborz Mountains (36°39' N, 53°17' E) (Fig. 1). This area located in the north with Caspian Sea, in the south with Tehran and Semnan Provinces, in the east with Golestan Province and in the west Guilan Province, including 19 Counties (Geography of Mazandaran Province 2013). Maximum and minimum temperatures were 27.6 and 9.1 °C in July and December 2009, respectively. The maximum annual rainfall has been reported 204.6 mm in October, and the least was 0.1 mm in July 2009. The most important crop of the region is rice, and Hyrcanian forest area and mild and rainy weather caused this Province to be a suitable mosquito habitat (Mazandaran Local Meteorological Office 2009).

Three fixed villages, namely Darvishan, Chalmardi, and Khairabad, were randomly selected for field survey. Moreover, six variable villages (Golbestan, Elmiyeh, Baghearab, Nimchah, Chalehpol, and Khorshid) were selected with regard to existing facilities in Neka County. In each fixed village, six fixed and two variable locations were selected in plain, slope and mountain conditions. Larvae sampling was collected from the natural larval habitats of fixed villages including river margins, ponds and water craters, craters rain water, marsh, tree holes, rice fields, worn tires, irrigation canals, wells, and sewage, using the standard dipper rate of 350 mm capacity every 15 days (WHO 1963). In addition, larval collection carried out in variable villages randomly.

We attempted to collect the data about the biological and physical characteristics of the larval habitats, including running water, residents (permanent or temporary), the type of larval habitats natural or artificial, presence or absence of the plant, type of substrate (sand, clay, or rocky), to be sunny (sunny or shady), water status (clear or opaque) and water temperature.

The larval stages III and IV were preserved in 100 % lactophenol solution and later identified morphologically using the keys of Shahgudian (1960), Zaim and Cranston (1986a), Harbach (1985, 1988), and Azari-Hamidian and Harbach (2009). The results were analyzed by SPSS 11.5 software. The abbreviations of the genus names were regulated based on Reinert (2009).

Results

A total of 32530 larvae were collected: 3625 larvae of anophelinae and 28905 culicinae. Table 1 shows the results of anophelinae and culicinae species in the fixed and variable breeding places and tree holes. Four species of *Anopheles* were collected: *An. claviger* (0.31%), *An. maculipennis* (0.54%), *An. plumbeus* (10.28%), and *An. superpictus* (0.01%).

Anopheles claviger collected 0.31 %, 2.78 % of total culicidae and *Anopheles* collection, respectively (Table 1). The characteristics of the habitats under different temperature and altitude conditions are shown in Table 2. The minimum and maximum temperature of water of the larval habitats was calculated as 19 °C and 22 °C, respectively (average 20.5 °C). On the other hand, the maximum and minimum altitude of the habitats was 300 and 200 meters, respectively (average 250 meters).

Anopheles maculipennis was collected 0.54 %, 4.82 % of the total and *Anopheles* collection, respectively. Maximum and mini-

imum water temperature was reported as 26 °C and 17 °C, respectively (average 22.5 °C). Maximum and minimum altitude was 300 and 180 meters respectively, (average 240 meters).

Anopheles plumbeus was collected 10.28 %, 92.27 % of total and *Anopheles* collection, respectively. Moreover, this species was collected in tree holes and the artificial habitats such as pond water. Maximum and minimum temperature was found as 16 °C and 14 °C, respectively (average 15 °C). The minimum and maximum sea level of larval habitat was calculated at least 180 and 250 meters, respectively and the average were 215 meters.

Anopheles superpictus was collected 0.01 %, 0.13 % of the total and *Anopheles* collection, respectively. Minimum and maximum temperature of the larval habitat was reported as 17 °C and 19 °C, respectively and average was 180 °C. The minimum altitude was 280 meters, and the maximum was 300 (average 290 meters). The optimum of temperature and favorable altitudes of other species are presented in Tables 1 and 2.

The majority of *An. plumbeus* was collected in natural, permanent, and stagnant water with vegetation, turbid and transparent water and semi-shade tree holes are shown in Table 3. *An. superpictus* larvae were collected from natural, permanent, stagnant water with vegetation, clear and transparent breeding places. On the other hand, *An. claviger* larvae were collected mostly in permanent, and stagnant, transparent water with semi-shade and vegetation. *Culex mimeticus* was found in the natural habitats such as river and ponds from Permanent River, and stagnant pools with vegetation. The larvae of *Cx. pipiens*, the most abundant species, were found in artificial habitats and sewage breeding places in agricultural lands and low abundance along the river. Moreover, *Oc. geniculatus*, the most dominant species, was found in permanent water with vegetation in slow running with clay substrate water, semi-

shade in natural breeding places. The larvae of *Cs. longiareolata* were found in most the larval habitats. The larval habitat characteristics of the other species are shown in Table 3.

Checklist of Culicidae in Mazandaran Province

Family culicidae

Subfamily Anophelinae

Subgenus *Anopheles* Meigen

- 1- *An. (Ano.) algeriensis*
- 2- *An. (Ano.) claviger*
- 3- *An. (Ano.) marteri*
- 4- *An. (Ano.) maculipennis*
- 5- *An. (Ano.) atroparvus*
- 6- *An. (Ano.) melanoon*
- 7- *An. (Ano.) messeae*
- 8- *An. (Ano.) persiensis*
- 9- *An. (Ano.) sacharovi*
- 10- *An. (Ano.) plumbeus*
- 11- *An. (Ano.) hyrcanus*
- 12- *An. (Ano.) pseudopictus*

Subfamily Culicinae

Subgenus *Cellia* Theobald

- 13- *An. (Cel.) pulcherrimus*
- 14- *An. (Cel.) superpictus*

Subgenus *Aedimorphus* Theobald

- 15- *Ae. (Adm.) vexans*

Subgenus *Finlaya*

- 16- *Oc. (Fin.) echinus*
- 17- *Oc. (Fin.) geniculatus*

Subgenus *Ochlerotatus*

- 18- *Oc. (Och.) caspius*
- 19- *Oc. (Och.) pulcritarsis*

Subgenus *Culex*

- 20- *Cx. (Cux.) pipiens*
- 21- *Cx. (Cux.) perexiguus*
- 22- *Cx. (Cux.) territans*
- 23- *Cx. (Cux.) theileri*
- 24- *Cx. (Cux.) mimeticus*
- 25- *Cx. (Cux.) tritaeniorhynchus*
- 26- *Cx. (Cux.) hortensis*

Subgenus *Allotheobaldia*

- 27- *Cs. (All.) longiareolata*

Subgenus *Culiseta*

- 28- *Cs. (Cus.) annulata*
- 29- *Cs. (Cus.) subochrea*

Subgenus *Coquillettidia*

- 30- *Cq. (Coq.) richiardi*

Subgenus *Pseudoficalbia*

- 31- *Ur. (Pfc.) unguiculata*




Fig. 1. Map of Iran and Neka County, northern Iran

Table 1. Abundance of larval mosquitoes (density per 10 dip) in Neka County, Mazandaran Province, northern Iran, 2009

Species	Villages				n	%
	Darvishan	Chalmardi	Khairabad	Variable		
<i>An. claviger</i>	2	6	2	0	100	0.31
<i>An. maculipennis</i>	10	4	3.5	0	175	0.54
<i>An. plumbeus</i>	1.5	0	333	0	3345	10.28
<i>An. superpictus</i>	0	0.5	0	0	5	0.01
<i>Cs. annulata</i>	2	25	8	0	350	1.07
<i>Cs. longiareolata</i>	50.5	21.5	202	16	2900	8.91
<i>Cx. mimeticus</i>	0	0.5	0.5	0	10	0.03
<i>Cx. pipiens</i>	527.5	494	991	69	20815	63.99
<i>Oc. geniculatus</i>	0	0	483	0	4830	14.85

Table 2. Species composition of mosquito larvae collected from different temperatures and altitudes in Neka County, northern Iran, 2009

Species	Temperature C			Altitude		
	Minimum	Maximum	Mean	Minimum	Maximum	Mean
<i>An. claviger</i>	19	22	20.5	200	300	250
<i>An. maculipennis</i>	17	26	22.5	180	300	240
<i>An. plumbeus</i>	14	16	15	180	250	215
<i>An. superpictus</i>	17	19	18	280	300	290
<i>Cs. annulata</i>	12	22	17	180	300	240
<i>Cs. longiareolata</i>	12	25	18.5	180	300	240
<i>Cx. mimeticus</i>	12	13	12.5	200	300	250
<i>Cx. pipiens</i>	12	25	18.5	180	300	240
<i>Oc. geniculatus</i>	11	14	12.5	230	250	240

Table 3. Larval habitat characteristics in Neka County, Mazandaran Province, northern Iran, 2009

	<i>An. claviger</i>	<i>An. maculipennis</i>	<i>An. plumbeus</i>	<i>An. superpictus</i>	<i>Cs. annulata</i>	<i>Cs. longiareolata</i>	<i>Cx. mimeticus</i>	<i>Cx. pipiens</i>	<i>Oc. geniculatus</i>
Habitat status									
Permanent	91	95.3	91.3	66.2	82.5	75.1	100	71.6	99
Temporary	9	4.7	8.7	33.8	17.5	24.9	0	28.4	1
Water									
Slow running	10	16	20	29.5	10	2	3	30	0
Stagnant	90	84	80	70.5	90	98	87	70	100
Vegetation									
Without vegetation	2	2	7.6	20	5	29	19.6	13	0
With vegetation	98	98	92.4	80	95	71	80.4	87	100
Type of bed									
Clay	47	1	84.5	50	98	88	66	56	94
Sand	3	17	15.5	20	2	5	30	33	6
Stone or cement	50	82	0	30	0	5	4	11	0

Water status									
Opaque	1	40	50	9.1	50	40	40	45	50
Transparent	99	60	50	90.9	50	60	60	55	50
Status Light									
Sunny	1	33.2	0	14.7	0	2.26	26.7	37.4	0
Shaded	34	28.4	5	26.1	5	26.9	19.4	18.9	5
Semi-shade	65	38.4	95	59.2	95	70.5	54.9	43.7	95
Habitat type									
Natural	99	55	98	89	98	97	70	41	100
Artificial	1	45	2	11	2	3	30	59	0

Discussion

In the present study, *An. plumbeus* larvae were collected from tree holes and artificial habitats such as permanent and stagnant water. However, the maximum and minimum of temperature of the larval habitat was as 16 °C and 14 °C, respectively. In parallel, Azari-Hamidian (2011a) found this species in tree holes and discarded tire. The most favorable temperature is 15–18 °C; however, the species has tolerated to freeze (up to -6° C to -8 °C) in the laboratory (Horsfall 1955). However, Garrett Jones reported the occurrence of this species in Kelardasht of Mazandaran Province (Garrett Jones 1951).

In our investigation, *Cs. annulata* was found in natural, permanent, stagnant, semi-shade habitats with vegetation. This species has been reported in Guilan Province, Alborz and Zagros Mountains, East Bushehr and Robat Tork of Isfahan Province (Azari-Hamidian et al. 2003b). Zaim and Cranston (1986) included *Cs. subochrea* in their systematic keys and *Cs. annulata* in their checklist. The taxonomy and distribution of *Cs. annulata*, *Cs. subochrea* and *Cs. alaskaensis* need to be more investigated in Iran (Azari-Hamidian 2005). Record of *Cs. annulata* in our research was based on adults, whereas *Cs. subochrea* were not found in our collection, and neither were the adults of this species (Table1). *Culiseta subochrea* and *Cs. annulata* were reported in Ardebil Province (Azari-Hamidian et al. 2009).

The larval habitats of *Cs. annulata* have also been reported from various places except tree holes (Azari-Hamidian et al. 2003b). However, *An. plumbeus* and *Oc. geniculatus* species have been reported in tree holes (Nikookar et al. 2010).

In our study, the larvae of *Cs. longiareolata* were collected from all larval habitats except tree holes, which comprised 8.91 %, 10.03 % of the total and Culicinae collection, respectively. This species was reported first time in north of the country by Gutzevich (1943). The occurrence of this species was also reported by the previous study in Mazandaran Province (Zaim 1987). The larval habitat of this species has been reported from many parts of Iran (Zaim 1987). At the same study, the larval habitat has been reported as contaminated with organic materials, in old pit located in outdoor places and rice fields (Kitron and Pender 1986, Mousakazemi et al. 2000, Azari-Hamidian 2005). It has also been reported from the North Khorasan and Qom Provinces (Azari-Hamidian et al. 2011, Saghafipour et al. 2012).

There is scatter information about the distribution and taxonomy of the three species of *Cs. alaskaensis*, *Cs. annulata*, and *Cs. subochrea* in Iran (Zaim et al. 1986, Azari-Hamidian et al. 2003b, Azari-Hamidian 2005). Moosa Kazemi et al. (2010) found the species of *Cs. subochrea* and *Cs. longiareolata*

in Sanandaj County of Kurdistan Province, northwestern Iran.

In our research, *An. claviger* was found in permanent, stagnant water with vegetation habitat, as well as in the clay and stone substrate of water. In parallel, Macan (1950) has reported the existence of this species in the sunny/shadowy springs, and slow running pools water temperature of 14–16 °C in Iraq and western Iran. Dow (1953) reported the larval habitat of this species in shallow and small pools with little vegetation. It was also found within the larval habitats with temperature of 12–15 °C (Horsfall 1955). The occurrence of this species in Mazandaran was reported previously by Garrett Jones (Garrett Jones 1951). Zaim (1987) collected *An. claviger* larvae from Kashan, central Iran.

In our study, *An. maculipennis* was found in permanent, transparent, semi-shady natural breeding places with vegetation habitat, as well as in the cement or stone substrate of water. In parallel, Azari-Hamidian (2011) reported the presence of this species in the stagnant, gravel substrate, sunny springs, and pools with stagnant water. *Anopheles maculipennis* complex formally comprises 12 Palearctic members, including *An. atroparvus*, *An. beklemishevi*, *An. labranchiae*, *An. maculipennis*, *An. martinius*, *An. melanoon*, *An. messeae*, *An. sacharovi*, *An. persiensis*, *An. daciae*, *An. lewisi* and *An. artemievi* (Dinparast Djadid et al. 2007). *Anopheles maculipennis* species associated by *An. hyrcanus*, *An. claviger* from Mazandaran Province previously (Nikookar et al. 2010). This species has been identified in Guilan (Azari-Hamidian et al. 2004), Mazandaran and Golestan Provinces (Zaim et al. 1986). *Anopheles messeae* and *Anopheles melanoon* reported from Guilan Province, and *An. sacharovi* and *An. hyrcanus* from Golestan Province (Saebi 1987). *Anopheles sacharovi* has been reported from Mazandaran Province (Sedaghat et al. 2003). Dow reported the species of *An. subalpinus* in Sari, Babolsar,

Chalus and Astaneh cities in Mazandaran Province. He also reported *An. melanoon* in Astaneh city, Guilan Province (Dow 1983). *Anopheles subalpinus* is a Synonyms *An. melanoon*. Initial studies mainly focused on identification of the mosquitoes and their role in disease transmission (Gutsevich 1943, Macan 1950, Dow 1953, Minar 1974).

In our investigation, *An. superpictus* was found as 0.01 %, 0.13 % of the total and *Anopheles* collection, respectively. This species was found in permanent, stagnant, with clay substrate, transparent water, semi-shade, natural with vegetation habitat. Studies in different parts of Mazandaran Province confirmed the occurrence of this species (Zolotarev 1945, Dow 1953). Moreover, Azari-Hamidian (2011) stated its presence in stagnant, transient, mud substrate, fully sunlight places with vegetation in natural habitats. Further support for our results comes from a study by Mousakazemi et al. (2000) in Zarrin-Shahr and Mobarakeh areas of Isfahan Province. Some information on the larval stage of *An. superpictus* was reported by Yaghoobi-Ershadi et al. (2001) in Ardebil Province, northwestern Iran. Some ecological aspects of this species have been in Rasht County of Guilan Province (Azari-Hamidian et al. 2002b). Ghanbari et al. (2005) reported the larval breeding places of mosquitoes including anophelines in Iranshahr, southeastern Iran. However, the edges of stony streams with sunlight and semi-shallow have been mentioned as the main breeding places of this species in western Iran (Macan 1950). Three genotypes named X, Y, and Z within *An. superpictus* have been reported in Iran by Oshaghi et al. (2008). There are no reports about the genotypes of this species in Mazandaran Province, however, it needs to be studied by the future investigations.

In our study, *Culex pipiens* was found as 63.99 % of the total and 72.01 % of Culicinae larvae collection in natural and

artificial larval habitats such as tree holes. In addition, hollow pipes, hot water, holes Hungarian sewage, stagnant water, and toilet field were reported as the breeding places of this species (Jupp 1970, Kulanin 1980, Zaim 1987, Vinogradova 2000, Mousakazemi et al. 2000). This species were reported in Natural habitat (rain pool, river edge) with Standing water Azari-Hamidian (2007b).

The species of *Cx. pipiens* and *Cx. quinquefasciatus* are very similar. There are no reports in regarding the existence of *Cx. quinquefasciatus* in Mazandaran Province. The shape of siphon of larvae in the tip, middle and the base has been mentioned to identification of two recent species (Zaim and Cranston 1986). However, Azari-Hamidian and Harbach (2009) mentioned the existence of two branches of seta 1 on abdominal segment III-VI for *Cx. pipiens* and one branch for *Cx. quinquefasciatus*.

The common characteristics among the species of *Cx. vagans*, *Cx. torrentium*, and *Cx. pipiens*, were reported as the presence of two pairs of long seta on the base of siphon (Zaim and Cranston 1986). The existence of seta 1 with 4–5 branches and seta 1-X with two branches is attributed to *Cx. torrentium* and two branches of seta 1 on abdominal segment III-V mentioned to *Cx. vagans* (Harbach 1985).

In our study, *Ochlerotatus geniculatus* was collected as comprising 14.85 %, 16.7 % of the total and Culicinae larvae collections, respectively. In addition, the most favorable temperature was 11–14 °C. Maximum and minimum altitude of the larval habitat was 250 meters and 230 meters, respectively and the average was 240 meters. Further support of this result comes from some previous studies (Gutsevich 1943, Zaim et al. 1984a, Zaim 1987, Azari-Hamidian et al. 2001, Nikookar et al. 2010). This species is very similar to *Oc. echinus* as reported in a recent study (Nikookar et al. 2010). *Ochlerotatus echinus* is distributed in Palaearctic, Europe,

north Africa, and south east Asia, and in Iran, it has been reported from Guilan Province in the north (Knight and Stone 1977, Azarai-Hamidian et al. 2002). In addition, this species were collected by janbakhsh in Sari County and reported by Zaim (1984a). Studies from some parts of Mazandaran Province showed the occurrence of this species (Zaim et al. 1984a, Zaim 1987, Nikookar et al. 2010). Moreover, the distribution of this species has been reported from both Guilan, Mazandaran Provinces (Azari-Hamidian et al. 2002a), as well as from Gulestan (Gutsevich 1943), and Ardebil (Azari-Hamidian et al. 2009).

In our study, *Culex mimeticus* was found as comprising 0.03 % of the total larvae collection in permanent, stagnant, transparent water, clay substrate, semi-shade natural breeding places. In parallel, Zaim (1987) stressed the permanent, slow running water, full sunlight, mud substrate, transient and fresh water with vegetation in natural stream bed pools and rice fields as the breeding places of this species. Studies from different parts of Mazandaran confirmed the occurrence of this species (Dow 1953, Zaim 1987).

Conclusion

Our results indicate that the presence of the dominant species of *Cx. pipiens* in larval habitat natural and artificial *Oc. geniculatus* and *An. plumbeus* in natural such as tree hole. These species dramatically increased and reach to peak in warm season. The recent species could be creating nuisance and health risks in this area.

Acknowledgements

The authors are grateful to Dr Azari-Hamidian for giving the useful comments, Dr Davood Nikzad, Mr Hamid Ravesh, Mr Mojtaba Asghari, Mr Mirramazan Nikookar,

Mr Mehrdad Badangiz, and Mr Mohammad Badangiz (Environmental Health Director of Neka) for supporting the investigation. We also would like to express our appreciation to the people of Darvishan, Khairabad and Chelmardi villages in Neka County for their kind cooperation during the study. This study was financially supported by Tehran University of Medical Sciences (No: 10513-27-01-89). The authors declare that there is no conflict of interests.

References

- Abai MR, Azari-Hamidian S, Ladonni H, Hakimi M, Mashhadi-Esmail K, Sheikhzadeh K, Kousha A, Vatandoost H (2007) Fauna and checklist of mosquitoes (Diptera: Culicidae) of East Azerbaijan, northwestern Iran. *Iran J Arthropod-Borne Dis*.1: 27–33.
- Azari-Hamidian S, Yaghoobi-Ershadi MR, Javadian E (2001) A faunistic study of mosquito (Diptera: Culicidae) fauna in Rasht County (Iran). *Modarres J Med Sci*. 3: 65–70 [Persian].
- Azari-Hamidian SH, Joeafshani MA, Mosslem M, Rassaei MR (2002a) Mosquitoes of the genus *Aedes* (Diptera: Culicidae) in Guilan. *J Med Fac Guilan Univ Med Sci*. 11(43): 29–39 [Persian].
- Azari-Hamidian S, Joeafshani MA, Mosslem M, Rassaei AR (2002b) Taxonomic survey of mosquitoes (Diptera: Culicida) in Guilan Province with reporting of a subgenus new to Iranian mosquito fauna. The 15th Iranian Plant Protection Congress, 7–11 September 2002, Razi University of Kermanshah, Kermanshah, Iran, pp. 319–320 [Persian].
- Azari-Hamidian S, Joeafshani MA, Mosslem M, Rassaei AR (2003a) Adult mosquito habitats and resting-places in Guilan Province (Diptera: Culicidae). *Hakim J*. 6(3): 55–62 [Persian].
- Azari-Hamidian S, Joeafshani MA, Rassaei AR, Mosslem M (2003b) Faunistic studies on the genus *Culiseta* (Diptera: Culicidae) in Guilan Province. *J Kerman Univ Med Sci*. 10: 225–233.
- Azari-Hamidian S, Joeafshani MA, Rassaei AR, Mosslem M, Mousavi-Eivanaki E (2004) Mosquito fauna of the genus *Anopheles* (Diptera: Culicidae) in Guilan Province. *Modarres J Med Sci*. 6: 11–22 [Persian].
- Azari-Hamidian S (2005) Larval habitat characteristics of mosquitoes of the genus *Culiseta* Felt, 1904 (Diptera: Culicidae) in the Caspian Sea littoral, Iran. *Zool Middle East*. 36: 59–66.
- Azari-Hamidian S (2007a) Checklist of Iranian Mosquitoes (Diptera: Culicidae). *J Vect Ecol*. 32 (2): 235–242.
- Azari-Hamidian S (2007b) Larval habitat characteristics of mosquito of the genus *Culex* (Diptera: Culicidae) in Guilan Province, Iran. *Iran J Arthropod-Borne Dis*. 1(1): 9–20.
- Azari Hamidian S, Yaghoobi-Ershadi MR, Javadian E, Mobedi I, Abai MR (2007) Review of dirofilariasis in Iran. *J Med Fac Guilan Univ Med Sci*. 15(60): 102–113.
- Azari-Hamidian S, Harbach RE (2009) Keys to the adult females and fourth instar larvae of mosquitoes of Iran. *Zootaxa*. 2078: 1–33.
- Azari-Hamidian SH, Yaghoobi-Ershadi MR, Javadian E, Abai MR, Mobedi I, Linton Y-M, Harbach RE (2009) Distribution and ecology of mosquitoes in a focus of dirofilariasis in northwestern Iran, with the first finding of filarial larvae in naturally infected local mosquitoes. *Med Vet Entomol*. 23: 111–121.
- Azari-Hamidian S (2011) Larval habitat characteristics of the genus *Anopheles* (Diptera: culicidae) and a checklist of mosquitoes in Guilan Province, northern Iran. *Iran J Arthropod-Borne Dis*. 5: 37–53.

- Azari-Hamidian S, Abai MR, Arzamani K, Bakhshi H, Karami H, Ladonni H, Harbach RE (2011) Mosquitoes (Diptera: Culicidae) of North Khorasan Province, northeastern Iran and the zoogeographic affinities of the Iranian and Middle Asian mosquito fauna. *J Entomol.* 8(3): 204–217.
- Chinikar S, Ghiasi SM, Moradi A, Madihi SR (2010) Laboratory detection facility of Dengue Fever (DF) in Iran: The first imported case. *Internet J Infect Dis.* 8 (1): 1–2.
- Chinikar S, Ghiasi SM, Shah-Hosseini N, Mostafavi E, Moradi M, Khakifirouz S, Rasi Varai FS, Rafeigh M, Jalali T, Goya MM, Shirzadi MR, Zainali M, Fooks A R (2013) Preliminary study of dengue virus infection in Iran. *Travel Med Infect Dis.* 11(3): 166–169.
- Dehghan H, Sadraeia J, Moosa-Kazemi SH (2011) The morphological variations of *Culex pipiens* (Diptera: Culicidae) in central Iran. *Asian Pacific J Trop Med.* 4(3): 215–219.
- Dinparast Djadid N, Gholizadeh S, Tafsiri E, Romi R, Gordeev M, Zakeri S (2007) Molecular identification of Palearctic members of *Anopheles maculipennis* in northern Iran. *Malar J.* 6: 6.
- Dinparast Djadid N, Jazayeri H, Gholizadeh S, Pashaei Rad SH, Zakeri S (2009) First record of a new member of *Anopheles hyrcanus* group from Iran: Molecular identification, diagnosis, phylogeny, status of kdr resistance and *Plasmodium* infection. *J Med Entomol.* 46(5): 1084–1093.
- Dow RP (1953) Notes on Iranian mosquitoes. *Am J Trop Med Hyg.* 2: 683–693.
- Eslami A (1997) Veterinary Helminthology. Vol. 3. Nematoda and Acanthocephala. Tehran University Publications, Tehran [Persian].
- Edrissian GH (2006) Malaria in Iran: Past and present situation. *Iran J Parasitol.* 1: 1–14.
- Eshghy N (1977) *Anopheles multicolor* Cambouliu and its role in the transmission of malaria in Iran. *J Entomol Soc Iran.* 4:87–88 [Persian].
- Faghih MA (1969) Malariology and Malaria Eradication. Tehran University Press. Tehran [Persian].
- Garret J (1951) An experiment in trapping and controlling *Anopheles maculipennis* in north Iran an experiment in north Iran. *Bull World Health Organ.* 4(4): 547–562.
- Ghaffary EN (1956) *Culex* in Iran. School of Public Health, Tehran University of Medical Sciences, Iran.
- Geography of Mazandaran Province (2013) Department of Culture of Mazandaran, Iran. Available at: <http://mazandaran.farhang.gov.ir>.
- Glick JI (1992) Illustrated key to the female *Anopheles* of southwestern Asia and Egypt (Diptera: Culicidae). *Mosq Syst.* 24: 125–153.
- Gutsevich AV (1943) On the mosquitoes of North Iran. *Compt Rend Acad Sci USSR.* 40(3): 123–125 [Russian].
- Harbach RE (1985) Pictorial keys to the genera of mosquitoes, subgenera of *Culex* and the species of *Culex* (*Culex*) occurring in southwestern Asia and Egypt, with a note on the subgeneric placement of *Culex deserticola* (Diptera: Culicidae). *Mosq Syst.* 17: 83–107.
- Harbach RE (1988) The mosquitoes of the subgenus *Culex* in southwestern Asia and Egypt (Diptera: Culicidae). *Contrib Am Ent Inst.* 24: 1–240.
- Harbach RE (2007) Mosquito Taxonomic Inventory. Available at: <http://mosquito-taxonomic-inventory.info/>.
- Horsfall WR (1955) Mosquitoes. Their Biology and Relation to Disease. Hafner Publishing, New York.
- Institute of Parasitology and Malariology (1953) Distribution of the anopheline of Iran. Publication No. 225, Tehran [Persian].

- Jupp PG (1970) The taxonomic status of *Culex* (*Culex*) *univittatus* Theobald (Diptera: Culicidae) in South Africa. *Mosq Syst News*. 2(1): 9–16.
- Karami M (2000) Studied species of fauna and seasonal variations of Exophagic Culicidae in Amol County. [M Sc Thesis]. School of Public Health, Tehran University of Medical Sciences, Iran.
- Kitron U, Pener H (1986) Distribution of mosquitoes (Diptera: Culicidae) in Northern Israel: A historical perspective. II. Culicine mosquitoes. *J Med Entomol*. 23(2): 182–187.
- Knight KL, Stone A (1977) A catalog of the mosquitoes of the world, Thomas Say Foundation. Entomological Society of America. Maryland.
- Kulanin VL (1980) Breeding of *Culex pipiens molestus* Forsk and *Culex pipiens pipiens* L. mosquitoes in Samarkand basements. *Med Parazitol (Mosk)*. 49(1): 25–28.
- Lotfi MD (1970) Iranian species of genus *Culex* (Culicinae: Diptera). *Bull Soc Pathol. Exot*. 63: 400–403.
- Lotfi M (1973) Iranian species of genus *Culex* (Culicinae: Diptera). II. Report of four species of larvae (including three new records) and 14 adult species. *Bull Soc Pathol Exot*. 66(1): 204–207.
- Lotfi MD (1976) Key to Culicinae of Iran, genus *Culex* and their biology (Diptera: Culicidae). *Iran J Public Health*. 5: 71–78.
- Macan TT (1950) *Anopheles* and malaria in the Near East III-the Anopheline mosquitoes of Iraq and north Persia, London. *Sch Hyg Trop Med Res Mem*. No 7, pp. 109–219.
- Mazandaran Local Meteorological Office (2009) Mazandaran Meteorological Office. Mazandaran, Iran. Available at: <http://www.mazandaranmet.ir>.
- Minar J (1974) Results of Czechoslovak-Iranian entomological expedition to Iran, Diptera: Culicidae, Acta Entomol Mus Nat. Pragae. 6: 87–89.
- Minar J (1981) Results of Czechoslovak-Iranian Entomological Expeditions to Iran, Diptera: Culicidae, Oestridae. *Acta Ent Musei Nat Pragae*. 40: 83–84.
- Momeni S, Manouchehri AV, Zaim M, Mottaghi M (1992) The distinguish between the species of *Anopheles maculipennis* complex by using of egg characters in Guilan and Mazandaran provinces, northern areas of Iran (1990–1991). *Iranian Cong Malaria, Zahedan*, pp. 29–30.
- Mousakazemi SH, Zaim M, Zahraini Ramazani AR (2000) Found and ecology of Culicidae mosquitoes in Zarrin-Shahr and Mobarakeh areas, Isfahan Province. *Armagan Danesh J Yasuj Uni Med Sci*. 5: 17–18.
- Moosa-Kazemi SH, Zaim M, Zahraini Ramazani AR (2005) Fauna and ecology of Culicidae mosquitoes of Zarrin-Shahr and Mobarakeh, Isfahan Province, Center of Iran, Medicine and Health in the Tropics Congress, Marseille, France.
- Moosa-Kazemi SH, Vatandoost H, Nikookar H, Fathian M (2009) Culicinae (Diptera: Culicidae) mosquitoes in Chabahar County, Sistan and Baluchistan Province, southeastern Iran. *Iran J Arthropod-Borne Dis*. 3(1): 29–35.
- Moosa-Kazemi SH, Karimian F, Davari B (2010) Culicinae mosquitoes in Sanandaj County, Kurdistan Province, western Iran. *J Vector Borne Dis*. 47: 103–107.
- Naficy K, Saidi S (1970) Serological survey on viral antibodies in Iran. *Trop Geogr Med*. 2: 183–188.
- Nikookar SH, Moosa-Kazemi SH, Oshaghi MA, Yaghoobi-Ershadi MR, Vatandoost H, Kianinasab A (2010) Species composition and diversity of mosquitoes in Neka County, Mazandaran Province, northern Iran. *Iran J Arthropod-Borne Dis*. 4(2): 26–34.

- Oryan A, Valinezhad A, Bahrami S (2008) Prevalence and pathology of camel filariasis in Iran. *Parasitol Res.* 103: 1125–1131.
- Oshaghi MA, Yaghoobi-Ershadi MR, Shemshad K, Pedram M, Amani H (2008) The *Anopheles superpictus* complex: Introduction of a new malaria vector complex in Iran. *Bull Soc Pathol Exot.* 101(5): 429–434.
- Reinert JF (2009) List of abbreviations for currently valid generic-level taxa in family Culicidae (Diptera) *Eur Mosq Bull.* 27: 68–76.
- Saidi S, Tesh R, Javadian E, Nadim A (1976) The prevalence of human infection of West Nile in Iran. *Iran J Public Health.* 5: 8–14.
- Saebi ME (1987) Morphological study on anopheline larvae and their distribution in Iran. [PhD Thesis]. School of Public Health, Tehran University of Medical Sciences, Iran. [Persian].
- Saghafipour A, Abai MR, Farzinnia B, Nafar R, Ladonni H, Azari-Hamidian S (2012) Mosquito (Diptera: Culicidae) fauna of Qom Province, Iran. *Iran J Arthropod-Borne Dis.* 6: 54–61.
- Sedaghat M, Linton M, Oshaghi MA, Vatandoost H, Harbach RE (2003) The *Anopheles maculipennis* complex (Diptera: Culicidae) in Iran: molecular characterization and recognition of a new species. *Bull. Entomol. Res.* 93: 527–535.
- Sedaghat MM, Harbach RE (2005) An annotated checklist of the *Anopheles* mosquitoes (Diptera: Culicidae) in Iran. *J Vect Ecol.* 30(2): 272–276.
- Service MW (1993) Mosquitoes (Diptera: Culicidae) In: Lane RP, Crosskey RW (Eds) *Medical Insects and Arachnids*. Chapman and Hall, London, pp. 120–240.
- Shahgudian ER (1960) A key to anophelines of Iran. *Acta Med Iran.* 3(3): 38–48.
- Siavashi MR, Massoud J (1995) Human cutaneous dirofilariasis in Iran: A report of two cases. *Iran J Med Sci.* 20: 85–86.
- Vatandoost H, Hanafi-Bojd AA (2008) Laboratory evaluation of 3 repellents against *Anopheles stephensi* in the Islamic Republic of Iran. *East Mediterr Health J.* 14: 260–7.
- Vatandoost H, Khazani A, Kebriain-Zadeh A, Rafinejad J, Khoobdel M, Abai MR, Hanafi-Bojd AA, Akhavan AA, Abtahi SM, Rafi F (2008) Comparative efficacy of Neem and dimethyl phthalate (DMP) against malaria vector, *Anopheles stephensi* (Diptera: Culicidae). *Asian Pacific J Trop Med.* 1: 1–6.
- Vatandoost H, Ramin E, Rassi Y, Abai MR (2009a) Stability and wash resistance of local made mosquito bednets and detergents treated with pyrethroids against *Anopheles stephensi*. *Iran J Arthropod-Borne Dis.* 3: 19–28.
- Vatandoost H, Abai MR, Abbasi M, Shaeghi M, Abtahi M, Rafie F (2009b) Designing of a laboratory model for evaluation of the residual effects of deltamethrin (K-othrine WP 5%) on different surfaces against malaria vector, *Anopheles stephensi* (Diptera: Culicidae). *J Vector Borne Dis.* 46: 261–267.
- Vinogradova EB (2000) *Culex pipiens pipiens* mosquitoes: taxonomy, distribution, ecology, physiology, genetic, applied importance and control. vol. 2. Pensoft Publishers. Bulgaria.
- WHO (1963) *Practical entomology in malaria eradication. Part I: Field and laboratory techniques.*
- WHO (2008) *World Malaria Report 2008.* Geneva, Switzerland. Available at: <http://malaria.who.int/wmr2008/malaria2008.pdf>.
- Yaghoobi-Ershadi MR, Namazi J, Piazak N (2001) Bionomics of *Anopheles sacharovi* in Ardebil Province, north

- western Iran during a larval control program. *Acta Trop.* 78(3): 207–215.
- Zaim M, Manouchehri AV, Yaghoobi-Ershadi MR (1984a) Mosquito fauna of Iran. 1- *Aedes*. *Iran J Public Health.* 13(1–4): 22–33.
- Zaim M, Manouchehri AV, Yaghoobi-Ershadi MR (1984b) Mosquito fauna of Iran. 2- *Culex*. *Iran J Public Health.* 13(1–4): 22–33.
- Zaim M, Manouchehri AV, Yaghoobi-Ershadi MR (1985) Mosquito fauna of Iran. 3- *Culexeta*, *Coquillettidia* and *Uranotaenia*. *Iran J Public Health.* 14(1–4): 1–12.
- Zaim M, Cranston PS (1986) Checklist and keys to the Culicinae of Iran (Diptera: Culicidae). *Mosq Syst.* 18: 233–245.
- Zaim M, Manouchehri AV, Yaghoobi-Ershadi MR (1986) Mosquito fauna of Iran (Diptera: Culicidae) other Culicinae. *Iran J Public Health.* 15(1–4): 1–10.
- Zaim M (1987) The distribution and larval habitat characteristics of Iranian Culicinae. *J Am Mosq Control.* 3(4): 568–573.
- Zaim M, Subbarao SK, Manouchehri AV, Cochrane AH (1993) Role of *Anopheles culicifacies s.l.* and *An. pulcherrimus* in malaria transmission in Ghassreghand (Baluchistan). *Iran. J Am Mosq Control.* 9: 23–26.
- Zolotarev ER (1945) *Anopheles maculipennis* of northern Iran. *Med Parasitol Parasitic Dis.* 14: 50–57 [Russian].