

[Research]

Assessing the sensitivity of Caspian Kutum (*Rutilus kutum*) and the endangered Caspian trout (*Salmo trutta caspius*) to acute toxicity of nonylphenol

I. Shirdel, M.R. Kalbassi*

Dept. of Aquaculture, Faculty of Marine Sciences, Tarbiat Modares University, Noor, Mazandaran, Iran

* Corresponding author's E-mail: kalbassi_m@modares.ac.ir

(Received: July. 29. 2013, Accepted: Dec. 27. 2013)

ABSTRACT

Toxicity tests are commonly used as a tool to determine the standards of water quality for chemicals and to discover appropriate organisms as bioindicators in toxicological studies, and also could be used as an essential tool for evaluation of the pollutant effects in aquatic ecosystems. The aim of the present study was to evaluate the sensitivity of two Caspian fish species, Caspian trout as an endangered species and Caspian Kutum using the static acute toxicity in response to nonylphenol, which is widely discharged into the Caspian Sea environment. The fish were exposed to various concentrations of nonylphenol for 96 hours to determine the LC₅₀. The experimental nonylphenol concentrations were consisted of five exposure groups for each species: 100, 150, 200, 250, and 300 µg/l for Caspian trout; 400, 800, 1200, 1600 and 2000 µg/l for Caspian Kutum, with a control group for each of them. Using probit analysis, the 96 h LC₅₀ of nonylphenol to Caspian trout and Caspian Kutum was determined to be 204.78 and 1262.36 µg/l, respectively. In addition, Caspian trout was approximately 6 times more sensitive than Caspian Kutum. Nonylphenol was reflected to be "highly toxic" to Caspian trout and "moderately toxic" to Caspian Kutum. The results could be considered in preparing plans for conservation and restocking management of Caspian Kutum and the endangered Caspian trout.

Keywords: Nonylphenol, Toxicity test, Caspian kutum, Caspian trout

INTRODUCTION

Widespread use of industrial, household and agricultural chemicals potentially poses a hazard to threatened and endangered species for the reason that the distribution of endangered species is limited and further adverse effects on their populations could lead to more extinction. Conservation of endangered species from chemical hazards is principally based on standardized toxicity tests employing standard test organisms as surrogates for other species. Nevertheless, the sensitivity of threatened and endangered species populations may not be represented by surrogate species (Sappington *et al.*, 2001). Therefore, determining the sensitivities of endangered and threatened species, could help us to protect their populations by using standard regulatory approaches (Dwyer *et al.*, 2005).

Toxicity tests are widely being utilized as a tool to determine water quality standards for chemicals and to discover appropriate organisms as bioindicators in toxicological studies, and also it could be used as an essential tool for evaluation of the pollutant effects in aquatic ecosystems. The median lethal concentration (96 h LC₅₀) tests carried out to determine the vulnerability and survival potential of organisms to particular toxicant (Boudou and Ribeyre, 1997). LC₅₀ values are indicative of the sensitivity and susceptibility of an organism to a particular toxicant. A species showing lower levels of LC₅₀ in response to a particular toxic material could be considered as a suitable sentinel organism in the field studies concerning that toxicant. Nonylphenol is a toxic microbial degradation product of

nonylphenoethoxylate, a compound which is extensively utilized in the production of non-ionic surfactants, paints, emulsifiers, lubricants, pesticides and detergents (Soares *et al.*, 2008). Nonylphenol is more stable and persistent in the aquatic environment than its parent compound (Soares *et al.*, 2008). Nonylphenol and its ethoxylates are acutely toxic to various organisms, including fish, mollusks, crustaceans and amphibians (Servos, 1999). Widespread use of afore-mentioned products, have been led to release of nonylphenol into the aquatic ecosystems through industrial, agricultural and municipal effluents (Mao *et al.*, 2012). In the Southern Caspian Sea waters, high concentrations of nonylphenol have been reported with concentrations ranging from 0.05 to 29 $\mu\text{g/g}$ dry weight of surface sediment samples in the vicinity of urban areas (Mortazavi *et al.*, 2012).

Caspian trout (*Salmo trutta caspius*) and Caspian Kutum (*Rutilus kutum*) are brackish water species of the southern Caspian Sea that their population has sharply been declined over the past two decades (Shirangi *et al.*, 2011; Vera *et al.*, 2011; Ghaninejad & Abdulmaleki, 2007). A variety of factors, especially pollution, are contributing to decline and extinction of these fish population (Kiabi *et al.*, 1999). These species are artificially being propagated in freshwater and in order to recruit their stock, released to the entering rivers of the Caspian Sea.

Regarding to the existence of high concentrations of nonylphenol in the southern Caspian Sea environment (Mortazavi *et al.*, 2012), short-term exposure to this substance could threaten the survival of the released fish during the critical period in which they adapt to their vicinity environmental conditions.

To author's knowledge, the sensitivity of the Caspian species in response to acute toxicity of nonylphenol has not been investigated to date. Thus, by simulating the freshwater condition of the rivers, the present study aimed to compare the sensitivity of Caspian trout and Caspian Kutum, the representative species of Salmonidae and Cyprinidae families, with high and lower sensitivity (Naylor, 1995; Dwyer *et al.*, 2005), respectively, against

nonylphenol, which is widely being discharged into the Caspian Sea environment.

MATERIALS AND METHODS

Test water characteristics

Salinity, conductivity, pH and dissolved oxygen were measured by Consort multi-parameter analyzer (Consort C863, Belgium). Total hardness was measured using Palintest Hardicol test by Palintest Photometer (Palintest 8000, England) based on a colorimetric method.

Acute toxicity test

Juvenile Caspian trout and Caspian Kutum (5 ± 2 g) were purchased from a local commercial hatchery and transported to laboratory in winter 2013, and maintained in dechlorinated tap water (Dechlorination using sodium thiosulphate together with aeration at 24 hours before using the water). The fish were acclimated for one week prior to the experiment and fed twice daily at the rate of 2% body weight by commercial feed until 24h prior to the initiation of the test. Over the experimental and acclimation period, fish were maintained in 12:12 (Light:Dark) photoperiod and average temperature of 13 ± 2 °C for Caspian trout and 20 ± 2 °C for Caspian Kutum. Before the experiment, a stock solution of nonylphenol (Sigma-Aldrich, PESTANAL, analytical standard, technical mixture, 85%, linear formula: $\text{CH}_3(\text{CH}_2)_8\text{C}_6\text{H}_4\text{OH}$ was prepared by dissolving in absolute ethanol. Static acute toxicity test in this study was conducted according to the procedure as described in the OECD (Organization for Economic Co-operation and Development) guideline (OECD guideline No. 203, 1992). Preliminary tests were carried out to determine the appropriate range of toxicity. According to these preliminary tests, concentrations of nonylphenol for acute toxicity test were selected for each species. In order to determine the median lethal concentration, fishes (10 fish per concentration for each species) were exposed for 96 h to various nominal concentrations of nonylphenol. The fish were maintained in 50L fiberglass tanks containing aerated water and were not fed during the toxicity test. The experiments were consisted of a control group and five

nonylphenol concentration groups (100, 150, 200, 250 and 300 $\mu\text{g/l}$) for Caspian trout and five concentrations (400, 800, 1200, 1600 and 2000 $\mu\text{g/l}$) for Caspian Kutum. Animal were exposed to these concentrations for 96 h without feeding and water exchange. The survival of fish in relation to different nonylphenol concentrations were evaluated and counted over the exposure period at 24, 48, 72 and 96 h, then the dead fish were

removed immediately. Median lethal concentrations of nonylphenol for these two species were calculated by Probit analysis using SPSS software (version 17.0).

RESULTS

Test water characteristics

The physical and chemical properties of the test water are presented in Table 1.

Table 1. Physical and chemical properties of the test water

Parameter	Values
pH	7.9-8.2
Salinity	0.4 (ppt)
Dissolved Oxygen	9 ± 0.3 (mg/l)
Conductivity	800 ($\mu\text{S/cm}$)
Total Hardness	102 (mg/l CaCO_3)

Acute toxicity test

There was a concentration-dependent increase and a time-dependent decrease in the mortality rate in each of the tests. For both species, it was observed that most of the mortalities occurred within the first 24h

of exposure. With increasing concentrations of nonylphenol, the mortality of both species was increased (Fig. 1). No mortality was observed in the control group for Caspian trout and Caspian kutum, and also




Fig 1. Mortality of Caspian trout (A) and Caspian Kutum (B) after 96 h exposure to nonylphenol

The 24, 48, 72 and 96 h LC_{50} (with 95% confidence limits) values for each species are shown in Tables 2 and 3. Calculated 96 h LC_{50} values of nonylphenol for Caspian 72h, were 218.78, 214.01 and 209.37 $\mu\text{g/l}$ trout and Caspian Kutum was 204.78 and 1262.36 $\mu\text{g/l}$, respectively. Also, the concentrations of nonylphenol that caused mortality in 50% of the fish within the 24, 48 and 1 to Caspian trout, and 1414.29, 1299.94 and 1262.36 $\mu\text{g/l}$ to Caspian Kutum, respectively.

Results show that, Caspian trout appeared to be approximately six times more sensitive than Caspian Kutum. Moreover, maximum acceptable toxicant concentration (MATC: one tenth of 96 h LC_{50}), no observed effect concentration (NOEC) and the lowest observed effect concentration (LOEC) of nonylphenol were determined as 20.48, 100 and 126.17 $\mu\text{g/l}$ to Caspian trout and 126.24, 400 and 670.08 $\mu\text{g/l}$ to Caspian Kutum, respectively.

Table 2. Lethal concentrations (LC₁₋₉₉) of nonylphenol for Caspian trout
Concentrations (µg/l) (95% of Confidence Limits)

LC	24 h	95% C.L.	48 h	95% C.L.	72 h	95% C.L.	96 h	95% C.L.
LC ₁	134.30		128.64		130.72		126.17	
LC ₁₀	167.20		161.68		161.52		156.83	
LC ₂₀	183.37		178.02		176.57		171.87	
LC ₃₀	195.99		190.81		188.28		183.60	
LC ₄₀	207.45		202.47		198.90		194.26	
LC ₅₀	218.78	193.97-245.93	214.01	188.56-240.84	209.37	185.12-234.11	204.78	179.99-229.11
LC ₆₀	230.72		226.21		220.39		215.88	
LC ₇₀	244.22		240.03		232.83		228.41	
LC ₈₀	261.02		257.28		248.28		244.00	
LC ₉₀	286.26		283.28		271.41		267.41	
LC ₉₉	356.41		356.04		335.36		332.39	

Table 3. Lethal concentrations (LC₁₋₉₉) of nonylphenol for Caspian Kutum
Concentrations (µg/l) (95% of Confidence Limits)

LC	24 h	95% C.L.	48 h	95% C.L.	72 h	95% C.L.	96 h	95% C.L.
LC ₁	942.90		705.23		670.08		670.08	
LC ₁₀	1131.20		928.14		890.55		890.55	
LC ₂₀	1221.34		1041.93		1003.86		1003.86	
LC ₃₀	1290.77		1132.54		1094.41		1094.41	
LC ₄₀	1353.20		1216.19		1178.22		1178.22	
LC ₅₀	1414.29	1234.64-1583.82	1299.94	1107.69-1504.29	1262.36	1066.38-1462.83	1262.36	1066.38-1462.83
LC ₆₀	1478.13		1389.46		1352.50		1352.50	
LC ₇₀	1549.63		1492.08		1456.08		1456.08	
LC ₈₀	1637.71		1621.84		1587.42		1587.42	
LC ₉₀	1768.21		1820.68		1789.40		1789.40	
LC ₉₉	2121.35		2396.16		2378.14		2378.14	

DISCUSSION

Caspian trout population has been disappeared from a large numbers of its historic areas in the southern part of the Caspian Sea, and its wild populations are close to complete extinction (Shirangi *et al.*, 2011; Vera *et al.*, 2011). In addition, Caspian kutum populations have been sharply declined over the past decades (Ghaninejad & Abdulmaleki, 2007). Elevated levels of pollutants, overfishing and destruction of natural spawning areas are considered as the major causes of the population decline for both species (Kiabi *et al.*, 1999). Evaluation of the tolerance of

endangered and threatened species against toxicants could be helpful for protecting their populations (Dwyer *et al.*, 2005). This study was conducted to compare the sensitivity of two Caspian fish species, Caspian trout (as an endangered species) and Caspian Kutum, to nonylphenol using the 96 h LC₅₀ values.

In the present study, calculated 96 h LC₅₀ values of nonylphenol for Caspian trout and Caspian kutum was 204.78 and 1262.36 µg/l, respectively, which were in the ranges observed in the previous studies. Previous studies using different test methods in various organisms showed

that nonylphenol is acutely toxic to fish with 96 h LC₅₀ values ranging from 17 to 3000 µg/l and for most species range from 100 to 500 µg/l (Baldwin *et al.*, 1997; Servos, 1999; Bhattacharya *et al.*, 2008). Winter flounder (*Pleuronectes americanus*) was reported to be the most sensitive fish species in response to nonylphenol acute exposure, with a obtained 96 h LC₅₀ of 17 µg/l (Lussier *et al.*, 2000), whereas the African catfish (*Clarias gariepinus*) was the least sensitive fish, with calculated 96 h LC₅₀ values of 3480 µg/l (Kumaran *et al.*, 2011). A recent study has reported 221, 128 and 209 µg/l as 96 h LC₅₀ values of nonylphenol to rainbow trout alevin (240 mg), juvenile fathead minnow (*Pimephales promelas*) and juvenile bluegill, respectively (Spehar *et al.*, 2010). Kashiwada *et al.* (2002) reported that the adult medaka LC₅₀ for *p*-nonylphenol at 72h was 850 µg/l. A study with rainbow trout and apache trout (*Oncorhynchus apache*) as well as two subspecies of the species *Oncorhynchus clarki*, reported LC₅₀s from 140 to 270 µg/l (Brooke, 1993; Dwyer *et al.*, 1995). Moreover, McLeese *et al.* (1981) reported a 96h LC₅₀ of 190 µg/l for nonylphenol in *Salmo salar*. Lech *et al.* (1996) obtained a median lethal concentration of 193.65 for rainbow trout after 72 h exposure to nonylphenol. According to the USEPA toxicity category, chemicals with LC₅₀ values ranging from 100 to 1000 µg/l have been considered as "highly toxic", and also chemicals with LC₅₀ values within 1000-10000 µg/l as "moderately toxic" (USEPA, 2010). Therefore, our results clearly indicated that nonylphenol is highly toxic to Caspian trout and moderately toxic to Caspian Kutum.

Our results also showed that the median lethal concentration of nonylphenol for Caspian kutum was much higher than nonylphenol LC₅₀ for Caspian trout. These results were consistent with previous studies which showed that salmonid and coldwater species are more sensitive than cyprinid and warmwater species (Naylor, 1995; Sappington *et al.*, 2001; Dwyer *et al.*, 2005). Sappington *et al.* (2001) studied several cold and warm water species to compare their sensitivity in response to acute 4-nonylphenol exposure and reported 96 h LC₅₀ values of 190, 170, 270

and 290 µg/l for rainbow trout, apache trout, fathead minnow and bonytail chub (*Gila elegans*), respectively. Their results showed that coldwater species were more sensitive than warmwater species. Naylor (1995) presented 96 h LC₅₀ values of 300, 310 and 230 µg/l for fathead minnow, sheepshead minnow and rainbow trout, respectively. It is concluded from these values that a salmonid fish is more sensitive than a cyprinid fish against nonylphenol. Dwyer *et al.* (2005) conducted 4-nonylphenol toxicity tests with 18 fish species from seven families including: Cyprinidae, Cyprinodontidae, Salmonidae, Acipenseridae, Catostomidae, Percidae and Poeciliidae. Their results showed that species of the families Acipenseridae and Salmonidae were the most sensitive species with 96 h LC₅₀ values ranging 50-80 and 150-190 µg/l, respectively. Fathead minnow, bonytail chub and Colorado pikeminnow representing Cyprinidae family showed 96 h LC₅₀ values of 270, 290 and 260 µg/l, respectively. Based on our results, it is obvious that Caspian trout and Caspian Kutum exhibited different tolerances in response to nonylphenol exposure. These results clearly demonstrated that much higher concentrations of nonylphenol are required to induce mortality in the 50% of Caspian Kutum than those needed in Caspian trout.

As mentioned above, various species show different sensitivity in response to a given toxicant. Also inter-laboratory variations may be observed in the toxicity of a particular chemical to a particular species (Sappington *et al.*, 2001). The differences in acute toxicity even for a particular toxicant and particular species may be due to the changes in water quality, method for preparation of the toxicant, method of data analysis, fish size, age and condition of the test species as well as the experimental factors, formulation and purity of the test chemicals (Lemke, 1981; Rathore and Khangarot, 2002).

CONCLUSION

According to these acute toxicity tests (96h LC₅₀ values), it is concluded that nonylphenol is highly toxic to Caspian trout. So, it could be introduced as suitable

sentinel fish for assessing the nonylphenol toxicity in the field studies. In addition, LC₅₀ values determined in the present study can be used to determine thenonylphenol chronic toxicity concentrations for Caspian trout and Caspian Kutum. Moreover, these findings may be useful in preparing appropriate regulations for protection and restocking management of these species.

ACKNOWLEDGMENTS

The authors would like to thank their colleagues at the Faculty of Marine Sciences, Tarbiat Modares University, Iran, for their financial support in this project.

REFERENCES

- Baldwin, W.S., Graham, S.E., Shea D. and LeBlanc, G.A. (1997) Metabolic androgenization of female *Daphnia magna* by the xenoestrogen 4-nonylphenol. *Environmental Toxicology and Chemistry*, 16: 1905-1911.
- Bhattacharya, H., Xiao, Q. and Lun, L. (2008) Toxicity studies of nonylphenol on rosy barb (*Puntius conchonius*): A biochemical and histopathological evaluation. *Tissue and Cell*, 40: 243-249.
- Boudou, A. and Ribeyre, F. (1997) Aquatic ecotoxicology: from the ecosystem to the cellular and molecular levels. *Environmental Health Perspectives*, 105: 21-35.
- Brooke, L. (1993) Acute and chronic toxicity of nonylphenol to ten species of aquatic organisms. Report to the US Environmental Protection Agency.
- Dwyer, F., Mayer, F., Sappington, L., Buckler, D., Bridges, C., Greer, I., Hardesty, D., Henke, C., Ingersoll, C. and Kunz, J. (2005) Assessing contaminant sensitivity of endangered and threatened aquatic species: Part I. Acute toxicity of five chemicals. *Archives of Environmental Contamination and Toxicology*, 48: 143-154.
- Dwyer, F., Sappington, L., Buckler, D. and Jones, S. (1995) Use of surrogate species in assessing contaminant risk to endangered and threatened fishes (EPA/600/R-96/029). United States Environmental Protection Agency, Washington, DC.
- Ghaninejad, D. and Abdulmaleki, S. (2007) Annual stocks assessment of bony fish in the Caspian Sea. Iranian Fisheries Research Organization (IFRO), Tehran, Iran 65p.
- Kashiwada, S., Ishikawa, H., Miyamoto, N., Ohnishi, Y. and Magara, Y. (2002) Fish test for endocrine-disruption and estimation of water quality of Japanese rivers. *Water Research*, 36: 2161-2166.
- Kiabi, B.H., Abdoli, A. and Naderi, M. (1999) Status of the fish fauna in the south Caspian Basin of Iran. *Zoology in the Middle East*, 18: 57-65.
- Kumaran, S.S., Kavitha, C., Ramesh, M. and Grummt, T. (2011) Toxicity studies of nonylphenol and octylphenol: hormonal, hematological and biochemical effects in *Clarias gariepinus*. *Journal of Applied Toxicology*, 31: 752-761.
- Lech, J.J., Lewis, S.K. and Ren, L. (1996) In vivo estrogenic activity of nonylphenol in rainbow trout. *Toxicological Sciences*, 30: 229-232.
- Lemke, A.E. (1981) Interlaboratory comparison: Acute testing set: US Environmental Protection Agency, Research and Development, Environmental Research Laboratory.
- Lussier, S.M., Champlin, D., Li, Volsi, J., Poucher, S. and Pruell, R.J. (2000) Acute toxicity of *para*-nonylphenol to saltwater animals. *Environmental Toxicology and Chemistry*, 19: 17-621.
- Mao, Z., Zheng, X-F., Zhang, Y-Q., Tao, X-X., Li Y. and Wang, W. (2012) Occurrence and biodegradation of nonylphenol in the environment. *International Journal of Molecular Sciences*, 13: 491-505.
- McLeese, D.W., Zitko, V., Sergeant, D.B., Burrige, L. and Metcalfe, C.D. (1981) Lethality and accumulation of alkylphenols in aquatic fauna. *Chemosphere*, 10: 723-730.
- Mortazavi, S., Riyahi Bakhtiari, A., Esmaeili Sari, A., Bahramifar, N. and Rahbarizade, F. (2012) Phenolic endocrine disrupting chemicals (EDCs) in Anzali Wetland, Iran: Elevated concentrations of 4-nonylphenol, octylphenol and bisphenol A. *Marine Pollution Bulletin*, 64: 1067-1073.

- Naylor, C.G. (1995) The environmental fate and safety of nonylphenol ethoxylates. *Textile Chemist and Colorist*, 27: 29-33.
- OECD guideline for testing of chemicals, No. 203 (1992) Fish, Acute toxicity test. Adopted by the council on 17th July 1992.
- Sappington, L.C., Mayer, F.L., Dwyer, F.J., Buckler, D.R., Jones, J.R. and Ellersieck, M.R. (2001) Contaminant sensitivity of threatened and endangered fishes compared to standard surrogate species. *Environmental Toxicology and Chemistry*, 20: 2869-2876.
- Servos, M.R. (1999) Review of the aquatic toxicity, estrogenic responses and bioaccumulation of alkylphenols and alkylphenol polyethoxylates. *Water Quality Research Journal of Canada*, 34: 123-177.
- Shirangi, S.A., Kalbassi, M.R. and Dorafshan, S. (2011) Microsatellite polymorphism reveals low genetic differentiation between fall and spring migratory forms of endangered Caspian trout, *Salmo trutta caspius* (Kessler, 1870). *Caspian Journal of Environmental Sciences*, 9: 9-16.
- Soares, A., Guieysse, B., Jefferson, B., Cartmell, E. and Lester, J.N. (2008) Nonylphenol in the environment: A critical review on occurrence, fate, toxicity and treatment in wastewaters. *Environment International*, 34: 1033-1049.
- Spehar, R.L., Brooke, L.T., Markee, T.P. and Kahl, M.D. (2010) Comparative toxicity and bioconcentration of nonylphenol in freshwater organisms. *Environmental Toxicology and Chemistry*, 29 :2104-2111.
- USEPA, (2010) USEPA Dispersed Oil Toxicity Testing. July 31. U.S. Environmental Protection Agency, Office of Research and Development.
- Vera, M., Sourinejad, I., Bouza, C., Vilas, R., Pino-Querido, A., Kalbassi, M.R. and Martínez, P. (2011) Phylogeography, genetic structure, and conservation of the endangered Caspian brown trout, *Salmo trutta caspius* (Kessler, 1877), from Iran. *Hydrobiologia*, 664: 51-67.

Arhive

ارزیابی حساسیت ماهی سفید دریای خزر (*Rutilus frisii kutum*) و ماهی در معرض خطر آزاد دریای خزر (*Salmo trutta caspius*) در برابر سمیت حاد نونیل فنول

۱. شیردل، م. کلباسی*

گروه آبی پروری، دانشکده علوم دریایی دانشگاه تربیت مدرس، نور، مازندران

(تاریخ دریافت: 92/4/7 - تاریخ پذیرش: 92/10/6)

چکیده

آزمون‌های سمیت به طور معمول برای انتخاب گونه‌های مناسب به عنوان شاخص‌های زیستی در مطالعات سم‌شناسی و همچنین جهت ارزیابی اثرات آلاینده‌ها بر اکوسیستم‌های آبی به کار گرفته می‌شوند. در مطالعه حاضر حساسیت دو گونه‌ی ماهی سفید دریای خزر و ماهی آزاد دریای خزر در برابر آلاینده نونیل فنول و با به‌کارگیری آزمون سمیت حاد به صورت ساکن مورد ارزیابی قرار گرفت. به منظور تعیین غلظت نیمه‌کشنده، ماهی‌ها به مدت 96 ساعت در معرض غلظت‌های مختلف نونیل فنول قرار داده شدند. غلظت‌های به کار گرفته شده شامل 100، 150، 200، 250 و 300 میکروگرم در لیتر برای ماهی آزاد و 400، 800، 1200، 1600 و 2000 میکروگرم در لیتر برای ماهی سفید بود. غلظت نیمه‌کشنده (LC₅₀) نونیل فنول در مدت زمان 96 ساعت برای ماهی آزاد و سفید با استفاده از آنالیز Probit محاسبه شد که به ترتیب برابر 204/78 و 1262/36 میکروگرم در لیتر بود. به علاوه مشخص شد که ماهی آزاد در برابر نونیل فنول 6 برابر حساس‌تر از ماهی سفید می‌باشد. بر اساس طبقه‌بندی سمیت USEPA، نونیل فنول برای ماهی آزاد دریای خزر بسیار سمی و برای ماهی سفید، دارای سمیت متوسط می‌باشد. نتایج این تحقیق می‌تواند در تهیه برنامه‌های حفاظتی و بازسازی ذخایر این دو گونه مورد استفاده قرار گیرد.

* مولف مسئول