

[Research]

## Habitat suitability index of *Barbus cyri* (Heckel, 1843) in Tootkabon River, the South Caspian Sea basin, Iran

H. Asadi<sup>1\*</sup>, M. Sattari<sup>1,3</sup>, S. Eagderi<sup>2</sup>

1- Department of Fisheries, Faculty of Natural Resources, University of Guilan, Sowmeh Sara, Guilan, Iran.

2- Department of Fisheries, Faculty of Natural Resources, University of Tehran, Karaj, Iran.

3- Department of Marine Sciences, The Caspian Sea basin Research Center, University of Guilan, Rasht, Iran.

\* Corresponding author's E-mail: asadi.shil@gmail.com

(Received: May. 03, 2015 Accepted: Oct. 21, 2015)

### ABSTRACT

Knowledge of habitat requirements of aquatic animals plays an important role in fisheries and protection programs of aquatic ecosystems. Hence, this study was conducted to investigate the suitability indexes of habitat use and habitat suitability index (HSI) of *Barbus cyri* in its distribution range in Tootkabon River. A number of environmental variables, including elevation, water depth, river width, river slope, velocity, substrate type, and average diameter of bed stone, riparian vegetation type and the relative abundance of *B. cyri* at 13 stations and three replications from the downstream to upstream were examined during November 2013. The results showed that *B. cyri* mostly selects upper stretches of the river with higher velocity, middle depth, lower width and bed rock substrate i.e. bed with boulder cover and grasslands and also its residential area is of riparian type compared with the available ranges. Habitat selectivity index for *B. cyri* showed that the elevation is 130-220 m, water depth 18-75 cm, channel width less than 12 m, channel slope 0.5-2.3%, water velocity less than 0.8 m.s<sup>-1</sup>, bed rock substrate, average diameter of bed stone larger 15-50cm. Presence of these conditions with HSI of 0.798 indicates that Tootkabon River is an excellent habitat for *B. cyri*.

**Key words:** Habitat selection, Suitability index, *Barbus cyri*, Tootkabon River.

### INTRODUCTION

Changes of the hydrological features in a riverine ecosystem by anthropological activities such as habitat destruction can negatively affect survive, reproductive success and growth rate of aquatic organisms particularly fish species (Rosenfeld, 2003, Ahmadi-Nedushan *et al.* 2006) threatening their survival.

Therefore, as first step to assess the impact of human activity on riverine ecosystem and protection of their biodiversity, understanding the habitat requirements of its organism is crucial (Vinagre *et al.* 2006). Hence, Habitat Suitability Indices (HSI) of an aquatic species is one of the primacy themes in river ecology to predict the effect of human manipulation

due to the alternation of the riverine hydrological features (Guay *et al.* 2000). Hydrological characteristics of a river play an important role in fish distribution affecting their metabolism rate, feeding and behavior (Jowetl *et al.* 2007). Habitat selection means that a fish species is more abundant in a specific habitat which is suitable (Rosenfeld 2003). Many environmental factors are important to affect fish in selecting its preferred habitat (Bovee 1982).

Nevertheless, most of the models to study the habitat suitability of an riverine species use the hydrological parameters such as depth, current velocity, river width, type of river bed, river slope, elevation, and river vegetation (Orth 1988) as important factors.

Kura barbel, *Barbus cyri* (De Filippi 1865) is a cyprinid species native to the Caspian Sea basin (Esmaeili *et al.* 2014a, b). This species bears a fusiform body shape, small scales, inferior mouth, and thin lips with two pairs of barbels (Fig. 1). Berg (1948-1949) refers Caspian Sea basin specimens to *Barbus lacerta cyri*, but this subspecies recognized as a valid species by Naseka & Bogutskaya (2009).

Little information is available regarding biology and ecology of *B. cyri*, particularly its habitat use and habitat suitability indexes are unknown. Therefore, this study was conducted to assess the habitat use and habitat suitability indexes of *B. Cyri* with respect to habitat availability of this species in Tootkabon River, a branch of the Sefidrud River in the Southern Caspian Sea basin.



Fig. 1. *Barbus cyri* caught from Tootkabon River.




Fig. 2. Location of Alborz Province, Tootkabon River and sampling points.

Table 1. Explanation and abbreviation for each categorical habitat variable.

Substrate (mm)		Riparian vegetation type	
Bedrock	>4000	BV	Deciduous forest
Boulder	256-4000	BM	Deciduous forest and residential area
Cobble	64-265		
Pebble	16-64		
Gravel	2-16	A	Grasslands or herbs
Sand	<2	BA	Largely no vegetation

## MATERIALS AND METHODS

### Study area

Tootkabon River is a branch of the Sefidrud River in the Southern Caspian Sea basin, originates from the Alborz Mountains. The approximate length of this river is 23 km with an average slope bed of 6.1% (Fig. 2).

### Sampling

For this study, we sampled fish and habitat variables in autumn 2014 during base flow conditions. Thirteen sampling stations were located across the Tootkabon River (Guilan Province, north of Iran) (Fig. 1). The elevation (m) and geographic coordinates were recorded for each station to  $\pm 10$  m using GPS (Global Positioning System; Garmin) according to Torgersen & Close (2004).

The river and the sampling point locations were mapped using ArcGIS 9.3 (Fig. 1).

Fish were sampled in 10-15 m stretches of the river using a backpack electrofishing device (Samus Mp750, 45cm diameter, aluminium ring anode) in the downstream-upstream direction using upstream and downstream stop-nets of 0.2 cm mesh. For sampling, one-removal method with similar catch-per-unit effort strategy was employed (Klaar *et al.* 2004).

Fish specimens were collected from each station during 30 min, anesthetized in clove powder solution ( $1 \text{ g.l}^{-1}$ ), identified according to Coad (2015), counted, photographed, and finally placed in slow - moving water along the river bank to recover and return to the river.

Since no comparable study on microhabitat use of this species was available, therefore, only the specimens larger than 40 mm were selected and counted in each station for further analysis; because the habitat use of fishes in lotic systems can be strongly affected by ontogeny (Copp & Vilizzi 2004; Gillette *et al.* 2006).

### Habitat Data

Since there is limited or no knowledge available on the studied species, the environmental variables were selected

according to the results of other studies conducted on other fishes (Chuang *et al.* 2006; Riffart *et al.* 2009; Tabatabaei *et al.* 2015). The habitat data were measured immediately after sampling (Yu & Lee 2002).

The measured variables included water depth (cm), river width (m), Elevation (m) river slope (%), velocity ( $\text{m.s}^{-1}$ ), substrate type (substrate index), average diameter of bed stone (cm), and riparian vegetation type.

Elevation of sampling sites were recorded by GPS (Garmin).

The mean depth (cm) of each station was estimated by measuring depth at 20 random points across sampling site using a measuring bar, considering their average as the river depth (Lotfi, 2012).

The mean width of river (m) was measured using a tapeline by measuring upper, middle and lower sections of each sampling station, considering their average as the river width. The surface velocity ( $\text{m.s}^{-1}$ ) was estimated by a simple float based on Hassanlie (1999), repeating three times to minimize the error. Dominant substrate type was determined both visually and randomly via measuring the diameter of the riverbed stones in 20 selected quadrat ( $50 \times 50 \text{ cm}$ ) based on Lotfi (2012), and then classified according to Johnston *et al.* (1996) and Tabatabaei *et al.* (2015).

The bed stone diameter average also were calculated by measuring diameter of bed stones in 20 selected quadrat ( $50 \times 50 \text{ cm}$ ) based on Lotfi (2012). Riparian vegetation type (based on the type of vegetation growing at riparian zone of the river or absence thereof), were classified according to our observations, photographs, and standard procedures with some modifications (Johnston *et al.* 1996). The first six variables were continuous, and riparian vegetation type was nominal. The abbreviation and description for each discrete variable are presented in Table 1.

### Data analysis

Habitat use, availability, and selection were calculated over the range of each environmental variable. Each environmental variable was divided into a series of intervals,

and the mean relative abundance of fish in each interval was calculated using habitat selection (Habsel) software 1.0 (Jowett, 2014). The formula  $S = (\%U_{c,i})/(\%A_{c,i})$ , where  $i$  is the interval of a given environmental variable  $c$ ,  $\%U_{c,i}$  is the percentage of utilization of a specific interval of an environmental variable utilized by fish, and  $\%A_{c,i}$  is the percentage of availability of this environmental variable (Guay *et al.* 2000; Waddle 2001; Tabatabaei *et al.* 2015), resulted in a selectivity value ( $S$ ) at each interval. The selectivity values were then scaled, so that the maximum value was one. The mean selectivity values of each environmental variable (scaled to the maximum value of one) was considered as SI value of the variables at each station. Habitat Suitability Index (HIS) was calculated using following formula (Guay *et al.* 2000):

$$HIS = \sqrt[n]{SI\ Width \times SI\ Depth \times \dots \times SI\ n}$$

In this formula, all variables of a habitat are considered essential and the product equation yields zero suitability for any given unsuitable habitat variable.

Since here is no data available regarding the value of each selected variable, therefore, in this research, the SI is multiply to obtain HIS.

## RESULTS

All collected fish during sampling belonged to 8 species viz. *Barbus cyri*, *Capoeta graceless*, *Alburnus filipi*, *Cobitis keyvani*, *Ponticola iranica*, *Ponticola cyrius*, *Alburnoides ssamiii* and *Oxynoemacheilus bergianus* which were returned to the river after identification and counting. A total of 144 specimens of *B. cyri* were collected. Habitat use, availability, and selection are presented in Fig. 3. Table 2 summarizes the average data obtained from each sampling station and environmental variables in the study area are shown.

**Table 2.** Minimum, Maximum, Mean and Standard deviation of variables used.

Variable	Min	Max	Mean	SD
Depth (cm)	15	72	34.71	17.44
Slop (%)	0.37	2.4	1.21	0.68
Width(m)	2.7	5.57	91.31	2.815
Velocity (m.s <sup>-1</sup> )	0.36	1.2	0.51	0.123
Stone D (cm)	4	48.1	20.05	12.65
Elevation	129	208	180.6	25.133

The studied habitats in Tootkabon River mostly occurred in an elevation range of 175-200 m above sea level, with 0.55-0.69 cm depth, 2.8-4.2 m width, 0.60-0.75 m.s<sup>-1</sup> water velocity, slope of 0.5-1.75% and stone diameter of 25-30 cm, and cobble and then boulder substrate type, deciduous riparian forest, and with most available cover type of boulder (Fig. 2 and Table 3). The habitat-use pattern of *B. Cyri* generally followed habitat availability. Considering the availability of environmental variables and the selectivity, the habitat selection pattern of this species mainly had the following features: elevation 130-220 m, water

depth 18-75 cm, channel width less than 12 m, channel slope 0.5-2.3%, water velocity less than 0.8 m.s<sup>-1</sup>, bed rock substrate, average diameter of bed stone 15-50cm, and deciduous forest and residential area riparian type (Fig. 3). Suitability index values of environmental variables for *B. cyri* in Tootkabon River are shown in Table 4. Among of the variables studied, stone diameter had the most (0.81) and elevation (0.57) had the least amount. Calculation showed that the HSI river habitat suitability for *B. cyri* in Tootkabon River is 0.80. The results revealed that *B. cyri* mostly selected upper stretches of the river with


higher velocity and bed rock substrate (Table 3).

Furthermore, this species selects lower depth and velocity, bed rock substrate i.e. bed with larger elements, deciduous forest and

residential area riparian type, and boulder cover compared to the available ranges. In some cases, the pattern of habitat use was different from the pattern of habitat selection i.e. in water depth, river width and velocity.

**Table 3.** Mostly selected variables of *B. cyri*.

Variable	Slop (%)	Velocity(m.s <sup>-1</sup> )	Stone D(cm)	Width(m)	Depth(cm)	Elevation (m)
mostly selected	0.5-1.75	0.60-0.75	25-30	2.8-4.2	0.55-0.69	175-200



**Fig. 3.** Habitat availability (blue line), used (red line) and selected (black line) by *B. cyri* for environmental variables.

**Table 4.** Selected index for *B. cyri* in different stations.

Variable	Slop (%)	Elevation(m)	Velocity(m.s <sup>-1</sup> )	Stone D(cm)	Width(m)	Depth(cm)	HSI <sub>total</sub>
SI	0.803	0.57	0.65	0.81	0.82	0.69	0.798

**Table 5.** Habitat suitability for *B. cyri* in different stations.

Station	1	2	3	4	5	6	7	8	9	10	11	12	13
HSI	0.29	0.65	0.84	9.67	0.63	0.7	0.97	0.81	0.56	0.45	0.41	0.64	0.34

## DISCUSSION

The present study has provided the habitat use and environmental factors affecting the distribution of *B. cyri*, a native fish of Iranian inland waters, in Tootkabon River. In recent years, the industrial effluents in Tootkabon region have been caused the disposal of industrial effluents and chemical pollution of water resources (surface and groundwater) including Tootkabon River. Therefore, the findings of the present study can show the importance of some factors for effective management and protection of this native species. The fish tend to be in the water with a higher rate velocity and rocky bed. However, the preferred habitat of *Barbus* is middle - deep water with high velocity and medium - to large - sized substrates such as rock. The results of this study showed that the water flow rate is an effective factor in the presence and distribution of *B. cyri* in Tootkabon River. In the West Rivers of Taiwan, velocity rate was found to be an important variable influencing the populations of *Sinogastromyzon puliensis* (Shyi-Liang Yu *et al.* 2002) which is in line with our results. Water flow play as an important factor directly on the water and aquatic organisms and indirectly by food supply transfers (Ahmadi-Nedushan *et al.* 2006).

The results revealed that *B. cyri* mostly selects deeper reaches with the bed rock substrate and larger bed stones. In-stream habitat structures provide a variety of functions for stream fishes (Quist *et al.* 2005; Tabatabaei *et al.* 2015); cover features, provide protection from predators or ameliorate adverse conditions of stream flow or seasonal changes in metabolic costs and thereby influence fish survival and movement (MacKenzie & Greenberg 1998; Tabatabaei *et al.* 2015). In addition, deep body shape of this specie can help to rapid turning

and maneuvering in tight quarters as deeper reaches with substrate consisting large bed rocks that provides dead spaces to establish proper habitat. Furthermore, substrate type can be important for fish spawning and feeding behavior (Quist *et al.*, 2005; Tabatabaei *et al.*, 2015). *B. cyri* mostly occupy area with lower river width i.e. less than 12 m. Researches showed that habitats with higher river width have little suitability for fishes such as *Varicorhinus barbatulus* (Littlejohn *et al.* 1985; Chuang *et al.* 2006). In addition, deeper reaches with lower current, less river width along with larger bed stones can provide transparent water to penetrate sunlight causing higher production of periphyton algae as main food items of the *Barbus* species (Treer *et al.* 2006). Habitat of *B. cyri* is an area with deciduous forest and residential area of riparian type. This can be due to providing organic matters that considered as base of the primary production in the riverine ecosystems (Wootton 1999).

The results of calculation of the HSI showed that almost the intermediate stations have higher amount of (= 1) HSI index. It can be deduced that the habitat of choice in the area between the rivers for Kura barbel is Tootkabon. The highest HIS index was at station 7 (0.97), while the lowest at station 1 (0.29). Presence of these conditions with HSI of 0.798 indicates that Tootkabon River provide an excellent habitat for *B. cyri* (Tables 4 & 5). The limitations of using an electrofishing device (Yu & Lee 2002; Mercado-Silva *et al.* 2008), considering the limited sampling period and the variability of the habitat features within each station, may affect the efficiency of the sampling procedure. Fish habitat-use patterns may vary by changing the environmental conditions and be affected by

seasonal patterns (Copp & Vilizzi 2004; Gillette *et al.* 2006). Seasonal patterns were not assessed here, but the habitat use and selection patterns of *B. cyri* are indicative of autumn. Therefore, we recommend investigation of the habitat use and preference patterns in other seasons as well.

## REFERENCES

- Ahmadi-Nedushan, B, St-Hilaire, A, Berube, M, Robichaud, E, Thie Monge, N & Bobe, B 2006, A review of statistical methods for the evaluation of aquatic habitat suitability for in-stream flow assessment. *River Research and Applications*, 22: 503–523.
- Breitenstein, ME & Kirchhofer, A 2000, Growth, age structure and species association of the cyprinid *Alburnoides bipunctatus* in River Aare, Switzerland. *Folia Zoologica*, 49: 59–68.
- Cihar, J 1999, *A field guide in color to freshwater fish*. Blitz ed., Leicester. 184p.
- Bovee, KD 1982, *A guide to stream habitat analysis using the in-stream flow incremental methodology*. Washington, DC: U.S. Fish and Wildlife Service. FWS/OBS-82/26.273p.
- Chuang LC, Lin YS & Liang SH 2006, Ecomorphological comparison and habitat preference of 2 cyprinid fishes, *Varicorhinus barbatulus* and *Candidia barbatus*, in Hapen Creek of Northern Taiwan. *Zoological Studies*, 45: 114–123.
- Coad B 2015, *Fresh water fishes of Iran*. Retrieved from [www.Briancoad.com](http://www.Briancoad.com). Accessed 25th Oct. 2015. 589 p.
- Copp G & Vilizzi L 2004, Spatial and ontogenetic variability in the microhabitat use of stream-dwelling spined loach (*Cobitis taenia*) and stone loach (*Barbatula barbatula*). *Journal of Applied Ichthyology*, 20: 440–451.
- Esmaili HR, Coad BW, Mehraban HR, Masoudi M, Khaefi R, Abbasi K, Mostafavi H & Vatandoust S 2014a, An updated checklist of fishes of the Caspian Sea basin of Iran with a note on their zoogeography. *Iranian Journal of Ichthyology*, 1: 152–184.
- Esmaili HR, Teimory A, Owfi F, Abbasi K & Coad BW 2014b, Alien and invasive freshwater fish species in Iran: Diversity, environmental impacts and management. *Iranian Journal of Ichthyology*, 1: 62–72.
- Gillette D, Tiemann J, Edds D & Wildhaber M 2006, Habitat use by a Midwestern USA riverine fish assemblage: effects of season, water temperature and river discharge. *Journal of Fish Biology*, 68: 1494–1512.
- Guay J, Boisclair D, Rioux D, Leclerc M, Lapointe M & Legendre P 2000, Development and validation of numerical habitat models for juveniles of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, 57: 2065–2075.
- Hasanli, AM, 1999, *Diverse methods to water measurement (Hydrometry)*. Shiraz University Publication. 265 p. (In Persian).
- Helfman, GS, Collette, BB, Facey, DE & Bowen, BW 2009, *the diversity of fishes: Biology, Evolution, and Ecology*. Blackwell Publishing, UK, Oxford, 736p.
- Johnston, N, Slaney, P & Forest Renewal B 1996, *Fish Habitat Assessment Procedures*.
- Jouladeh-Roudbar A, Vatandoust S, Eagderi S, Jafari Kenari S & Mousavi-Sabet, H 2015b. Freshwater fishes of Iran; an updated checklist. *AACL Bioflux*, 8: 855–909.
- Jowett, I 2014, Jowett Consulting. [www.jowettconsulting.co.nz](http://www.jowettconsulting.co.nz). Accessed on December 9, 2014.
- Jowett, IG, Parkyn, SM & Richardson, J 2007, Habitat characteristics of crayfish (*Paranephoves Planifrons*) in New Zealand streams using generalized additive models (GAMs). *Hydrobiologia*, 291: 264–177.
- Kirchhofer, A 1997, the assessment of Fish vulnerability in Switzerland based on distribution data. *Biological Conservation*, 80: 1–8.
- Klaar, M, Copp, GH & Horsfield, R 2004, Autumnal habitat use of non-native pumpkinseed *Lepomis gibbosus* and associations with native fish species in

- small English streams. *Folia Zoologica*, 53: 189-202.
- Li, F, Cai, O, Fu, X & Liu J 2009, Construction of habitat suitability models (HSMs) for benthic macro invertebrate and their applications to in-stream environmental flows: A case study in Xiangxi River of Three Gorges Reservoir region, China. *Progress in Natural Science*, 19: 359-367.
- Lotfi, A 2012, *Guideline on rapid assessment of environmental features of rivers*. Environment Protection Department of Iran Publication, 120p. (In Persian)
- Littlejohn, S, Holland, L, Jacobson, R, Huston, M & Hornung, T 1985, *Habits and habitats of fishes in the Upper Mississippi River*. US Fish and Wildlife Service, La Crosse, Wisconsin. 17: 304-324
- Lusk, S, Lusková, V, Halacka, K, Šlechta, V & Šlechtová V 1998, Trends and production of fish communities of the barbell zone in a stream of the Czech Republic. *Folia Zoologica*, 47: 67-72.
- MacKenzie AR & Greenberg L 1998, the influence of instream cover and predation risk on microhabitat selection of stone loach *Barbatula barbatula* (L.). *Ecology of Freshwater Fish*, 7: 87-94.
- Mercado-Silva N, Gilroy DJ, Erdenebat M, Hogan Z, Chandra S & Vander Zanden MJ 2008, Fish community composition and habitat use in the Eg-Uur River system, Mongolia. *Mongolian Journal of Biological Sciences*, 6: 21-30.
- Naseka, AM & Bogutskaya, NG 2007, Eco-zoogeographic eco-regions of continental water bodies of North Eurasia based on data on fish assemblages content. *Sbornik Nauchnykh Trudov FGNU GosNIORKh*, 337: 211-242. (In Russian).
- Orth, B 1988, Use of habitat guilds of fishes to determine in-stream flow requirements. *North American Journal of Fisheries Management*, 8:399-409.
- Quist, M, Rahel, F & Hubert, W 2005, Hierarchical faunal filters: an approach to assessing effects of habitat and nonnative species on native fishes. *Ecology of Freshwater Fish*, 14: 24-39.
- Rashleigh, AP 1995, *Conservation and Biodiversity*. Scientific American Library, New York. 264 p.
- Riffart, R, Carrel, G, Le Coarer, Y & Fontez, BNT 2009, Spatio-temporal patterns of fish assemblages in a large regulated alluvial river. *Freshwater Biology*, 54: 1544-1559.
- Rosenfeld, J 2003. Assessing the habitat requirement of stream fishes: An overview and evaluation of different approaches. *Transaction of the American Fisheries Society*, 132: 953-968.
- Smokorowski, KE & Pratt, TC 2007, Effect of a change in physical structure and cover on fish and fish habitat in freshwater ecosystems- a review and meta-analysis. *Environmental Reviews*, 15: 15-41.
- Shyi-Liang, Yu & Pefers, EJ 2002, Diel and Seasonal Habitat Use by Red Shiner (*Cyprinella lutrensis*). *Zoological Studies*, 41: 229-235.
- Tabatabaei, SN, Hashemzadeh Segherloo, I, Eagderi, S & Zamani Faradonbeh, M 2015, Habitat use of two nemacheilid fish species, *Oxynoemacheilus bergianus* and *Paracobitis* sp. in the Kordan River, Iran. *Hydrobiologia*, 762: 183-193.
- Tahami MS, Esmaeili HR & Safaie, M 2015, Population dynamic parameters of the highly endemic fish, *Alburnoides qanati* Coad & Bogutskaya 2009, (Teleostei: Cyprinidae) in the Kor River Basin, Iran. *International Journal of Aquatic Biology*, 3: 119-128.
- Tejerina-Garro, FL, Maldonado M, Ibanez C, Pont D, Roset N & Oberdorff T 2005, Effects of natural and anthropogenic environmental changes on riverine fish assemblages: a framework for ecological assessment of rivers. *Brazilian Archives of Biology and Technology*, 48: 91-108.
- Torgersen CE & Close DA 2004, Influence of habitat heterogeneity on the distribution of larval Pacific lamprey (*Lampetra tridentata*) at two spatial scales. *Freshwater Biology*, 49: 614-630.



- Treer T, Piria M, Anicic I, Safner R, Tomljanovic T 2006, Diet and growth of spirin, *Alburnoides bipunctatus* in the barbel zone of the Sava River. *Folia Zoologica*, 55: 97-106.
- Thurrow, RF 1997, Habitat utilization and dial behavior of juvenile bull trout (*Salvelinus confluentus*) at the onset of winter. *Ecology of Freshwater Fish*, 6: 1-7.
- Vinagre, C, Fonseca, V, Cabral, H & Costa, MJ 2006, Habitat suitability index models for the juvenile soles, *Solea solea* and *Solea senegalensis*, in the Tagus estuary: Defining variables for species management. *Fisheries research*, 82:140-149.
- Waddle, T 2001, *Phabsim for Windows, User's Manual and Exercises*. US Department of the Interior, US Geological Survey.
- Wootton, RJ, 1991, *Fish Ecology*. Springer Science & Business Media. 212 p.
- Yu, SL & Lee, TW 2002, Habitat preference of the stream fish, *Sinogastromyzon puliensis* (Homalopteridae). *Zoological Studies-Taipei*, 41: 183-187.

Archive of SID

## بررسی شاخص مطلوبیت زیستگاه سس ماهی کورا (*Barbus cyri*) در رودخانه توتکابن، حوضه جنوبی دریای خزر، ایران

ه. اسدی<sup>۱\*</sup>، م. ستاری<sup>۱</sup>، س. ایگدری<sup>۲</sup>

۱- گروه شیلات، دانشکده منابع طبیعی، دانشگاه گیلان، واحد صومعه سرا، ایران

۲- گروه شیلات، دانشکده منابع طبیعی، دانشگاه تهران، واحد کرج، ایران

(تاریخ دریافت: ۹۴/۲/۱۳ تاریخ پذیرش: ۹۴/۷/۲۹)

### چکیده

در اجرای طرح‌های حفاظت از اکوسیستم‌های آبی و بهره‌برداری‌های پایدار وجود دانش کافی درباره نیازهای زیستگاهی گونه‌های آبی اهمیت زیادی دارد. بنابراین، مطالعه حاضر با هدف شناخت ترجیح زیستگاهی و شاخص‌های مطلوبیت ویژگی‌های زیستگاهی گونه سس ماهی کورا (*Barbus cyri*) در محدوده پراکنش آن در رودخانه توتکابن انجام شد. برای بررسی ویژگی‌های زیستگاهی این گونه، فراوانی این ماهی و متغیرهای زیستی، شامل ارتفاع، عمق، عرض، سرعت جریان، قطر متوسط سنگ و دما با نمونه برداری از ۱۳ ایستگاه و با سه تکرار از پایین دست بسمت بالادست رودخانه توتکابن در آبان‌ماه ۱۳۹۲ بررسی شد و محدوده زیستگاه انتخابی برای این گونه بدست آمد. با توجه به نتایج بدست آمده، شاخص‌های مطلوبیت زیستگاهی این گونه ارتفاع ۲۲۰-۱۳۰ متر، عمق در دامنه ۷۵-۱۸ سانتی‌متر، عرض از رودخانه کمتر از ۱۲ متر، شیب ۲،۳-۵،۰٪، سرعت کمتر از ۰،۸ متر بر ثانیه، بستر سنگلاخی و میانگین قطر سنگ بستر در دامنه ۵۰-۱۵ سانتی‌متر قرار داشت. وجود این شرایط با شاخص مطلوبیت ۰/۷۹۸ نشان می‌دهد که رودخانه توتکابن برای گونه ماهی زردپر زیستگاهی بسیار عالی است.

\*مؤلف مسئول