



Effects of xenostrogen treatment (4-Nonylphenol, Bisphenol A) on vitellogenin expression in juvenile *Cyprinus carpio*

Samar Mortazavi^{1*}, Ali Reza Riyahi Bakhtiari², Abbas Esmaili Sari², Fatemeh Rahbarizadeh³, Nasrin Hassanzadeh¹

1. Department of Environmental Sciences, Faculty of Natural Resource and Environment, Malayer University, Hamedan, Iran

2. Department of Environmental Sciences, Faculty of Natural Resource and Marine Science, Tarbiat Modares University, Noor, Mazandaran, Iran

3. Department of Medical Biotechnology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

* Corresponding author's E-mail: mortazavi.s@gmail.com

ABSTRACT

The aim of this study was to examine some potential effects of 4-Nonylphenol (4-NP), Bisphenol A (BPA), and mixture of them on common carp, *Cyprinus carpio* and demonstrate induction of vitellogenin (Vtg) as a biomarker for screening aquatic ecosystems. These compounds with other estrogenic chemicals may be responsible for disrupting effects observed in fish. A total of 100 juvenile carp were treated experimentally with 17 β -estradiol (E2), and increasing doses of 4-NP, BPA and also mixture of them. Then the plasma Vtg levels were measured using indirect competitive ELISA. Results showed a significant ($P < 0.05$) increase in Vtg levels of exposed carps with E2, maximum 4-NP, BPA and their mixture. The group treated with E2 showed high induction (490%) with respect to controls. The groups treated with maximum dose of 4-NP and BPA showed also induction of Vtg, whereas the group treated with their mixture (3 mg kg⁻¹ b.w. 4-NP and 0.01 mg kg⁻¹ b.w. BPA) showed the highest induction (2126%). Finally, results showed that mixture of 4-NP and BPA exhibited stronger (synergistic) effects. So that, large scale monitoring of estrogenic effects caused by contamination may be considered as a biomarker in urban and industrial effluents.

Key words: Vitellogenin (Vtg), *Cyprinus carpio*, 17 β -Esteradiol (E2), 4-Nonylphenol (4-NP), Bisphenol A (BPA), ELISA.

INTRODUCTION

Environmental problems caused by estrogenic compounds are mostly related to aquatic ecosystem (Matozzo *et al.* 2008). During the last decades, the use of aquatic systems as final receptors of urban and industrial waste waters has increased noticeably, leading to a decrease in the general quality of shores (De los Rios *et al.* 2012), ending in alterations in the composition and structure of communities and finally, damaged ecosystems (Estácio *et al.* 1997; Smith & Shackley 2006; Okus *et al.* 2008; Stein & Cadien 2009; DelPilar Ruso *et al.* 2010).

In such conditions, treated and untreated municipal and industrial sewage, agricultural activities, livestock wastes and system treatment plants (STPs) are considered as the major sources of xenoestrogens entering aquatic ecosystem (Ying *et al.* 2002; Atkinson *et al.* 2003; Tashiro *et al.* 2003; Kuster *et al.* 2004; Matozzo *et al.* 2008). Since sewage and manure are often used as fertilizer, detergents and agricultural effluent are considered as other sources of hormonal pollution in aquatic environments (Orlando *et al.* 2004; Soto *et al.* 2004). Consequently, large amounts of xenoestrogens are discharged in aquatic environments, where they may adsorb to sediments and last for quite a long time (Lai *et al.* 2000).

Among chemicals proven to be endocrine disrupting chemicals (EDCs), alkylphenols (APs) and alkylphenoethoxylates (APEs) are two of the most important classes, not only for the quantity released into the environment, but also for their proved estrogenic activity (Solé *et al.* 2000; Riva *et al.* 2010). Almost all of the

APEs manufactured worldwide are nonylphenolet- hoxylates (NPEs), a large group of nonionic surfactants used in lubricating oils, emulsifiers, plastics, latex paints, household and industrial detergents, and in the paper and textile industry (Lee, 1999). After usage, NPEs are released into municipal and industrial sewage treatment plants, gradually degrading to nonylphenols (NPs), whose eventual compartment is the aquatic environment (Ahel *et al.* 1994; Taylor & Harrison 1999; Matozzo *et al.* 2008). Among NPs, 4-nonylphenol (4-NP) has been identified as the most important metabolite due to its high resistance to biodegradation, toxicity and intense estrogenic effects (Taylor & Harrison 1999). In addition, 4-NP is a lipophilic compound ($\log Kow=4.48$, (Ahel & Giger 1993) and may therefore be taken up and bio-accumulated by aquatic organisms.

Another important compound existing in the effluent is Bisphenol A (BPA). While primarily applied as an intermediate and additive monomer in plastics and epoxy resins (Hohenblum *et al.* 2004), it is also used as a photo-initiator, or as an additive in insecticides, pharmaceuticals, and adhesives. In 1993 the annual universal production of BPA hit 640,000 tons, out of which an estimated 0.017% (109 tons) were discharged into the environment (Benjonathan & Steinmetz 1998; Staples *et al.* 1998; Matozzo *et al.* 2008).

Due to phenyl ring in their chemical structures, BPA and APs usually possess similar endocrine disrupting activities in general, and estrogenic ones in particular (Gutendorf & Westendorf 2001; Kwack *et al.* 2002; Xu *et al.* 2005). By binding to estrogenic receptors, these compounds can mimic the action of endogenous estrogens.

In fish, one of the best-proven effects of xenoestrogens is the induction of vitellogenins (Vtg), base of the egg-yolk proteins, which provide energy reservoirs for embryo development in oviparous organisms. Vtgs. are high-density glycolipophosphoproteins having Ca and Zn ligands (Montorzi *et al.* 1994; Denslow *et al.* 1999; Matozzo *et al.* 2008; Scognamiglio *et al.* 2016).

Vtg is basically a female particular serum protein. Although the liver synthesizes Vtg in females in reaction to stimulation by 17β -estradiol (E2) (Truscott *et al.* 1992), in males and juveniles Vtg is normally not noticeable. However, exposure to EDCs may cause the induction of Vtg to measurable concentrations even in male and juvenile fish. This has been shown in various laboratory and field studies (Jensen & Ankley 2006; Vega-Lopez *et al.* 2006; Zha *et al.* 2006; Reinen *et al.* 2012).

As pollutants usually occur as mixtures in real aquatic ecosystems, fish are unlikely to be exposed to just one estrogenic chemical. Since we ideally need to know how a fish responds to a mixture of estrogenic chemicals, rather than to an individual chemical, we should try to stimulate the real world, considering the interactions between different chemicals in a mixture, which may result in either a weaker (antagonistic) or a stronger (synergistic) effect.

The main objective of the current study was to examine some potential mechanism(s) of action for 4-NP, BPA, and the mixture of these compounds in sediments of Anzali Wetland, on *Cyprinus carpio*, inhabiting this aquatic ecosystem. The results of the study will relate to the use and design of in vitro bioassays to characterize the endocrine modulating potential of these chemicals and similar compounds.

MATERIALS AND METHODS

4-Nonylphenol (4-NP), Bisphenol A, and 17β -estradiol were purchased from Sigma Aldrich Company (St. Louis, Missouri, USA). Stock solutions of these chemicals were prepared in peanut oil (Sigma) and stored at 20°C in refrigerator in tightly-closed and dark-glass bottles. Juvenile common carps (*Cyprinus carpio*) of 9-10-month old's weighing 80-110 g were obtained from Nasr fish farming, around (airport road) Sari city of Iran, the hatchery of the fish culture. The fish were transferred to the research center of Tarbiat Modares University and were kept in a big fiber glass tank, held in a 12:12 h light-dark photoperiod and not fed during the experiment. The water supply was spring water, quality parameters of which including ammonia, nitrite, dissolved oxygen (DO), pH, hardness, and conductivity were monitored weekly (Table 1). No significant tank-to-tank or week-to-week differences in any of the water quality parameters were indicated (Villeneuve 2000; Casini *et al.* 2002). Water temperature was 17°C and dissolved oxygen was preserved above 60% of the saturation level by continuously aerating the test solutions.

After 5 days, the carps were divided into 9 groups of 9 specimens kept in separate tanks (20 L). The control group was injected intraperitoneally with peanut oil; of the 8 exposed groups, one was injected with 10 mg kg^{-1} b.w. 17β -estradiol in peanut oil and three others were injected with 3, 30 and 60 mg kg^{-1} b.w. 4-nonylphenol in peanut oil respectively. Three of the remaining groups were injected with 0.1, 1 and 10 mg kg^{-1} b.w. Bisphenol A, and

the last group was injected with 3 and 0.1 mg kg⁻¹ b.w. mixtures of 4-Nonylphenol and Bisphenol A. The treatment was repeated after intervals of 7 days for 3 times. 3 days after the last injection, blood samples were obtained from the dorsal vein by puncture for vitellogenin analysis. Blood was collected into heparinized 5ml glass tubes and centrifuged for 4 min at 4000g in a centrifuge. Plasma was pipetted into 1 mL Eppendorf centrifuge tubes, and stored at -80°C until analysis.

Table 1. Average water quality during exposure in weekly monitoring.

| Water Quality | Mean ± S.D |
|--|-------------|
| Temperature (°C) | 17 ± 0.5 |
| pH | 7.61 ± 0.02 |
| Dissolved Oxygen (mg L ⁻¹) | 9 ± 0.4 |
| Ammonia (mg L ⁻¹) | 0.57 |
| Nitrite (mg L ⁻¹) | 0.01 |
| Nitrate (mg L ⁻¹) | 0.22 |
| Hardness (mg L ⁻¹) | 120 ± 5 |
| Phosphate (mg L ⁻¹) | 0.36 |

Vitellogenin Analysis: Plasma levels of Vtg were determined by a commercial enzyme-linked immune sorbent assay developed for *Cyprinus carpio* (ELISA, Biosense Laboratories, Bergen, Norway). This assay had a quantification limit of 0.1 ng mL⁻¹, which did not allow for the accurate determination of the low Vtg concentrations found in the control and some treatments. The procedure of the ELISA was as follows (Christensen et al. 1998; Lindhost et al. 2002; Mandich et al. 2007): The 96-well polystyrene microtiter plates were supplied and precoated with *C. carpio* Vtg monoclonal antibody. Vtg. standard solutions (range 0.24 to 250 ng mL⁻¹ linear two-fold serial dilution) were distributed (50 µL) into dedicated wells while two nonspecific banding (NSB) wells received 100 µL dilution buffer (DB). The plasma samples were diluted 1:10; 1:100 in DB and 100 µL of the diluted samples were dispersed.

Plates were incubated for 1 h at 37 °C and then washed three times with 300 µL washing solution (Phosphate Buffered Saline (PBS), 0.05% Tween-20. In the next step, detecting anti-Vtg antibodies (100 µl) were distributed to each well, incubated for 1 h at 37°C and then washed three times with 300 µL washing solution. Secondary anti-IgG antibodies, horseradish peroxidase conjugate (100 µl), were instantly distributed to each well, incubated for 1 h at 37 °C and then washed five times with 300 µL washing solution for staining. Each well received 100 µL substrate solution (OPD-peroxidase substrate in urea hydrogen peroxide) and the plates were incubated for 30 min at room temperature (20–25 °C), then the reactions were stopped by adding 50 µL 2M H₂SO₄ to all wells. Finally, the absorbance at 492 nm was read with microplate reader.

The working range of the Vtg standard is normally 0.98–125 ng Vtg mL⁻¹ and, the intra and inter assay coefficients of variation (% CV) are between 5.6 and 24% within the working range (Table 2).

Table 2. Carp Vtg standard.

| Vtg concentration (ng mL ⁻¹) | absorbance | NSB-corrected absorbance |
|--|------------|--------------------------|
| 250 | 2.062 | 2.017 |
| 62.5 | 1.738 | 1.725 |
| 15.6 | 0.816 | 0.771 |
| 1.95 | 0.190 | 0.145 |
| 0.49 | 0.0875 | 0.0425 |
| 0.24 | 0.063 | 0.018 |

Statistical analysis

The SPSS version 11.7 was used for statistical analysis. In case of inhomogeneity of variances, the nonparametric Kruskal-Wallis test was performed, followed by the Mann-Whitney U test.

RESULTS AND DISCUSSION

The results obtained from ELISA analysis are shown in (Figs. 1, 2 and, 4). The group treated with 17β-estradiol had high absorbance values (490% variation) (Fig 3). The Mann-Whitney U test results indicated significant difference ($Z = -2.138$, $p < 0.05$) in Vtg levels between groups treated with 17β-estradiol and the control group. On the other hand, groups exposed to different (low to high) concentrations of 4-Nonylphenol showed enhanced

absorbance values (Fig. 1). In addition, the Mann-Whitney U test results pointed out significant difference ($Z = -2.132$, $p < 0.05$) between groups treated with maximum 4-Nonylphenol and the control group in them.

Vtg levels. The treatment group injected by maximum concentration of Bisphenol A had high absorbance (Fig. 2). Moreover, Mann-Whitney U the test results showed significant difference ($Z = -2.123$, $p < 0.05$) between groups treated with maximum Bisphenol A and the control group in Vtg levels. The most important result of the present study belonged to the group exposed to the mixture of Bisphenol and 4-Nonylphenol. This final group had the highest absorbance value (2126%) (Fig. 3). This is while the results of Kruskal-Wallis test ($H = 6.4$; $df = 2$; $p < 0.05$) proved significant difference between this latter group and groups treated with 17β -estradiol and control (Fig. 4). All in all, comparison of estrogenic potentials of the four selected chemical compounds in ELISA revealed that relative potency of these chemicals was decreased in the following order: mixture of 4-NP and PBA > 17β -Esteradiol > 4-NP > BPA (Fig. 4). Statistically significant correlation ($p < 0.01$) was found between dose and response in plasma Vtg levels in treatments injected with increasing dose of 4-NP and BPA (Fig. 5). 100 Juvenile female carps were chosen for this study. Carps are commonly-occurring fish species in Europe and Asia, relatively easy to keep in the laboratory, and large enough to provide tissue and plasma volumes adequate for analyses. In addition, laboratory studies with carp support constant field work involving the use of vitellogenin in wild carp as a biomarker of exposure to xenoestrogens (Folmar *et al.* 1996; Goodbred *et al.* 1997; Snyder 2001; Villeneuve *et al.* 2002; Li *et al.* 2004; Li *et al.* 2010; Gilannejad *et al.* 2016). In this study the sensitivity of carp to xenoestrogens was compared by measuring the effects of natural estrogens and anthropogenic compounds on Vtg induction in primary hepatocytes. The fish tested in this study were exposed to environmentally relevant concentrations of 4-NP and BPA in sediments of Anzali Wetland. 4-NP and BPA concentrations ranging from 0.05 to $29 \mu\text{g g}^{-1}$ d.w. and 0.01 to $6.97 \mu\text{g g}^{-1}$ d.w. have been reported respectively (Mortazavi *et al.* 2012).

A representative range of chemicals reported to be estrogenic have been tested to determine whether they are also estrogenic to carp or not, and if so, to assess approximately how potent. Some of the results obtained are shown in Fig. 1. All of the chemicals examined stimulated synthesis of Vtg in a dose-dependent manner. However, some of these chemicals stimulated Vtg synthesis at concentrations reported to be present in the aquatic environment. In natural situations, fish are continuously subjected to any estrogenic chemicals present in the water. The results revealed that exposure to minimum concentrations of 4-NP and BPA in sediments did not cause significant Vtg induction in juvenile female carp (Figs 1 and 2). This is in agreement with other studies demonstrating significant Vtg induction only at greater exposure concentrations. $10 \mu\text{g L}^{-1}$ NP was reported to be the threshold exposure concentration needed to cause Vtg induction in 2-year-old rainbow trouts (Jobling *et al.* 1996; Villeneuve *et al.* 2002). Another study proved significant induction of Vtg in rainbow trout exposed to concentrations greater than $25 \mu\text{g L}^{-1}$ NP (Tremblay & Van DerKraak 1998).




Fig. 1. Plasma Vtg levels (mean \pm standard error) in juvenile carps: control, experimentally treated 17β -estradiol (E2) (10 mg kg^{-1} b.w.) and increasing dose of 4-NP (3, 30, 60 mg kg^{-1} b.w.).




Fig. 2. Plasma Vtg levels (mean ± standard error) in juvenile carps: control, experimentally treated 17β-estradiol (E2) (10 mg kg⁻¹ b.w.) and increasing dose of BPA (0.1, 1, 10 mg kg⁻¹ b.w.).




Fig. 3. Percentage variation of plasma Vtg in treated groups with respect to control group.

Another study with fathead minnows exhibited no significant induction of Vtg in males at exposure concentrations up to 2.4 μg L⁻¹ NP, but a significant one in females (Giesy *et al.* 2000). Therefore, compared to other reports in the literature, it seems that exposure to high concentrations of the chemicals used for this study were adequate to induce Vtg in juvenile female carp. Other studies also detected marked increases in plasma Vtg concentrations in Atlantic cod, *Gadus morhua*, Atlantic salmon, and *Salmo salar*, after injection of E2 (peaking on day 6) and 4-NP (Hylland & Haux 1997; Ratna *et al.* 2016). Vtg induction has also been observed in various fish species exposed to diluted effluents from wastewater treatment (Jobling *et al.* 2003; Robinson *et al.* 2003; Marin *et al.* 2004). In the present study, mixture of 4-NP and BPA produced significant elevation in plasma Vtg levels. In fact, there was no evidence of synergistic influences of the examined chemicals with respect to their ability to induce plasma Vtg levels. In addition, when NP was injected in combination with BPA, like research of Arukwe *et al.* (2000), Vtg levels were more elevated. Since environmental samples are mixtures of many compounds, including agonists and antagonists, several studies have dealt with this issue. Seifert *et al.* (1999) and Li *et al.* (2004) examined the mixtures of different estrogenic chemicals including antiestrogen tamoxifen with ELRA.




Fig. 4. Plasma Vtg levels (mean \pm standard error) in juvenile carps: control, experimentally treated 17 β -estradiol (E2) (10 mg kg⁻¹ b.w.) and increasing dose of 4-NP, BPA, and mixture of them (3 and 0.1, mg kg⁻¹ b.w.) respectively.




Fig. 5. Significant correlation between dose and response in plasma Vtg levels in Juvenile carps experimentally treated with increasing doses of 4-NP and BPA.

In all cases, synergism was found. In another study, Silva *et al.* (2002) examined the mixtures of several agonists including Bisphenol-A with YES assay, synergism was also detected. These results validate findings of previous studies on marine fish, especially those by Arukwe *et al.* (1997) in which the treatment of *Salmo salar* with the same two compounds showed induction of Vtg in plasma of the same amount.

Similar to maturely differentiated specimens (Folmar *et al.* 1996; Solè *et al.* 2000; Villeneuve *et al.* 2002), juvenile common carps are also responsive to estrogenic incentives. The fact that this species is also sensitive to relatively low doses of xenoestrogens, such as 4-NP, might have an outcome on natural populations.

With further evidence, this fish might be proposed for large-scale monitoring of estrogenic effects caused by contamination in urban and industrial sewage. Carp is in fact pervasive in most fresh water courses, easily available, and does not migrate (Casini *et al.* 2002) and can therefore represent a good sentinel organism.

CONCLUSION

The findings of laboratory studies show that a rapidly increasing number of chemicals, entering water ways through different sources of effluent, or their degradation products, are being recognized as estrogenic.

As a result, Vtg levels may increase when animals are exposed to a wide group of pollutants having similar feminizing effects. This is while Vtg induction may be considered as a useful tool in evaluating endocrine disruption in aquatic animals, and therefore the potential risk associated with exposure to xenoestrogens. The repercussions of this exposure and the responses to it are unknown for the present time, but they could include opposing effects on physiological processes, particularly reproduction.

In addition, there is a considerable relevance between the ecological and the physiological responses, considering that hormonal changes may cause variations in the sex ratio of a feral population, its reproductive capability, and even the presence of the species in the aquatic ecosystem.

Moreover, using only one (or a few) biomarkers can reduce expensive chemical analyses of individual pollutants and mostly allow detection of their interactions in the aquatic ecosystem.

Further work is required to clarify the effects of other estrogenic chemicals present in the environment on aquatic animals.

ACKNOWLEDGMENT

This study received a partial financial support from the Ministry of Science, Research and Technology, Iran, for which we are grateful.

We would like to thank Zohreh Hasanabadi, Asma Karami, Shahedeh Abdolazizi and Maryam Saghafi from the Faculty of Natural Resources and Marine Sciences, Tarbiat Modares University, for their kind assistance in preparing the samples. We would also like to thank Azin Kazemi, for her editorial suggestions on the first draft and Dr. Kazempour on the final draft of the paper.

REFERENCES

- Ahel, M, Scully, FE, Hoigné, J & Giger, W 1994, Photochemical degradation of nonylphenol and nonylphenol polyethoxylates in natural waters. *Chemosphere*, 28: 1361–1368.
- Arukwe, A, Celius, T, Walther, BT & Goksøyr, A 2000, Effects of xenoestrogen treatment on zona radiata protein and vitellogenin expression in Atlantic salmon (*Salmo salar*). *Aquatic Toxicology*, 49: 159–170.
- Arukwe, A, Knudsen, FR & Goksoyr, A 1997, Fish zona radiate (egg shell) protein: A sensible biomarker for environmental estrogens. *Environmental Health Perspectives*, 105: 418–422.
- Atkinson, S, Atkinson, MJ & Tarrant, AM 2003, Estrogens from sewage in coastal marine environments. *Environmental Health Perspectives*, 111: 531–535.
- Benjonathan, N & Steinmetz, R 1998, Xenoestrogens the emerging story of bisphenol A. *Trends in Endocrinology and Metabolism*, 9: 124–128.
- Casini, S, Fossi, MC, Mori, G & Bjornstad, A 2002, Vitellogenin duction in *Cyprinus carpio* treated with 17 β -Estradiol and 4-Nonylphenol. *Environmental Monitoring and Assessment*, 75, 235–239.
- Christensen, LB, Pedersen, KL, Korsgaard, B & Bjerregaard, P 1998, Estrogenicity of xenobiotics in rainbow trout (*Oncorhynchus mykiss*) using in vivo synthesis of vitellogenin as a biomarker. *Marine Environmental Research*, 46: 137–140.

- De los Rios, A, Juanes, JA, Ortiz-Zarragoitia, M, de Alda, ML, Barcel, D & Cajaraville, MP 2012, Assessment of the effects of a marine urban outfall discharge on caged mussels using chemical and biomarker analysis. *Marine Pollution Bulletin*, 64: 563–573.
- Del Pilar Ruso, Y, de la Ossa, JA, Giménez, F & Sanchez-Lizaso, JL 2010, Sewage treatment level and flow rates affect polychaete assemblages. *Marine Pollution Bulletin*, 60: 1930–1938.
- Denslow, ND, Chow, MC, Kroll, KJ & Green, L 1999, Vitellogenin as a biomarker of exposure for estrogen or estrogen mimics. *Ecotoxicology*, 8: 385–398.
- Esacio, FJ, Garcia-Diego, EM, Garcia-Gomez, DA, Daza, JL, Hortas, F & Gomez-Ariza, JL 1997, Ecological analysis in a polluted area of Algeciras Bay (Southern Spain): external ‘versus’ internal outfalls and environmental implications. *Marine Pollution Bulletin*, 34: 780–793.
- Folmar, LC, Denslow, ND, Rao, V, Chow, M, Crain, DA, Enblom, J, Marcino, J & Guillette Jr, LJ 1996, Vitellogenin induction and reduced serum testosterone concentrations in feral male carp (*Cyprinus carpio*) captured near a mayor metropolitan sewage treatment plant. *Environmental Health Perspectives*, 104: 1096–1101.
- Giesy, JP, Pierens, SL, Snyder, EM, Miles-Richardson, SM, Kramer, VJ, Snyder, SA, Nichols, KM & Villeneuve, DL 2000, Effects of 4-nonyl phenol on fecundity and biomarkers of estrogenicity in fathead minnows (*Pimephales promelas*). *Environmental Toxicology and Chemistry*, 19: 1368–1377.
- Gilannejad, N, Dorafshan, S, Heyrati, FP, Soofiani, NM, Asadollah, S, Martos-Sitcha, JA, Prat, F, Yúfera, M & Martínez-Rodríguez, G 2016, Vitellogenin expression in wild cyprinid *Petroleuciscus esfahani* as a biomarker of endocrine disruption along the Zayandeh Roud River, Iran. *Chemosphere*, 144: 1342–1350.
- Goodbred, SL, Gilliom, RJ, Gross, TS, Denslow, NP, Bryant, WB & Schoeb, TR 1997, Reconnaissance of 17-betaestradiol, 11-ketotestosterone, vitellogenin, and gonad histopathology in common carp of United States streams: potential for contaminant-induced endocrine disruption. United States Department of the Interior-US Geological Survey, Sacramento, CA, USA, pp. 20–28.
- Gutendorf, B & Westendorf, J 2001, Comparison of an array of in vitro assays for the assessment of the estrogenic potential of natural and synthetic estrogens, phytoestrogens and xenoestrogens. *Toxicology*, 166: 79–89.
- Hohenblum, P, Gans, O, Moche, W, Scharf, S & Lorbeer, G 2004, Monitoring of selected estrogenic hormones and industrial chemicals in ground waters and surface waters in Austria. *Sci. Total Environment*, 333: 185–193.
- Hylland, K & Haux, C 1997, Effects of environmental oestrogens on marine fish species. *Trends in Analytical Chemistry*, 16: 606–612.
- In the fathead minnow (*Pimephales promelas*). *Ecotoxicology Environmental Safety*, 64: 101–105.
- Jensen, KM, Ankley, GT 2006, Evaluation of a commercial kit for measuring vitellogenin.
- Jobling, S, Casey, D, Rodgers-Gray, T, Oehlmann, J, Schulte-Oehlmann, U, Pawlowski, S, Baunbeck, T, Turner, AP & Tyler, CR 2003, Comparative responses of molluscs and fish to environmental estrogens and an estrogenic effluent. *Aquatic Toxicology*, 65: 205–220.
- Jobling, S, Sheahan, D, Osborne, JA, Matthiessen, P, Sumpter, JP 1996, Inhibition of testicular growth in rainbow trout (*Oncorhynchus mykiss*) exposed to estrogenic alkylphenolic chemicals. *Environmental Toxicology and Chemistry*, 15: 194–202.
- Kuster, M, Lopezde Alda, MJ & Barceló, D 2004, Analysis and distribution of estrogens and progestogens in sewage sludge, soils and sediments. *Trends in Analytical Chemistry*, 23: 790–798.
- Kwack, SJ, Kwon, O, Kim, HS, Kim, SS, Kim, SH, Sohn, KH, Lee, RD, Park, CH, Jeung, EB, An, BS, Park, KL 2002, Comparative evaluation of alkylphenolic compounds on estrogenic activity in vitro and in vivo. *J. Toxicology and Environmental Health, A*, 65: 419–431.
- Lai, KM, Johnson, KL, Scrimshaw, MD & Lester, JN 2000, Binding of water born steroid estrogens to solid phases in river and estuarine systems. *Environmental Science and Technology*, 34: 3890–3894.
- Lee, HB 1999, Review of analytical methods for the determination of nonylphenol and related compounds in environmental samples. *Water Quality Research Journal of Canada*, 34: 3–35.
- Li, J, Ma, M & Wang, Z 2010, In vitro profiling of endocrine disrupting effects of phenols. *Toxicology, in Vitro*, 24: 201–207.

- Li, W, Seifert, MXu, Y & Hock, B 2004, Comparative study of estrogenic potencies of estradiol, tamoxifen, bisphenol-A and resveratrol with two in vitro bioassays. *Environmental International*, 30: 329–335.
- Lindholst, C, Pedersen, KL & Pedersen. SN 2000, Estrogenic response of bisphenol A in rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology*, 48: 87–94.
- Mandich, A, Bottero, S, Benfenati, E, Cevasco, A, Erratico, C, Maggioni, S, Massari, A, Pedemonte, F & Viganò, L 2007, In vivo exposure of carp to graded concentrations of bisphenol A. *General and Comparative Endocrinology*, 153: 15–24.
- Marin, MG & Matozzo, V 2004, Vitellogenin induction as a biomarker of exposure to estrogenic compounds in aquatic environments. *Marine Pollution Bulletin*, 48: 835–839.
- Matozzo, M, Gagné, F, Marin, MG, Ricciardi, F & Blaise, C 2008, Vitellogenin as a biomarker of exposure to estrogenic compounds in Aquatic invertebrates: A review. *Environmental International*, 34: 531–545.
- Montorzi, M, Falchuk, KH & Vallee, BL 1994, *Xenopus laevis* vitellogenin is a zinc protein. *Biochemical and Biophysical Research Communications*, 200: 1407–1413.
- Mortazavi, S, Riyahi Bakhtiari, AR, Esmaili Sari, A, Bahramifar, N & Rahbarizade, F 2012, Phenolic Endocrine Disrupting Chemicals (EDCs) in Anzali Wetland, Iran: Elevated concentrations of 4-Nonylphenol, Octylphenol and Bisphenol A. *Marine Pollution Bulletin*, 46: 1067–1073
- Okus, E, Ozturk, I, Ibrahim, H, Yukseka, A, Tasa, S, Aslan-Yilmaza, A, Altioka, H, Balkisa, N, Dogana, E, Ovezb, S & Aydinb, AF 2008, Critical evaluation of waste water treatment and disposal strategies for Istanbul with regards to water quality monitoring study results. *Desalination*, 226: 231–248.
- Orlando, EF, Kolok, AS, Binzcik, GA, Gates, JL, Horton, MK, Lambright, CS, Gray Jr. LE, Soto, AM, Guillette, Jr & Louis, J 2004, Endocrine-disrupting effects of cattle feed lot effluent on an aquatic sentinel species, the fathead minnow. *Environmental Health Perspectives*, 112: 353–358.
- Ratna, WN, Bhatt, VD, Chaudhary, K, Ariff, AB, Bavadekar, SA & Ratna, HN 2016, Estrogen-responsive genes encoding egg yolk proteins vitellogenin and apolipoprotein II in chicken are differentially regulated by selective estrogen receptor modulators. *Theriogenology*, 85(3): 376–383.
- Reinen, JI Suter, MJFI Voeli, AC, Fernandez, MF, Kiviranta, H, Eggen, RIL & Vermeulen, NPE 2012, Endocrine disrupting chemicals—Linking internal exposure to vitellogenin levels and ovotestis in *Abramis brama* from Dutch surface waters. *Environmental Toxicology and Pharmacology*, 30: 209–223.
- Riva, C, Porte, C, Binelli, A & Provini, A 2010, Evaluation of 4-nonylphenol in vivo exposure in *Dreissena polymorpha*: Bioaccumulation, steroid levels and oxidative stress. *Comparative Biochemistry and Physiology Part C*, 152: 175–181.
- Robinson, CD, Brown, E, Craft, JA, Davies, IM, Moffat, CF, Pirie, D, Robertson, F, Stagg, RM & Struthers, S 2003, Effects of sewage effluent and ethynyl oestradiol upon molecular markers of oestrogenic exposure, maturation and reproductive success in the sand goby (*Pomatoschistus minutus*, Pallas). *Aquatic Toxicology*, 62: 119–134.
- Scognamiglio, V, Antonacci, A, Patrolecco, L, Lambrea, MD, Litescu, SC, Ghuge, SA & Rea, G 2016, Analytical tools monitoring endocrine disrupting chemicals. *Trends in Analytical Chemistry*, 80: 555–567.
- Seifert, M, Haindl, S & Hock, B 1999, Development of an enzyme linked receptor assay (ELRA) for estrogens and xenoestrogens. *Analytica Chimica Acta*, 386: 191–199.
- Silva, E, Rajapakse, N, Kortenkamp, A 2002, Something from nothing eight weak estrogenic chemicals combined at concentrations below NOECs produce significant mixture effects. *Environmental Science and Technology*, 15: 1751–1756.
- Smith, J, Shackley, SE 2006, Effects of closure of a major sewage outfall on sublittoral, soft sediment benthic communities. *Marine Pollution Bulletin*, 52: 645–658.
- Snyder, EM 2001, Use of fish as bioassay organisms to assess wastewater effluents for reproductive endocrine modulating chemicals. Ph.D. Dissertation. Michigan State University, East Lansing, MI, USA.
- Solé, M, Lopez de Alda, MJ, Castillo, M, Porte, C, Ladegaard- Pedersen, K & Barcelo, D 2000, Estrogenicity determination in sewage treatment plants and surface waters from the Catalanian area (NE Spain). *Environmental Science and Technology*, 34: 5076–5083.

- Solè, M, Porte, D & Barceló, D 2000, Vitellogenin duction and other biochemical responses in Carp, *Cyprinus carpio*, after experimentally injection with 17β -Ethinylestradiol. *Arch. Environmental Contamination and Toxicology*, 38: 494–500.
- Soto, AM, Calabro, JM, Prechtel, NV, Yau, AY, Orlando, EF, Daxenberger, A, Kolok, AS, Guillette Jr, LJ, Bizec, BL, Lange, IG & Sonnenschein, C 2004, Androgenic and estrogenic activity in water bodies receiving cattle feed lot effluent in eastern Nebraska, USA. *Environmental Health Perspectives*, 112: 346–352.
- Staples, CA, Dorn, PB, Klecka, GM, O'Block, ST & Harris, LR 1998, A review of the environmental fate, effects, and exposures of bisphenol A. *Chemosphere*, 36: 2149–2173.
- Stein, ED & Cadien, DB 2009, Ecosystem response to regulatory and management actions: the southern California experience in long-term monitoring. *Marine Pollution Bulletin*, 59: 91–100.
- Tashiro, Y, Takemura, A, Fujii, H, Takahira, K & Nakanishi, Y 2003, Livestock wastes as a source of estrogens and their effects on wild life of Manko tidal flat, Okinawa. *Marine Pollution Bulletin*, 47: 143–147.
- Taylor, MR & Harrison, PTC 1999, Ecological effects of endocrine disruption: current evidence and research priorities. *Chemosphere*, 8: 1237–1248 .
- Tremblay, L & Van Der Kraak, G 1998, Use of a series of homologous in vitro and in vivo assays to evaluate the endocrine modulating actions of b-sitosterol in rainbow trout. *Aquatic Toxicology*, 43: 149–162.
- Truscott, BSo, YP, Nagler, JJ & Idler, DR 1992, Steroids involved with final oocyte maturation in the winter flounder. *The Journal of Steroid Biochemistry and Molecular Biology*, 42: 351–356.
- Vega-Lopez, A, Martinez-Tabche, L, Dominguez-Lopez, ML, Garcia-Latorre, E Ramon-Gallegos, E & Garcia-Gasca, A 2006, Vitellogenin induction in the endangered goodeid fish *Girardinichthys viviparus*: Vitellogenin characterization and estrogenic effects of polychlorinated biphenyls. *Comparative Biochemistry and Physiology C, Toxicology and Pharmacology*, 142: 356–364.
- Villeneuve, DL, Villalobos, SA, Keith, TL, Snyder, EM, Fitzgerald, SD, Giesy, JP 2002, Effects of waterborne exposure to 4-nonylphenol on plasma sex steroid and vitellogenin concentrations in sexually mature male carp (*Cyprinus carpio*). *Chemosphere*, 47: 15–28.
- Xu, LC, Sun, H, Chen., JF, Bian, Q, Qian, J, Song, L & Wang, XR 2005, Evaluation of androgen receptor transcriptional activities of bisphenol A, octylphenol and nonylphenol in vitro. *Toxicology*, 216: 197–203.
- Ying, GG, Kookana, RS & Ru, YJ 2002, Occurrence and fate of hormone esterooids in the environment. *Environment International*, 28: 545–51.
- Zha, J, Wang, Z, Schlenk, D 2006, Effects of pentachlorophenol on the reproduction of Japanese medaka (*Oryzias latipes*). *Chemico-Biological Interactions*, 161: 26–36.

اثرات ترکیبات زنواستروژنی (۴-نونیل فنل، بیسفنل-آ) بر تولید ویتلوژنین در کپورماهیان نابالغ

ثمر مرتضوی^{۱*}، علی رضا ریاحی بختیاری^۲، عباس اسماعیلی ساری^۲، فاطمه رهبری زاده^۳، نسرين حسن-زاده^۱

۱- گروه محیط زیست، دانشکده منابع طبیعی و محیط زیست، دانشگاه ملایر، همدان، ایران

۲- گروه محیط زیست، دانشکده منابع طبیعی و علوم دریایی، دانشگاه تربیت مدرس، نور، مازندران، ایران

۳- گروه بیوتکنولوژی پزشکی، دانشکده علوم پزشکی، دانشگاه تربیت مدرس، تهران، ایران

(تاریخ دریافت: ۹۷/۰۷/۱۰ تاریخ پذیرش: ۹۷/۱۱/۲۳)

چکیده

هدف از مطالعه حاضر بررسی برخی از اثرات بالقوه ترکیبات نونیل فنل (NP-4)، بیسفنل-آ (BPA) و مخلوط آنها بر کپورماهی معمولی در القاء ویتلوژنین (Vtg) به عنوان بیومارکری جهت پایش بوم‌سازگان‌های آبی است. این ترکیبات و همچنین سایر مواد شیمیایی استروژنی ممکن است مسئول اثرات مخرب مشاهده شده در ماهی باشد. جهت آزمایش ۱۰۰ عدد کپور ماهی نابالغ مورد تزریق با ۱۷ بتا استرادیول (E2)، دوزهای افزایشی BPA، NP-4 و همچنین مخلوط آنها قرار گرفت. سپس سطح Vtg پلاسما با استفاده از الایزا به روش غیرمستقیم اندازه گیری شد. نتایج نشان داد میزان Vtg افزایش معناداری ($P < 0.05$) در برابر میزان E2، حداکثر دوز BPA، NP-4 و مخلوط آنها دارد. در مقایسه با گروه کنترل، گروه در معرض E2 دارای القاء بالا (۴۹۰٪) بود. گروه های با تزریق حداکثر دوز NP-4 و BPA نیز القاء Vtg را نشان دادند، در حالیکه گروه با تزریق مخلوط آنها ۳ ($\text{mg kg}^{-1} \text{ b.w}$) ۴-نونیل فنل و ۰/۰۱ ($\text{mg kg}^{-1} \text{ b.w}$) بیسفنل آ، بیشترین القاء (۲۱۲۶٪) ویتلوژنین را در قیاس با گروه کنترل نشان داد. در نهایت، نتایج نشان داد که مخلوطی از NP-4 و BPA اثرات قوی تر (هم افزایی) را نشان می‌دهد. بنابراین نظارت گسترده بر اثرات استروژنی ناشی از آلودگی ممکن است بعنوان بیومارکدر پساب‌های شهری و صنعتی قابل ملاحظه باشد.

* مؤلف مسئول

Bibliographic information of this paper for citing:

Mortazavi, S, Riyahi Bakhtiari, AR, Esmaili Sari, A, Rahbarizadeh, F, Hassanzadeh, N 2019, Effects of xenostrogen treatment (4-Nonylphenol, Bisphenol A) on vitellogenin expression in juvenile *Cyprinus carpio*. Caspian Journal of Environmental Sciences, 17: 43-53

Copyright © 2019