ORIGINAL ARTICLE

# **Prevalence and risk factors of trichomoniasis among women in Tabriz**

Abdol-Samad Mazloumi Gavgani<sup>1\*</sup>, Asiye Namazi<sup>1</sup>, Ardavan Ghazanchaei<sup>2</sup>, Sakineh Alizadeh<sup>3</sup>, Fahimeh Sehhati<sup>3</sup>, Sakineh Rostamzadeh<sup>1</sup>, Afsaneh Dolatkhah<sup>2</sup>

<sup>1</sup> Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>2</sup> Tropical and Infectious Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>3</sup> Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran

#### ABSTRACT

**Background**: We investigated frequency of trichomoniasis among non-pregnant women in health centers of Tabriz, Iran.

**Patients and methods**: 1000 non-pregnant women aged 15-49 in health centers of Tabriz, Iran were examined by wet smears and Diamond culture methods for Trichomonas vaginalis, during the period of March to September 2005.

**Results:** Among 1000 specimens 92 cases were revealed to be positive for Trichomonas vaginalis by culture method and 31 cases by wet smear method. There was no significant difference in the isolation rate of trichomoniasis in women according to age, occupation, husband education level, abortion, parity, menstrual status and contraception use. The difference in the isolation rate of trichomoniasis in women with marriage age of more than 18 years (10.9%) and in women with marriage age of less than 18 years (8%) was statistically significant. Infection rates in different education levels did not show statistically significant difference.

**Conclusion**: Trichomonas vaginalis is one of the important diseases with a high prevalence in women in Tabriz. Eradication of this disease is possible with extensive public health education and administration of specific therapeutic agents to the infected patients.

Keywords: Trichomoniasis, Prevalence, Iran. (Iranian Journal of Clinical Infectious Diseases 2008;3(2):67-71).

#### INTRODUCTION

Lower genital tract related complaints among women account for majority of outpatient women health care visits in the United States (1). Trichomonas vaginalis is a protozoan parasite transmitted almost exclusively through vaginal intercourse. It is the most common non viral sexually transmitted disease (STD) in humans (24). Infection with the organism can cause vaginitis in women and urethritis in men (5). Trichomoniasis also impacts upon birth outcomes and is a co-factor immunodeficiency in human virus (HIV) transmission and acquisition (6-8). Approximately 180 million women world wide and 3 million women in the United States are infected every year by T.vaginalis. Of note, up to 50% of infection may be asymptomatic (4, 9, 10).Diagnosis and elimination of the cause of the problem rely heavily on an accurate and thorough history and physical

Received: 22 May 2007 Accepted: 22 December 2007 Reprint or Correspondence: Abdol-Samad Mazloumi Gavgani. Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. E-mail: mazloumi@tbzmed.ac.ir

examination (11). Despite its limited sensitivity (1, 7), direct microscopic examination of the vaginal fluid remains the most widely utilized diagnostic test for this infection. Culture of the organism using vaginal specimens is the current "gold standard" (1,4,7,10). The purpose of this study was to describe the prevalence and common risk factors, signs and symptoms of trichomoniasis in a randomly selected population of women in Tabriz.

## **PATIENTS and METHODS**

This was a cross-sectional study that 1000 non pregnant women between the ages of 15 and 49 years who had  $\geq 1$  week untreated genital complaints were enrolled under a research protocol. Then written and verbal informed, consent was obtained. Women were enrolled at 3 sites. Inclusion criteria for enrollment included the presence of vaginal discharge, abnormal vaginal odor, vaginal itching, or lower genital tract burning sensation. An extensive questionnaire on current and past medical and personal, general, obstetric, sexual and contraceptive history was completed by each patient. Each woman also underwent a speculum examination that included direct observation for vaginal inflammation, cervicitis and evaluation of vaginal secretions for color, viscosity. Among discharges, odor was also assessed before and after the addition of 10% potassium hydroxide (KOH). During examination, vaginal specimens were collected using sterile cotton swabs. One of these was used to inoculate culture medium (Diamonds media) for diagnosing T.vaginalis. The tube was then incubated at 37°C for 8 days and observed microscopically every 2 days for the presence of motile trichomonads.

The second swab was used to direct observation of a wet mount to diagnose T.vaginalis by observing motile trichomonads. Data were registered in a database and analyzed with SPSS statistical software using Chi-square and Fisher's exact test if needed.

# RESULTS

A total of 1000 women were included in the study. The mean age of the patients was  $31.6\pm6.1$  years. The overall infection frequency diagnosed by culture method (the gold standard) among the 1000 women was 92 (9.2%) and the frequency of trichomoniasis by wet mount method was 31 (3.1%).

Symptoms varied among the 1000 women. The most frequent reported symptoms included malodor, low abdominal pain, itching, dysparonia, burning sensation and dysuria (Table 1). The infection rate of T.vaginalis among women involved in this study did not show any statistically significant differences with age (p>0.05). The high prevalence rate of infection (10.3%) was found in women of less than 20 years old group. The women with the marriage age over the 18 years showed the highest infection rate (10.9%). There was a significant difference in relation to the age of marriage (p<0.05).

**Table 1.** Frequency of symptoms in Trichomonasvaginalis infected patients

Patient complaint	Number (%)
Malodor	526 (52.6)
Dysuria	230 (23)
Urinal signs	180 (18)
Itching	360 (36)
Itching during intercourse*	231 (23.5)
Low abdominal pain	458 (45.8)
Burning during intercourse*	268 (27.1)
Dysparonia*	272 (27.6)

\* 16 women were not married and not included.

The infection rate in housewives (9.2%) was not significantly higher than the rate in employed women. Also overall prevalence of infection in women with moderate education level (under diploma) was 13.3% and the infection rate in women with high education level (academic education) was 6%. There were significant differences between low, moderate and high education levels, but among husband education levels there were not statistically significant differences.

Of the 741 women with negative history of abortion 68 (9.2%) were infected with T.vaginalis, while 24 (9.3%) of the 259 women with an abortion previously, had the infection. The difference between these two groups was not statistically significant (p>0.05). Furthermore, there was no statistically significant difference related to sexual intercourse (p>0.05), although T.vaginalis infection was more common in women with sexual intercourses more than 6 times per week.

T.vaginalis infection was more common among women using depomedroxy progesterone acetate (16.7%) and intrauterine devices (11.6%), than among those using oral contraceptive (6%). The difference between these groups was not statistically significant ((p>0.05).

### DISCUSSION

The prevalence estimates of trichomoniasis vary between populations studied, but range from 5-74% in women, with the highest rates reported in either sex from STI clinics and in other high risk populations (6,12,13).

The results of this study have shown that among women attending a health center in Tabriz, prevalence of trichomoniasis was 9.2% by culture and 3.1% by the wet mount method. Different studies in Iran showed that prevalence rate varied widely from 0.5 to 42% in different provinces due to the unreliability of detection methods (14-20). According to the minimum concentration of organisms required for a positive result, culture is more sensitive than wet mount preparation (21). Majority of the surveys revealed that the culture method was a useful tool for accurate diagnosis of trichomoniasis (14,16,18,22-24). This presents an important public health problem, which should be drawn to the attention of the public as well as health authorities.

The available data suggest that reproductive hormone levels may be partly responsible for higher prevalence of trichomoniasis in older women (25). In this study higher rate of isolation was found in women less than 20 years old. This is probably related to the higher level of sexual activity in these women and maybe also due to the transmission from their husbands. Other studies reported the same results (15,17,19,20) except one study in Bushehr which showed high prevalence rate in females more than 45 years old (26).

We found no statistically significant difference in the frequency of trichomoniasis in relation to occupation and level of husband education. Similar lack of significance was also reported in other studies (27,28).

It is concluded that Trichomonas vaginalis in women in Tabriz is one of the important diseases with a high prevalence, and that the eradication of this disease is possible with extensive public health education and administration of specific therapeutic agents to the infected patients.

The question remains about the prevalence rate of infection in men in Tabriz. A study of epidemiology and risk factors of trichomoniasis in men can clarify this question.

## ACKNOWLEDGEMENTS

Special thanks to the staff of health centers of Tabriz for their participation in the elaboration of the study protocol. This study was supported by a grant from Tabriz University of Medical Sciences.

#### **REFERENCES** =

1. Landers DV, Wiesenfeld HC, Heine P, Krohn MA, Hillier SHL. Predictive value of the clinical diagnosis of lower genital tract infection in women. Am J Obstet Gynecol 2004;190:1004-10.

2. Gilbert RO, Ella G, Beach DH, Klaessig S, Singh BN. Cytopathogenic effect of trichomonas vaginalis on human vaginal epithelial cells cultured in vitro. Infect Immun 2000;68:4200-206.

3. Dehaan MS, Anderson DG. The CDC 2002 Guidelines for the Treatment of Sexually Transmitted Diseases: Implications for Women's Health Care. JMWH 2003;48:96-104.

4. Mariani SM. Vaginal infections-- How to diagnose and treat them. Annual Meeting of the American Society of clinical pathology. 2003 Sep 18 - Sep 21 2003; New Orleans, Louisiana.

5. Cates WJR. Estimates of the incidence and prevalence of sexually transmitted diseases in the United States. American Social Health Association Panel. Sex Transm Dis 1999;26:52-57.

6. Swygard H, Sena AC, Hobbs MM, Cohen MS. Trichomoniasis: clinical manifestations, diagnosis and management. Sex Transm Infect 2004:80:91-95.

7. Lawing LF, Hedgas SR, Schwebke JR. Detection of trichomoniasis in vaginal and urine specimens from women by culture and PCR. J Clin Microbiol 2000;38:3585-88.

8. Mayta H, Gilman RH, Calderon MM, Gottlieb A. 18S ribosomal DNA-based PCR for diagnosis of Trichomonas vaginalis. J Clin Microbiol 2000;38:2683-87.

9. Sorvillo F, Smith L, Kerndt P, Ash L. Trichomonas vaginalis, HIV and African- Americans. Emerg Infect Dis 2001;7:927-32.

10. Sobel JD. Vaginitis. N Engl J Med 1997;337:1896-903.

11. Plourd DM. Practical guide to diagnosing and treating vaginitis. Med Gen Med 1999:1:2.

12. Anorlu RI, Fagbenro-Beyioku AF, Fagorala T. Prevalence of Trichomonas vaginalis in patients with vaginal discharge in Lagos, Nigeria. Nigerian Postgard Med J 2001;8:183-6.

13. El Seoud Sf, Abbas MM, Habib FS. Study of trichomoniasis among Egyptian male patients. J Egypt Soc Parasitol 1998;28:263-70.

14. Rasti S, Thagriri A, Behrashi M. Trichomoniasis in parturients referring to Shabihkhani Hospital in Kashan, 2001-2. Feyz Journal 2003;26:21-25. [In Persian]

15. Sharifi I, Khatami M, Tahmores Kermani E. Prevalence of Trichomonas vaginalis in women referred to Vali-Asr polyclinic and the health center number 3 in Sirjan city. Journal of Kerman University of Medical Sciences 1994;3:125-32. [In Persian]

16. Ziaee H, Rezaeian M. Study of trichomoniasis in women referring to gynecology centers of Sari and comparison of laboratory diagnostic methods. Journal of Mazandaran University of Medical Sciences 1998;19:34-40. [In Persian]

17. Shahbazi A, Fallah E, Safaian R. Infection rate of Trichomonas vaginalis in females referring to Tabriz and Basmeng health care centers, 1998-99. Journal of Shahid Beheshti University of Medical Sciences 2001;4:231-34. [In Persian]

18. Hazrati Tappeh KH, Mohammad Zadeh H, Mostaghim M, Fereidoni J, Mehri E. A comparative study on the sensitivity of two different diagnostic ways of Diamond culture and wet mount in Trichomonas vaginalis diagnosis and correlation between infection and clinical findings. Urmia Medical Journal 2004;1:7-13. [In Persian]

19. Moshfe AA, Hosseini S. Comparison of clinical and microscopic diagnosis of trichomoniasis referred to the Yasuj women clinic. Armaghane-Danesh 2004;33:71-82. [In Persian]

20. Sharbat Daran M, Shefaei SH, Samiei H, Haji Ahmadi M, Ramezan Pour R, Mersadi N, et al. Comparison of clinical presentations, wet smear, Papanicolaou smear with Dorset's culture for diagnosis of Trichomonas vaginalis in doubtful women to trichomoniasis. Journal of Babol University of Medical Sciences 2005;27:46-49. [In Persian]

21. Ohlemeyer CL, Hornberger LL, Lynch DA, Swierkosz EM. Diagnosis of Trichomonas vaginalis in adolescent females: in pouch TV culture versus wet mount microscopy. J Adolesc Health 1998;22:205-8.

22. Ameercheli M, Jahani MR, Motevalian SA. The study of Trichomonas vaginalis infection in pregnant and non-pregnant out-patients of two gynecologic clinics in Quazvin. Kowsar Medical Jurnal 1999;4:247-51. [In Persian]

23. Zangi Abadi M, Ghoreishi M, Khoushideh M, Roudbari M, Bahrami Sh. Survey of sensitivity of wet smear and Dorset medium in comparison with Diamond medium for diagnosis of Trichomonas vaginalis. Tabibe-Shargh 2002;3:141-47. [In Persian] 24. Memar Pour H, Maraghi SH, Shahabi S, Khazan H. Evaluation of the sensitivity of wet smear, Diamond medium and Giemsa stain in diagnosis of Trichomonas vaginalis. Hakim 1998;2:135-40. [In Persian]

25. Spinillo A, Bernuzzi AM, Cevini C, Gulminetti R, Luzi S, De Santolo A. The relationship of bacterial vaginosis, Candida and Trichomonas infection to symptomatic vaginitis in postmenoposal women attending a vaginitis clinic. Maturitas 1997;27:253-60.

26. Foulad Vand MA. Diagnosis of trichomoniasis by three parasitologic methods and evaluation of indirect fluorescent antibody (IFA) in Bushehr and Kangan ports. Iranian South Medical Journal 1997;1:23-29. [In Persian]

27. Mahdi NK, Gany ZH, Sharief M. Risk factors for vaginal trichomoniasis among women in Basra, Iraq. Eastern Med health J 2001;7:918-24.

28. Buv'e A, Weiss HA, Laga M, Van Dyck E, Musonda R, Zekeng L, et al. The epidemiology of trichomoniasis in women in four African cities. AIDS 2001;15:589-96.