

Genetic structure of *Plasmodium vivax* population assessed by sequence analysis of the merozoite surface protein 3 β gene

Abbas Shahbazi¹, Ahmad Raeisi², Mehdi Nateghpour³, Mehdi Mohebbi³, Mehdi Asmar⁴, Hossein Mirhendi^{3*}

¹Department of Parasitology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

²National Malaria Control Program, Center for Disease Management, Ministry of Health, Tehran, Iran

³Department of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

⁴Department of Parasitology, Pasteur Institute, Tehran, Iran

ABSTRACT

Background: The endemicity and transmission intensity levels of malaria are related to genetic diversity of the parasites. Merozoite surface protein 3 β (MSP3 β) is an important marker for assessing the polymorphic nature of *Plasmodium vivax* while it is also a vaccine candidate against the parasite.

Patients and methods: In this study we investigated the genetic structure of *P. vivax* population by sequence analysis of a polymorphic region of the *P. vivax* MSP3 β gene in isolates from Iran. Blood samples were collected from 100 patients with clinical symptoms. DNA was extracted and the target gene was amplified by polymerase chain reaction (PCR). The sequences of 17 samples were used for sequence analysis using nucleotide Blast search and ClustalW multiple alignment. Phylogenetic tree was derived to describe the geographical branching and relationships.

Results: A large number of nucleotide insertions and deletions were observed in the sequences of polymorphic region of PvMSP3 β gene that were not specific in each biotype. Single nucleotide polymorphism (SNP) was found extensively in the sequences. The phylogenetic analysis did not show any significant geographical branching.

Conclusion: The lack of any geographical branching and extensive polymorphism in MSP3 β gene of *P. vivax* isolates suggests that more investigations are needed to find a more suitable gene in order to develop a vaccine.

Keywords: *Plasmodium vivax*, Merozoite surface protein 3 β , Malaria, Vaccine.

(Iranian Journal of Clinical Infectious Diseases 2010;5(3):126-132).

INTRODUCTION

Although it is usually a non-lethal disease, *Vivax* malaria has major injurious effects on human well-being growth and on the economic progress at individual, family, community, and national levels because of its extended and recurrent infection (1).

At present, *Plasmodium vivax* causes an estimated 80-90 million cases each year and is the most widely distributed human malaria species in the world. In the Americas and Asia, *P. vivax* is the most prevalent malaria species (2). Lately, re-emerging of *vivax* malaria has occurred in many regions where it had been largely eradicated during the malaria control programs such as Korea, temperate provinces of China, and some former Soviet republics (3-6). Although, there are rare and

Received: 15 November 2009 Accepted: 20 June 2010

Reprint or Correspondence: Hossein Mirhendi, PhD.
Department of Medical Parasitology and Mycology, National
Institute of Health Research, Tehran University of Medical
Sciences, Tehran, 14155-6446, Iran.

E-mail: mirhendi@tums.ac.ir

sporadic reports of severe and fatal *P. vivax* infections (7), as resources for malaria control are currently often directed against the more lethal *P. falciparum* malaria, the hidden burden of *P. vivax* malaria is likely to become increasingly apparent.

For monitoring the distribution of drug resistance and for evaluation the performance of the vaccines under development, understanding of population structures of the parasite is important (8). It has been shown that different *P. vivax* isolates exhibit distinctive biological characteristics (e.g., relapse patterns or transmissibility to mosquitoes). This is a matter of fact even within a local population, which are often used to specify geographical strains and subspecies (9). Such a biological diversity is demonstrated by genotyping approach using various polymorphic genetic markers including the genes coding for circum sporozoite protein (CSP) (10), the apical membrane antigen 1 (AMA1) (11), the Duffy binding protein (DBP) (12), the merozoite surface protein 1 (MSP1) (13), and the merozoite surface protein 3 α (MSP3 α) (14-16) and merozoite surface protein 3 β (MSP3 β) (17). The *P. vivax* MSP3 α , β and γ constitute a protein family with the central Ala-rich region with an extensive coiled-coil structure (18, 19). The genes encoding these proteins are highly polymorphic (17-23). Sequences of MSP3 β in *P. vivax* isolates from various geographical regions of the world are radically divergent with the majority bearing large insertion/deletion mutations in the central Ala-rich region (21).

In regions where *P. vivax* and *P. falciparum* are coexistent, transmission of *P. vivax* has been become more prevalent. In Iran, more than 80% of all malaria cases causes by *P. vivax* (24). Today, transmission of *P. vivax* is almost located in southern and east-southern provinces of the country including Sistan and Baluchestan, Hormozgan, Kerman and Boushehr (24). As malaria studies are of practical importance in the development and utilization of control programs (8), and because of the great importance of MSP3 β gene as a genetic

polymorphic marker and a vaccine candidate (20, 21,17), we have described the polymorphic nature of MSP3 β gene based on the partial sequence analysis of the gene in *P. vivax* isolated from malarious areas of Iran. These data can enrich the databases obtained from other malarious countries.

PATIENTS and METHODS

For this descriptive study, a total of 100 positive-blood samples were collected from patients with clinical symptoms of malaria attending to Malaria clinics in the malarious areas of Sistan and Baluchestan, Hormozgan, Kerman, and Boushehr provinces. Under the supervision of Ethical Committee of Tehran University of Medical Sciences, sample collection was achieved after obtaining informed consent from each subject. Blood specimens were taken by experienced laboratory technicians with use of venipuncture or finger prick. Blood slides were stained with Giemsa and examined by well-experienced microscopists. When revealed positive, approximately 1000 μ l of venous blood was mixed with EDTA and stored in -20°C for further studies.

For DNA extraction and PCR amplification, QIAamp® DNA blood mini kit 50 (Qiagen, Germany) was used. All samples were rechecked by nested-PCR using plasmodium genus specific (primary PCR) and *P. vivax* and *P. falciparum* species-specific primers (nested PCR) (25) for reconfirming of primary microscopy diagnosis of the parasite.

The fragment was amplified through PCR by forward (5'- AACTTGAGAAACGGATG-3') and reverse (5'-TGCGAGTGTTCATGCG-3') specific primers (21), which amplify a small fragment of *P. vivax* MSP3 β (PvMSP3 β) by binding at positions 102-118 and 1943-1961 of corresponding region of PvMSP3 β in the Belem strain (GenBank accession number AF099662). The PCR was performed with an initial denaturation of 2.5min at 95°C, followed by 35

128 *MSP3 β gene of P.vivax*

cycles of 95°C for 30sec, 56°C for 30sec and 72°C for 2.5min (26).

Seventeen samples selected according to the previous study (17), were subjected to DNA sequencing. DNA sequencing of the target polymorphic DNA region was performed through Cinnagen Company (Tehran, Iran). The program BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) was used for pair wise sequence alignment and comparison with all similar sequences deposited in GenBank. The version 1.83 of ClustalW program (<http://www.ebi.ac.uk/Tools/clustalw2/index.html>) was used for multiple sequence alignment and comparison between obtained sequences and published relevant sequences of *P. vivax* including Bengal isolates (Accession number AY454084), Belem reference strain (Accession number AF099662) and Sali strain (Accession number AY454095). A phylogenetic tree was derived from the aligned nucleotides to describe geographical branching and relationships.

RESULTS

Totally, 17 PCR products corresponded to nucleotides 1780-2800 of the Belem reference strain (Accession number AF093584), 1-1030 of the Bangal strain (Accession number AY454084) and Sali strain (Accession number AY454095) were selected for sequencing. In our setting, associated gene sequenced part contained approximately 830-1025 bp (Figure 1A). A large number of nucleotide insertions and deletions were observed, of which none matched with previously described biotypes (17) (Figures 1B and 1C). In some cases the blocks of nucleotide insertions and deletions were respondent to the strains of other countries such as Bangal strain (Figure 1C). Single nucleotide polymorphism (SNP) was seen extensively in the sequences (Figures 1B and 1C). The phylogenetic analysis did not show any significant biotype or geographical specific branching (Figure 2). Only four sequences (isolates

619, 69, 63, and 604 from Boushehr province) were assigned to a separate group or specific branch in the phylogenetic tree (Figure 2). The range of homology score among the isolates through ClustalW program was between 41-97%. The average of the sequence homology score through ClustalW program was 72% (in the range of 45-82) between the isolates of Iran and Belem reference strain and 71% (in the range of 48-82) between Iran and Sali strain.




Figure 2. Phylogenetic tree of PvMSP-3 β gene sequences using 17 isolates from malarious areas of Iran. A significant geographical branching of the parasite populations is not seen in geographically separated isolates. Only four isolates (619, 69, 63, and 604) from similar geographical origin (Kangan district in Boushehr province) were grouped together in a separate branch

DISCUSSION

The endemicity and transmission intensity levels of malaria may have some relationship with genetic diversity of the malaria parasites (27). Diversity and multiplicity of *P. vivax* in hyper-endemic areas such as Papua New Guinea is high (14). In hypo-endemic areas such as Thailand and Iran, diversity and polymorphism of the parasite and its complex genetic structures has been reported (6,16,17,23,28).

In the present study we report high level of single nucleotide polymorphism (SNP) and several blocks of insertions and deletions among

38B1	-----TCTGCACGTTCTGCAT	16	A
39B1	-----TAANGCCCGTCTCTGCA-	16	
217B1	-----GGAGTGTCCTCC	13	
52B1	-----NATCTGCCGNTCCGCCA	16	
43B1	-----NGCCGATCTGN	11	
65B1	-----TTGATCGNGCTCAANCCGC	19	
620B1	-----AACTGTCT	9	
Bangal	-----GTCAATGAAAAACGTGTCT	19	
SalI	-----GTCAATGAAAAACGTGTCT	19	
610B1	-----TGAAACGTGTCT	12	
604B1	-----GAAAACTGTCT	12	
63B1	-----CTGTCT	6	
69B1	-----T	1	
619B1	-----TGAAACCTGTCT	12	
113B1	-----CTCANCATCCGTAG	15	
29B1	-----CACGACCACCCGTTT	15	
211B1	-----GAATGCCGNTTC	12	
612B1	-----TGAA--ACGTGTCT	12	
Belem	GAAATTGTAAACCTTAAAAATCCCAACTTGAGAAACGGATGGTCAATGAATAACCTGTCT	1800	
53B1	-----GAGAGNCACNTCTGTTG	17	
38B1	TTGGANATTAGAA--ACTTTAAAGGNGATGTAGANA-NTGCAGAAAAGNCTGCAAAAAA	452	B
39B1	TTGCTANTTTAGACAAANTNNAGAAGGGAAGGTAGAC--CTGCAGAAANNCN-GCAAAAAA	455	
217B1	ATGGAGAGTTTAA--AAAGTTAAAGATGAAGTAAAAA-AGGCAGAAAAGGCTGCAAAAAA	474	
52B1	ANGGAGAGTTTACA-AAAATTAAAGAGGGAAGTAGAAN-AGGCTGAAAAGGCTGCCAAAAA	457	
43B1	ATGGAGAGTTTCA-NAAGTTANAAGATNAAGTAGATA-AGGCTGACAAGGCTGCAAAAAA	427	
65B1	ATGGNGAGTTTCA-NAAGTTAAAGATGAAGTAGATA-AGGCTGAAAAGGCTGCAAAAAA	439	
620B1	-----AGTTAAAGATGAAGTAGATA-AGGCTGAAAAGGCTGCAAAAAA	419	
Bangal	-----AGTTAAAGATGAAGTAGATA-AGGCTGAAAAGGCTGCAAAAAA	429	
SalI	ATGGAC-----AGTTAGAAAAGTTAAAAATG-AAGAAAGAAAGACTGCAAAAAA	438	
610B1	ATGGAGAGTTTACA-AAAATTAAAGAGGGAAGTAGAAA-AGGCTGAAAAGGCTGCAAAAAA	443	
604B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-AGGCAGAAAAGGCTGCAAAAAA	440	
63B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-AGGCAGAAAAGGCTGCAAAAAA	431	
69B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-AGGCAGAAAAGGCTGCAAAAAA	437	
619B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-AGGCAGAAAAGGCTGCAAAAAA	440	
113B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-AGGCAGAAAAGGCTGCAAAAAA	442	
29B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-ATGCAGAAAAGGCTGCAAAAAA	445	
211B1	ATGAAGATTTACA-AAAATTAAAACTGAAGTAGAGA-ATGCAGAAAAGGCTGCAAAAAA	455	
612B1	ATGGAGATTTAGG-AAAATTAAAAAGTGAAGTAGAGA-ATGCAGAAAAGGCTGCAAAAAA	440	
Belem	TTGTAGAGTTTACA-AAAAGTTAAAGATGAAGTAGATA-A-----GGCTGCAAAAAA	2222	
53B1	TTCTACTTCTNNT--NACTNGAAGNNTNGACNANANACCNGGAGGNNAGCCTGCGNAGAA	453	
38B1	GAACTCNGNCTNGAAATTTAAACCGTGA-----AGACTNGCC	780	C
39B1	--AANTCGAN--TNAATTTNAAACCGGGA-----ANACTG--C	764	
217B1	TGAAAATAAAGAAA-TATCACAACCTAGAA-----AATGAAATAA	787	
52B1	TGAAGATAAAAAA-TATCAGATCTAGAA-----AACGAAATAT	771	
43B1	TGAGGATAAAGAAAATAACAGAACTAGGA-----GATATCTTTN	742	
65B1	TGAAGATAAAGAAA-TAACAGAACTAGGA-----GATATCTTAA	753	
620B1	TGAAGATAAAGAAA-TAACAGAACTAGGAGA-----TATCTTAA	733	
Bangal	TGAAGATAAAGAAA-TAACAGAACTAGGAGA-----TATCTTAA	743	
SalI	TGACGATAAAGAA-G-TATCACAACCTAGAAAGAAATAAAGAACTAGAACTATATTAA	773	
610B1	TGAAGATAAA-AAA-TATCAGATCTAGAAAACGAAATATCAGAACTAGAAATATCTTAA	777	
604B1	TGAAAGNTAAGAAA-A-TTAGAACTAGAA-----GNN-TCTTAA	750	
63B1	TGAAAGTAAAGAAA-AATTAGAACTAGAA-----GGTATCTTAA	745	
69B1	TGAAAGTAAAGAAA-A-TTAGAACTAGAA-----GGTATCTTAA	749	
619B1	TGAA--GTAAGAAA-A-TTAGAACTAGAA-----GGTATCTTAA	750	
113B1	TGAAAGTAAAGAAA-A-TTAGAACTAGAA-----GGTATCTTAA	755	
29B1	TGAAAGTAAAGAAA-AATTAGAACTAGAA-----GGTATCTTAA	760	
211B1	TGAAAGTAAAGAAA-AATTAGAACTAGAA-----GGTATCTTAA	769	
612B1	TGAAAGTAAAGAAA-CATCAGATCTAGAA-----GATATATTAA	775	
Belem	TGAAAATAAAGAAA-TATCACAACCTAGAAAATGAAATAACAAAACCTAGGAGATATATTAA	2557	
53B1	GGAAGGNAANGAAA-ANTTAGAACTAGAA-----GGNATCTTAA	753	

Figure 1. Partial nucleotide sequences alignment of *PvMSP-3β* gene in *P. vivax* isolated from malarious areas of Iran and corresponded part of the gene in Belem reference strain (Accession number AF093584), Bangal strain (Accession number AY454084) and SalI strain (Accession number AY454095). High level of single nucleotide polymorphism among the isolates and a block of deletion in sequence of the isolate 620 is noticeable in part B and large block of insertion is noticeable in isolate 610 correspond to nucleotide 2526-2547 of Belem strain and to nucleotide 742-763 of SalI strain in part C. None of the other isolates entailed this block.

nucleotide sequences of the *PvMSP3 β* locus in the Iranian isolates (Fig. 2). Such diversity has been reported previously in only two reports (21,22); therefore this is the first sequence-based study for the gene in Iran and the third one all over the world.

Malaria in Iran is hypo-endemic and transmission is seasonal and experiences a break during the winter (24), hence, generation and maintenance of considerable genetic diversity at the nucleotide sequences of the *PvMSP3 β* locus in malarious areas of the country is partially due to increased travel followed by the introductions of diverse parasite strains from neighboring endemic countries (6,24,29).

In our study the most homology in the *PvMSP3 β* locus was found between the isolates 619 and 63 (98%) both from Kangan district, and minimum homology was found between isolate 43 and 52 both from Minab district. Therefore, in case of this parasite, belonging to the same geographical origin can not necessarily cause a dramatic homology in genetic structure. A considerable number of insertions, deletions and SNP in the sequences was observed, but was locality or biotype specific variety (17) (Fig. 1B). Such a variety has been formerly reported in other studies (21,27). Existence of blocks of insertion and deletions means that the diversity of *PvMSP3 β* locus can be easily assessed by size polymorphism or restriction fragment length polymorphism (RFLP) (2), as we have already shown the utility of these methods for diversity assessment of this locus (17).

We compared our sequences with the sequences of Belem reference strain, Bangal strain and Sall strain. Maximum and minimum identity of the sequences with Belem reference strain were 85% and 45% corresponding to isolates number 619 and 53 from Kangan in Boushehr province and Minab in Hormozgan province, respectively. Comparison of sequences with Bangal strain and Sall strain

showed the maximum identity score of 95% and 84% (both from Kangan), respectively.

According to phylogenetic dendrogram, we found that there is not a significant geographical branching of the parasite populations (Fig. 2) and only four isolates (619,69,63 and 604) from similar geographical origin (all from Kangan district in Boushehr province) were grouped together in a separate branch. Up to now, attempts to reveal phylogenetic branching and relationships among the global *P. vivax* isolates failed to demonstrate any geographical structure of the parasite populations, and sequences obtained from distinct geographical areas cluster together in the phylogenetic trees (22). In the present study isolates 612 from Boushehr, 53 from Hormozgan and 38 from Sistan and Baluchestan with distance of more than 1000Km have clustered together (Fig. 2). However, it cannot be ignored that the similarities between our isolates and the isolates from other localities of the world suggests that some of geographic isolation may exist.

In conclusion, based on the high diversity and lack of adequate genetic data about the parasite, more extensive and comprehensive studies are needed to demonstrate polymorphic and population genetic aspects of the *P. vivax* in Iran and other geographical countries.

†

ACKNOWLEDGEMENT

This study was supported by Tehran University of Medical Sciences and the Center for Diseases Management in Iranian Ministry of Health. We thank the Malaria clinic health care workers in Sistan and Baluchestan, Hormozgan, Boushehr and Kerman provinces for their kind help during sample collection.

REFERENCES

1. Mendis K, Sina BJ, Marchesini P, Carter R. The neglected burden of *Plasmodium vivax* malaria. Am J Trop Med Hyg. 2001;64 (Suppl. I, II):97-106.

2. Souza-Neiras W, Melo LMS, Machado RLD. The genetic diversity of *Plasmodium vivax*; A review. Mem Inst Oswaldo Cruz, Rio de Janeiro. 2007;102(3):245-54.
3. Sleigh AC, Liu XL, Jackson S, Li P, Shang LY. Resurgence of vivax malaria in Henan Province, China. Bull World Health Organ. 1998;76:265-70.
4. Chai JY. Re-emerging *Plasmodium vivax* malaria in the Republic of Korea. Korean J Parasitol. 1999;37:129-43.
5. Sabatinelli G, Aliev S, Ejov M, Verbitsky V. The malaria epidemiological situation in Tajikistan. Parassitologia 2000;42 (suppl 1):148.
6. Leclerc MC, Menegon M, Cligny A, Noyer JL, Mammadov S, Aliyev N, et al. Genetic diversity of *Plasmodium vivax* isolates from Azerbaijan. Malaria J. 2004;3:40.
7. Beg A. Cerebral involvement in benign tertian malaria. Am J Trop Med Hyg. 2002;67:230-32.
8. Cui L, Escalante AA, Imwong M, Snounou G. The genetic diversity of *Plasmodium vivax* populations. Trends Parasitol 2003;19:220-26.
9. Li J, Collins WE, Wirtz RA, Rathore D, Lal A, McCutchan TF. Geographic subdivision of the range of the malaria parasite *Plasmodium vivax*. Emerg Infect Dis 2001;7:35-42.
10. Rosenberg R, Wirtz RA, Lanar DE, Sattabongkot J, Hall T, Waters AP, et al. Circumsporozoite protein heterogeneity in the human malaria parasite *Plasmodium vivax*. Science. 1989;245:973-76.
11. Figtree M, Pasay CJ, Slade R, Cheng Q, Cloonan N, Walker J, et al. *Plasmodium vivax* synonymous substitution frequencies, evolution and population structure deduced from diversity in AMA 1 and MSP 1 genes. Mol Biochem Parasitol. 2000;108:53-66.
12. Xainli J, Adams JH, King CL. The erythrocyte binding motif of *Plasmodium vivax* Duffy binding protein is highly polymorphic and functionally conserved in isolates from Papua New Guinea. Mol Biochem Parasitol. 2000;111:153-60.
13. Putaporntip C, Jongwutiwes S, Sakihama N, Ferreira MU, Kho WG, Kaneko A, et al. Mosaic organization and heterogeneity in frequency of allelic recombination of the *Plasmodium vivax* merozoite surface protein-1 locus. Proc Natl Acad Sci USA. 2002;99:16348-53.
14. Bruce MC, Galinski MR, Barnwell JW, Donnelly CA, Walmsley M, Alpers MP, et al. Genetic diversity and dynamics of *Plasmodium falciparum* and *P. vivax* populations in multiply infected children with asymptomatic malaria infections in Papua New Guinea. Parasitology. 2000;121:257-72.
15. Mueller I, Kaiok J, Reeder JC, Cortes A. The population structure of *Plasmodium falciparum* and *Plasmodium vivax* during an epidemic of malaria in the Eastern Highlands of Papua New Guinea. Am J Trop Med Hyg 2002;67:459-64.
16. Cui L, Mascorro CN, Fan Q, Rzomp KA, Khuntirat B, Zhou G, et al. Genetic diversity and multiple infections of *Plasmodium vivax* malaria in Western Thailand. Am J Trop Med Hyg. 2000;68:613-19.
17. Shahbazi A, Raeisi A, Nateghpour M, Mirhendi H, Mohebbali M, Asmar M. Diversity of merozoite surface protein-3 β gene of *Plasmodium vivax* isolates from Iran. Iranian J Publ Health. 2007;36(4):1-5.
18. Galinski MR, Corredor-Medina C, Pova M, Crosby J, Ingravall P, Barnwell JW. *Plasmodium vivax* merozoite surface protein-3 contains coiled-coil motifs in an alanine-rich central domain. Mol Biochem Parasitol. 1999;101:131-47.
19. Galinski MR, Ingravall P, Corredor-Medina C, Al-Khedery B, Pova M, Barnwell JW. *Plasmodium vivax* merozoite surface protein-3 β and -3 γ share structural similarities with *P. vivax* merozoite surface protein-3 α and define a new gene family. Mol Biochem Parasitol. 2001;115:41-53.
20. Rayner JC, Corredor V, Feldman D, Ingravall P, Iderabdullah F, Galinski MR, et al. Extensive polymorphism in the *Plasmodium vivax* merozoite surface coat protein MSP-3 α is limited to specific domains. Parasitology. 2002;125:393-405.
21. Rayner JC, Huber CS, Feldman D, Ingravall P, Galinski MR, Barnwell JW. *Plasmodium vivax* merozoite surface protein-3 β is radically polymorphic through mutation and large insertions and deletions. Infect Gen Evol. 2004;4:309-19.
22. Mascorro CN, Zhao K, Khuntirat B, Sattabongkot J, Yan G, Escalante AA, et al. Molecular evolution and intragenic recombination of the merozoite surface protein MSP-3 α from the malaria parasite *Plasmodium vivax* in Thailand. Parasitology. 2005;131:25-35.
23. Shahbazi A, Raeisi A, Nateghpour M, Mirhendi H, Mohebbali M, Asmar M. Polymorphism of merozoite surface protein-3 α gene of *Plasmodium vivax* in isolates of Iran. Iranian Journal of Parasitology. 2008;3(2):15-20.
24. Diseases Management Center, Ministry of Health, I.R. Iran. Annual Reports of Malaria. 2007.
25. Snounou G, Pinheiro L, Goncalves A, Fonseca L, Dias F, Brown KN, et al. The importance of sensitive

132 MSP3 β gene of *P.vivax*

detection of malaria parasites in the human and insect hosts in epidemiological studies, as shown by the analysis of field samples from Guinea Bissau. Trans Roy Soc Trop Med Hyg. 1993;87:649-53.

26. Altschul FS, Madden LT, Schäffer AA, Zhang J, Zhang Z, Miller W. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res. 1997;25:3389-402.

27. Yangl Z, Miao J, Huang Y, Li X, Putaporntip C, Jongwutiwes S, et al. Genetic structures of geographically distinct *Plasmodium vivax* populations assessed by PCR/RFLP analysis of the merozoite surface protein 3 β gene. Acta Trop. 2006;100(3):205–12.

28. Shahbazi A, Raeisi A, Nateghpour M, Mohebbali M, Asmar M, Mirhendi H. Partial sequence analysis of merozoite surface proteine-3 α gene in *Plasmodium vivax* isolates from malarious areas of Iran. Iranian Journal of Parasitology 2008;3(4):1-8.

29. Zakeri S, Djadid ND, Zeinali S. Sequence heterogeneity of the merozoite surface protein-1 gene (MSP-1) of *Plasmodium vivax* wild isolates in southeastern Iran. Acta Trop. 2003;88:91–97