

Detection of *H. pylori* infection and cagA strains seropositivity in adult dyspeptic patients in east Azerbaijan, northwest of Iran

Mohamadreza Bonyadi*, Zohreh Babaloo, Ebrahim Fattahi, Manochehr Khoshbaten, Fateme Abbasalizade, Shaindokht Poozesh

Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

ABSTRACT

Background: *Helicobacter pylori* (*H. Pylori*) bacteria colonize human stomach mucosa and may establish acute or chronic gastric inflammation. Cytotoxic-associated gene A (cag A) is associated with higher grades of gastric inflammation and carcinoma. In the present study we determine cag A seropositive strain in dyspeptic patients with *H. pylori* infection.

Patients and methods: Six hundred adult dyspeptic patients examined for anti *H. Pylori* and anti-cag A antibodies by enzyme-linked immunoassay (ELISA) method. All cases resided in east Azarbijan in northwest of Iran and were enrolled in a 5-year period (2003–2008).

Results: A total of 85.5% of dyspeptic patients were positive for *H. pylori* infection. Anti-cag A antibody was detected in 35.6% of patients with *H. pylori* infection.

Conclusion: Screening of *H. pylori* infection by ELISA method revealed that the vast majority of (85.5%) dyspeptic patients are seropositive for *H. pylori*. Determining of photogenic strains of *H. pylori* by anti-cag A antibody could be diagnostic in severe gastric infections.

Keywords: *Helicobacter pylori*; Cytotoxic-associated gene A (Cag A); Gastric inflammation.
(Iranian Journal of Clinical Infectious Diseases 2010;5(4):228-230).

INTRODUCTION

Helicobacter pylori (*H. pylori*) colonizes human stomach mucosa where it induces acute or chronic gastric inflammation. *H. pylori* infection increases the risk of peptic ulcers, stomach adenocarcinoma and lymphoproliferative disease of stomach mucosa (1,2).

H. pylori heterogenic are associated with a variety of clinical manifestations. Cytotoxic associated gene A (cag-A), one of the critical

pathogenic genes of *H. pylori*, has close relationship with peptic ulcers. This gene encodes a toxic vocoulizing protein, Vag A, in about 50% of *H. pylori* infections (1,2). P₃₀ and P₃₃ are the other pathogenic genes which participate in *H. pylori* pathogenicity. Cag A gene's length is 40 KD and most of the ulcerogenic strains of *H. pylori* have this gene which increases the risk of peptic ulcers, gastric atrophia and gastric adenocarcinoma (3).

In the present study, cag-A positive *H. pylori* strains have been evaluated in adult dyspeptic patients.

Received: 13 July 2010 Accepted: 17 October 2010

Reprint or Correspondence: Mohamadreza Bonyadi
Department of Immunology, Drug Applied Research Center,
Tabriz University of Medical Sciences, Tabriz, Iran.

E-mail: bonyadir@tbzmed.ac.ir

PATIENTS and METHODS

Six hundred dyspeptic patients for whom gastric disorder has been confirmed by clinical examinations and endoscopy, enrolled in this study during a five year period (2003-2008). Anti-*H. pylori* IgG and anti-cag A antibody were analyzed by ELISA method (Padten and Diapro Kits) using six standards: 0, 15, 30, 60 and 100 unit (arbu/ml).

Data were entered and analyzed using SPSS software (version 14, SPSS Inc., Chicago, USA).

RESULTS

Six hundred dyspeptic adults (280 females and 320 males) aged 25-75 years old (mean: 43.2 ± 12.1 years) and 200 controls with no signs of dyspepsia (mean: 42.1 ± 15.2) were enrolled.

Anti-*H. pylori* IgG was positive in 85.5% (n=513) of patients and 81% (n=162) of controls. Totally, anti-cag A was positive in 30.5% of patients and 28% of controls. The difference did not reach a statistically significant level. Meanwhile, cag A-IgG was positive in 183 cases (30.5%) and 56 controls (28%).

Of 513 *H. pylori*-positive patients, 183 (35.6%) were also positive for cag A-IgG, however, this figure was 34.6% (56 of 162) in controls.

All *H. pylori* seronegative patients and controls were negative for cag A.

DISCUSSION

Frequency of cag A positive *H. pylori* strains was 35.6% and 34.6% among cases and controls, respectively.

Invasive and noninvasive methods have been developed for diagnosis of *H. pylori* infection. Serologic tests are among the best noninvasive methods. Because of the long half-life of *H. pylori* IgG it could be a good indicator of infection severity. Serologic detection of cag A could be very important in prognosis of *H. pylori* infection (4). Cag A strain is a pathologic marker that is

associated with strong immunologic response (5,6), hence, detection of this gene not only in *H. pylori* infected patients but also in asymptomatic cases, plays a critical role in treatment follow up and prevention of gastric ulcers and carcinomas. Frequency of cag A seropositivity has been variable in different societies: 50% in Turkish population (2), 40.2% in Iranian GERD patients and 43% in normal cases (7). In a study in Italy, frequency of cag A seropositivity was 86.1% in patients with duodenal ulcers, 96.4% in peptic ulcers and 52.4% in normal controls (8). In a study from China, *H. pylori* IgG in patients and controls was 88.9% and 45%, and cag A seropositivity was 78.1% and 31.3%, respectively (9).

European and American studies showed that 50% of *H. pylori* strains contain vag A gene and cag A (1).

Meta analysis of 21 study reports showed that specificity and sensitivity of ELISA testes were 79% and 85%, respectively (2). New commercial kits have higher sensitivity and specificity (98%) therefore, they could be more suitable to diagnose *H. pylori* infection. On the other hand, detection of pathogenic strains, cag A and vag A, with ELISA method and other genes which are associated with *H. pylori* pathogenicity, P₁₂₀ (cag A), P₃₀ (OMP) and P₃₃, could be helpful.

ACKNOWLEDGEMENT

We wish to thank our colleagues in Drug Applied Research Centre, Liver and Gastrointestinal Disease Research Center in Tabriz University of Medical Sciences.

REFERENCES

1. Lamarque D, Peek Jr RM. Pathogenesis of *Helicobacter pylori* infection. *Helicobacter*. 2003;8(suppl. 1):21-30.
2. Nazime OY, Ahmet S, Ali K, Iikay S. Detection of *H. pylori* infection by ELISA and Western blot techniques and evaluation of anti cagA seropositivity in adult

Turkish dyspeptic patients. World Gastroenterol 2006; 12(33):5375-78.

3. Sicinschi LA, Correa P, Bravo LE, Schneider BG. A positive assay for identification of *cagA* negative strains *Helicobacter pylori*. J Microbial Methods 2003; 55:625-33.

4. Peters TM, Owen RJ, Slater E, Varea R, Teare E, Saverymutu S. Genetic diversity in the *Helicobacter pylori* *cag* pathogenicity island and effect on expression of anti-*cagA* serum antibody in UK patient with dyspepsia. J Clin Pathol. 2001;54:219-23.

5. Herbrink P, van Doorn LJ. Serological methods for diagnosis of *Helicobacter pylori* infection and monitoring of eradication therapy. Eur J Clin Microbiol Infect Dis. 2000;19:164-73.

6. Monteiro L, Bergey B, Gras N, Megraud F. Evaluation of the performance of the Helico Blot 2.1 as a tool to investigate the virulence properties of *Helicobacter pylori*. Clin Microbiol Infect. 2002;8:676-79.

7. Somi MH, Fattahi E, Fouladi R, Karimi M, Bonyadi R, Baballou Z. An inverse relation between *CagA*+ strains of *Helicobacter pylori* infection and risk of erosive GERD. Saudi Medical Journal. 2008; 29(3):393-96.

8. Orsini B, Ciano G, Surrenti E, Macrí G, Biagini MR, Milani S, et al. Serologic detection of *cagA* positive *Helicobacter pylori* infection in a northern Italian population. Helicobacter. 1998;3(1):15-20.

9. Abraham MY, Guillermo I, Lee J. Relation between *Helicobacter pylori* *cagA* status and risk of peptic ulcer disease. Chinese Medical Journal 2004;117(2):301-2.