Variable Number Tandem Repeat (VNTR) Genotyping of Hydatidiform Mole in Iranian Patients

Zahra Pakzad¹, Hossein Mozdarani^{1*}, Narges Izadi-Mood², and Shirin Niromanesh²

1. Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran 2. Mirza Koochak Khan Hospital, Iran University of Medical Sciences, Tehran, Iran

Abstract

Background: Classification of molar gestation into Complete Hydatidiform Mole (CHM) and Partial Hydatidiform Mole (PHM) is done according to clinical, ultrasonographic, histologic and genetic criteria. However, making a distinction between CHM and PHM using histologic criteria alone may be difficult and several studies have shown that misclassifications are frequent, even for experienced pathologists. CHM is the most common precursor to choriocarcinoma and heterozygous moles carry an increased predisposition to transformation.

Methods: Formalin-fixed, paraffin-embedded tissue sections of patients as well as peripheral blood of patients and their partners' were collected in EDTA tubes. Tissue samples were obtained by curettage. Histological evaluation was performed on routine section stained with Hematoxylin and Eosin. Variable Number Tandem Repeats (VNTRs) genotyping was performed for 30 cases in two groups of CHM (n=21) and PHM (n=9), with Polymerase Chain Reaction (PCR) amplification of 2 different polymorphic loci, namely the Col2A1 and D1S80.

Results: The results of DNA analysis by VNTR genotyping showed that in 16 cases of CHM, amplification of the VNTR polymorphic loci showed androgenetic mono-spermic moles (homozygote) and in 5 cases of CHM androgenetic dispermic moles (heterozygote) in molar tissue. In cases of PHM, 6 samples were triploid dispermic and 3 samples were diploid biparental.

Conclusion: This study confirmed that VNTR genotyping can identify the parental source of polymorphic alleles in hydatidiform mole. Compared to STR genotyping, VNTR genotyping was performed by PCR amplification of several minisatellite markers of DNA. This method significantly requires less time and is cost-effective.

Avicenna J Med Biotech 2014; 6(4): 246-253

Keywords: Hydatidiform mole, Minisatellite, Variable Number Tandem Repeats, Genotyping technique

Introduction

Hydatidiform Mole (HM) (OMIM 231090) is an abnormal gestation characterized by significant hydropic enlargement and variable trophoblastic proliferation involving part of or all chorionic villi. HMs are sub-classified into Complete HM (CHM) and Partial HM (PHM), based on clinical features, histopathology and genetic differences ¹⁻³.

HMs are relatively common and occur in approximately 1 in 1500 pregnancies in Europe and North America. This incidence varies between ethnic groups and is 2-10 times

* Corresponding author: Hossein Mozdarani, Ph.D., Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran Tel: +98 21 82883830 Fax: +98 21 82806544 E-mail: mozdarah@modares.ac.ir Received: 17 Mar 2014 Accepted: 3 Jun 2014



higher in some countries of Latin America, the Middle East, and the Far East with the highest frequencies being in Mexico, Iran and Indonesia⁴⁻⁷.

Karyotype and genotype analyses have shown that sporadic moles may have different genotypic types with the majority of the CHMs being diploid androgenetic and the majority of the PHMs being triploid diandric dispermic. Among androgenetic moles, the majority are mono-spermic (homozygous) and 10-20% are dispermic (heterozygous)⁸⁻¹². Studies have suggested that heterozygous mole may have a more malignant potential than its homozygous counterpart ^{13,14}.

With the increased use of high resolution ultrasonography in early pregnancy, the diagnosis and sub-classification of hydatidiform moles have become more difficult, as CHMs are now commonly evacuated at an earlier stage and before the development of classical morphologic and histologic features. Recently, pathologists have relied on molecular techniques, such as DNA flow cytometry, chromosome *in situ* hybridization, polymerase chain reaction-based genotyping or HLA typing, showing DNA content differences for CHM and PHM ¹⁵⁻¹⁷. Approximately 75% of clinically ascertained molar pregnancy are CHMs and mostly of diploid, androgenetic origin, and the remaining 25% are PHMs and mostly of diandric, triploid origin ^{18,19}. Vanda Repiska et al suggested that PCR amplification of polymorphic VNTR regions will be a useful tool in helping to determine relative risk of choriocarcinoma in patients with complete hydatidiform moles. Closer clinical surveillance of patients with heterozygous moles may prove to be an important part of management in the future ²⁰.

Persistent gestational trophoblastic disease develops after a CHM in 10 to 30% of cases and after a PHM in 0.5 to 5%. Choriocarcinoma arises in up to 3% of CHM and is rare but reported to be one of the consequences of PHM ^{21,22}. Despite the difference in risk for persistent disease or metastases, the post evacuation surveillance recommendations are

identical with frequent monitoring of serum human chorionic gonadotropin levels until normalization and monthly monitoring for 6 month thereafter ^{23,24}.

Each human somatic cell contains 6.4 billion base pairs of DNA. Portions of this DNA encode over 30,000 genes, while the remainders are non-coding. The human genome contains hundreds of highly polymorphic segments, each of which is characterized by allelic variation and attributed to the difference in the copy number of the tandemly repeated DNA sequences, such as Variable Number of Tandem Repeats (VNTRs). There are two principal families of VNTRs: minisatellite and microsatellite. Minisatellites are composed of different repeated "core" DNA sequences arranged in tandem. The size of the core sequence can vary from 7 to 100 bp in different VNTRs. Due to their high polymorphic content, VNTR is a useful tool in linkage analysis, forensic identification, paternity testing and population genetic studies ²⁵⁻²⁷. VNTRs or minisatellite loci are polymorphic and due to the variation in the number of copies of their repeat unit, VNTRs are multiallelic, so they display variation in length of the repeats and don't vary from one generation to the next. They are easily amenable to Polymerase Chain Reaction (PCR). Regarding the above mentioned criteria, these markers would be useful and informative in studying genetic variation among human population²⁸. In a study performed in Iran, it was shown that both ApoB and D1S80 are very polymorphic in Iranian populations ²⁶. The Col2A1 minisatellite as one of VNTR markers, located at 3' of last exon of Col2A1 gene has been shown to be highly variable and informative 29

Therefore, the aim of this study was to show the value of VNTR amplification for genotyping of CHM and PHM.

Materials and Methods

Patients

Thirty cases were selected from gynecologic oncology service. Formalin-fixed, paraffin-



embedded tissue sections of the patients were collected from the routine gynecologic pathology service files of the Mirza Koochak Khan Hospital, a referral gynecologic hospital in Tehran, Iran. Before sample collection, the study was reviewed and approved by the Tarbiat Modares University Ethics Committee.

The mean age of patients with complete mole was 26.4 years (range, 15-41 years) and in patients with partial mole, it was 30 years (range, 24-37 years). Peripheral blood of patients and their partners' were collected in EDTA tubes. Tissue samples were obtained by curettage. Histological evaluation was performed on routine section stained with Hematoxylin and Eosin and diagnoses were made by gynecologic pathologist.

For each specimen, the maternal deciduas and molar villi on paraffin block were separated using scalpel and a consecutive section stained with Hematoxyline and Eosin (Figure 1). Then three, 20 μm thick sections from FFPE tissue blocks were collected in 1.5 *micro* tubes for each case.

DNA isolation

DNA was isolated from FFPE tissue as well as from peripheral blood of patients and their partners. DNA extraction from peripheral blood was performed using modified salting out method. For each extraction, 500 μl blood (EDTA, whole blood) was used. Then, 1000 μl cell lysis buffer was added into the sterile 1.5 *ml* tube and centrifuged at 7000 RPM for 4 *min* and the supernatant was discarded. This step was repeated once more. To



Figure 1. H&E staining of hydatidiform mole. A) CHM characterized by, 1: hydropic villi with circumferential trophoblastic hyperplasia $\times 100$ B) PHM characterized by, 1: hydropic and 2: fibrotic villi with focal mild trophoblastic hyperplasia $\times 100$

the remaining white pellet, 300 µl nuclei lysis buffer was added and incubated at RT for 30 min until the pellet was re-suspended. Then, 100 µl NaCl (Merck, Germany) 6M and 600 μl choloroform (Merck, Germany) were added and centrifuged at 7000 RPM for 4 min. The supernatant (upper layer) was transferred to a new 1.5 tube and 1000 μl cold Ethanol (Merck, Germany) 100% (-20°C) was added. The tube was inverted 5-6 times gently until Hank of DNA was obtained, then centrifuged at 1000 RPM for 2 min. The supernatant was discarded and cold Ethanol (Merck, Germany) 70% (-20°C) was added, again tube was centrifuged. The supernatant was discarded and 100 μl distilled water was added to resuspended DNA.

DNA was prepared from tissue using modified Phenol-chloroform method. In each case, three, 20 µm-thick sections from FFPE tissue blocks were collected in 1.5 micro tubes. Briefly, the tissue sections were deparaffinized with Xylene (Merck, Germany), washed with 100%, 96% and 70% Ethanol (Merck, Germany) and dried in thermo block. The dried tissue was then incubated in 200 *µl* tissue lysis buffer (Genetbio, Korea) and 50 μl proteinase K (Genetbio, Korea). After overnight incubation at $46^{\circ}C$, the tube was incubated at $75^{\circ}C$ for 1 hr until complete digestion of tissue. Then, 250 µl Phenol-Choloroform-Isoamyl alcohol (Sinaclon, Iran) was added and centrifuged. The upper layer was transferred to a new 1.5 tube and equal volume of Isopropanol alcohol (Merck, Germany) was added and then incubated at $4^{\circ}C$ overnight. The next day, the tube was inverted several times and then centrifuged at 13000 RPM for 30 min. The supernatant was discarded and 50 µl distilled water was added to eluted DNA.

PCR amplification

The DNA samples were subjected to polymerase chain reaction (PCR) amplification of 2 VNTRs (variable number tandem repeats) polymorphic loci, namely the Col2A and D1S80. The primer sequences for 2VNTR locus ²⁹⁻³¹ are given in table 1.

Table 1. Minisatellite primers

Locus	Location	Repeat	Primer sequence		
D1S80	1P	16 bp	5'-GTC TTG TTG GAG ATG CAC GTG CCC CTT GC-3' 5'-GAA ACT GGC CTC CAA ACA CTG CCC GCC GCC G-3'		
Col2A1	12q13	26 bp	5'-GCT TCC TCC AGG TTA AGG TTG ACA G-3' 5'-ACC TGC TCT CCT CCG ACC C-3'		

All the DNA samples were subjected to PCR with slight modification in a thermocycler (CreaCon- Netherlands). To determine the genetic origin of each HM, 1 μl DNA from the patient and her partner and 1.5 μl DNA from each of the HM were amplified using two pairs of primers (BioNEER, South Korea) in a final reaction volume of 25 μl containing DNA, 2× Taq Master mix (Amplicon, Denmark) 12.5 μl and 50 *pmols* of each primer.

The cycle conditions were as follows: for Col2A1, 30 cycles of denaturation at $94^{\circ}C$ for 30 s; annealing at $60^{\circ}C$ for 50 s; extension at $72^{\circ}C$ for 50 s, for D1S80, 30 cycles of denaturation at $94^{\circ}C$ for 30 s; annealing at $70^{\circ}C$ for 50 s; extension at $72^{\circ}C$ for 50 s. The initial denaturation and final extension were carried out at $94^{\circ}C$ for 5 *min* and $72^{\circ}C$ for 10 *min* for Col2A1 and D1S80.

Following amplification, 8 μl of each PCR reaction product was analyzed by electrophoresis in a 2% agarose gel with a 100 *bp* molecular weight marker, stained with ethidium bromide and documented in a gel documentation system. In the interpretation of genotyping results, the allele sizes of minisatellite polymorphisms found in the molar villi were



Figure 2. Results of the PCR amplification of D1S80 VNTR locus, line 1: mother; line 2: CHM of heterozygous; line 3: father; line 4: mother; line 5: CHM of homozygous; line 6: father; line 7: ladder; line 8: blank

then compared with the corresponding allele sizes observed in the maternal and paternal samples as shown in figures 2-4.

Results

The results of DNA analysis by VNTR genotyping as well as other related information about patients including clinical pa-



Figure 3. Results of the PCR amplification of D1S80 VNTR locus, line 1: mother; line 2: PHM of triploid dispermic; line 3: father; line 4: ladder

1 Mother	2 Mole diploid biparental	3 Father	4 Ladder	
				500 <i>bp</i>
See.				

Figure 4. Results of the PCR amplification of Col2A1 VNTR locus, line 1: mother; line 2: PHM of diploid biparental; line 3: father; line 4: ladder

Avicenna Journal of Medical Biotechnology, Vol. 6, No. 4, October-December 2014

VNTR Genotyping of Hydatidiform Mole

Patient no.	Patient age(yrs)	Weeks of gestation	Pathologic diagnosis	B-hCG (mlU/ml)	Genotyping
1	29	6W	СНМ	200000	A.mono S
2	24	6W+6d	CHM	120000	A.mono S
3	30	12W+3d	CHM	140000	A.mono S
4	27	7W+4d	CHM	100000	A.mono S
5	27	7W	CHM	98000	A.di S
6	20	9W	CHM	20700	A.di S
7	20	10W	CHM	40000	A.mono S
8	41	11W+5d	CHM	140000	A.mono S
9	29	9W+4d	CHM	14800	A.mono S
10	26	8W+5d	CHM	150000	A.mono S
11	27	10W+5d	CHM	180000	A.mono S
12	29	8W+1d	CHM	110000	A.mono S
13	22	11W+1d	CHM	150000	A.di S
14	23	11W+4d	CHM	140000	A.di S
15	38	10W+2d	СНМ	120000	A.mono S
16	20	9W+6d	СНМ	200000	A.mono S
17	22	11w+3d	СНМ	20400	A.mono S
18	28	11W	CHM	300000	A.di S
19	28	9W+5d	СНМ	273000	A.mono S
20	31	11W+6d	СНМ	30000	A.mono S
21	15	12W	СНМ	180000	A.mono S
22	33	7W+4d	РНМ	225	Diploid biparental
23	30	9W+2d	РНМ	1600	Triploid di S
24	31	8w+5d	РНМ	1608	Diploid biparental
25	37	10W+3d	РНМ	2000	Triploid di S
26	25	8W	PHM	1624	Triploid di S
27	35	11W+2d	PHM	1200	Triploid di S
28	30	9W	PHM	1524	Diploid biparental
29	31	10W+3d	PHM	17000	Triploid di S
30	24	11W	PHM	2650	Triploid di S

Table 2. Clinical parameter, histological diagnosis and genotyping findings of patients examined

rameter and histologic diagnosis are shown in table 2. In 16 cases of CHM, amplification of the VNTR polymorphic loci showed androgenetic mono-spermic moles (homozygote) in molar tissue (76.2%). In 5 cases of CHM, androgenetic dispermic moles (heterozygote) were seen in molar tissue (23.8%). In cases of PHM, 6 samples were triploid dispermic (66.7%) and 3 samples were diploid biparental (33.3%).

Discussion

CHMs are generally diploid and androgenetic in origin, all 46 chromosomes being derived from the father. They may be monospermic, arising by fertilization of an enucleate egg by a single spermatozoon which then doubles to provide a diploid chromosome complement, or dispermic arising from fertilization of an enucleate egg by two spermatozoa. PHMs are usually triploid, the extra haploid set of chromosome can have either a maternal origin (digynic triploidy) or a paternal one (diandric triploidy)^{14,15}.

Our results showed that 76.2% cases of CHM were homozygous and 23.8% heterozygous. Also, in cases of PHM, 66.7% were triploid dispermic and 33.3% diploid biparental. These results are consistent with reports in other investigations ^{8,11}. Persistent gestational trophoblastic disease develops after a CHM in 10 to 30% of cases and after a PHM in 0.5 to 5% ^{21,22}. Homozy-gous and heterozygous CHM are two genetically distinct entities which can only be distinguished on the basis of genetic analysis and heterozygous mole may have a more malignant potential than its homozygous counterpart ^{16,17}. In this study, five of 21 CHM cases, analyzed by genotyping, were found to be heterozygous.

Although many cases of hydropic abortion can be diagnosed on the basis of histology alone, it is difficult to estimate the proportion of cases evacuated at an early gestational age ^{1,2,15}. Thus, accurate diagnosis of abnormal product of conception specimens and subsequent classification as complete or partial mole is mandatory to ascertain the actual risk of persistent GTD and to determine the appropriate nature and duration of clinical follow-up care. More advanced molecular genetics techniques have subsequently been applied to resolve this differential diagnosis problem ¹⁵⁻¹⁷.

The PCR amplification of polymorphic VNTR region will be a useful tool in helping to determine relative risk of choriocarcinoma in patients with complete hydatidiform moles ²⁰. VNTRs are the most informative markers for genetic studies. Although each VNTR locus is usually associated with a large number of different alleles in any population, each individual carries two alleles, one on each of the two homologous chromosomes. The major methodological advantage of VNTR is that all alleles are stable across generation, they are inherited in a Mendelian fashion and do not vary from parents to offspring ^{25,27}.

Other techniques that can be used for genetic evaluation of molar pregnancy include Fluorescence In Situ Hybridization (FISH), flow cytometry and fluorescent microsatellite genotyping. However, comparison of VNTR genotyping with other methods showed that VNTR method is very simple, thus does not require special device and also it is less timeconsuming and more cost-effective.

In this study, it was shown that VNTR genotyping can identify the parental source of polymorphic alleles and thus discern androgenetic diploidy (homozygous and heterozygous), diandric triploidy, and biparental diploidy, which allows for specific diagnosis of CHMs, PHMs, and NMs, respectively. In a similar study, choriocarcinoma analyzed by PCR amplification of polymorphic VNTR in 9 out of 10 cases was identified as herterozygous²⁰.

The reliability of the diagnosis is crucial for appropriate counseling and to determine if a patient falls in to a "short-term" or "longterm" follow-up. In some instances, for example in women having a "lesion suspicious for HM", the diagnosis could be confirmed or ruled out, thus avoiding an unnecessary follow-up. This may be of particular importance in "older patients" having difficulties in conceiving and for whom a one-year wait may be extremely distressing ^{9,13,16,24,32}.

Conclusion

In conclusion, the majority of cases of hydropic abortion can be diagnosed based on histological feature alone, but molecular biological techniques will be necessary to evaluate the proportion of cases evacuated at an early gestational age. Our results indicate that VNTR genotyping, a relatively simple, reliable, cost-efficient and rapid procedure, can be used successfully for diagnosis of HM and distinguishing CHM from its mimics, which is clinically important because the majority of persistent trophoblastic diseases and choriocarcinomas arise from CHM.

Acknowledgement

This work was supported by Research Department of the Faculty of Medical Sciences, Tarbiat Modares University. The authors would sincerely thank for contribution of all participating individuals, especially patients in this study.

References

- 1. Landolsia H, Missaouia N, Brahemc S, Hmissaa S, Gribaac M, Yacoubia MT. The usefulness of p57 KIP2 immunohistochemical staining and genotyping test in the diagnosis of the hydatidiform mole. Patho Res Pract 2011;207(8):498-504.
- Muhlstein J, Golfier F, Rittore C, Hajri T, Philibert L, Abel F, et al. The spectrum of NLRP7 mutations in French patients with recurrent hydatidiform mole. Eur J Obstet Gynecol Reprod Biol 2011;157 (2):197-199.
- Qian JH, Cheng Qi, Murdoch Sh, Xu CM, Jin F, Chebaro W, et al. The genetics of recurrent hydatidiform moles in China: correlations between NLR P7 mutations, molar genotypes and reproductive outcomes. Mol Hum Reprod 2011;17(10):612-619.
- Slim R, Mehio A. The genetics of hydatidiform moles: new lights on an ancient disease. Clin Genet 2007;71:25-34.
- 5. Fallahian M. Familial gestational trophoblastic disease. Placenta 2003;24(7):797-799.
- Javey H, Sajadi H. Hydatidiform mole in southern Iran: a statistical survey of 113 cases. Int J Gynaecol Obstet 1978;15(5):390-395.
- Slim R, Coullin Ph, Diatta AL, Chebaro W, Courtin D, Abdelhak S, and Garcia A. NLRP7 and the genetics of post-molar choriocarcinomas in Senegal. Mol Hum Reprod 2012;18(1):52-56.
- 8. Slim R, Wallace EP. NLRP7 and the genetics of hydatidiform moles: recent advances and new challenges. Front Immunol 2013;4:1-12.
- 9. Baasanjav B, Usui H, Kihara M, Kaku H, Nakada E, Tate S, et al. The risk of post-molar gestational trophoblastic neoplasia is higher in heterozygous than in homozygous complete hydatidiform moles. Hum Reprod 2010;25(5):1183-1191.
- Lai CY, Chan KY, Khoo US, Ngan HY, Xue WC, Chiu PM, et al. Analysis of gestational trophoblastic disease by genotyping and chromosome in situ hybridization. Mod Pathol 2004;17(1):40-48.
- 11. Devriendt K. Hydatidiform mole and triploidy: the role of genomic imprinting in placental development. Human Reprod Update 2005;11:137-142.
- Fukunaga M. Early partial hydatidiform mole: prevalence, histopathology, DNA ploidy, and persistence rate. Virchows Arch 2000;437(2):180-184.
- Wake N, Seki T, Fujita H, Okubo H, Sakai K, Okuyama K, et al. Malignant potential of homozygous and heterozygous complete moles. Cancer Res 1984;44(3):1226-1230.
- 14. Fisher RA, Lawler SD, Povey S, Bagshawe KD. Genetically homozygous choriocarcinoma follow-

ing pregnancy with hydatidiform mole. Br J Cancer 1988;58(6):788-792.

- 15. Sarmadi S, Izadi-Mood N, Abbasi A, Sanii S. p57 KIP2 immunohistochemical expression: a useful diagnostic tool in discrimination between complete hydatidiform mole and its mimics. Arch Gynecol Obstet 2011;283(4):743-748.
- Petignat P, Billieux MH, Blouin JL, Dahoun S, Vassilakos P. Is genetic analysis useful in the routine management of hydatidiform mole? Hum Reprod 2003;18(2):243-249.
- 17. Howat AJ, Beck S, Fox H, Harris SC, Hill AS, Nicholson CM, et al. Can histopathologists reliably diagnose molar pregnancy? J Clin Pathol 1993;46 (7):599-602.
- Ulker V, Gurkan H, Tozkir H, Karaman V, Ozgur H, Numanoglu C, et al. Novel NLRP7 mutations in familial recurrent hydatidiform mole: are NLRP7 mutations a risk for recurrent reproductive wastage? Eur J Obstet Gynecol Reprod Biol 2013;170 (1):188-192.
- 19. Murphy KM, McConnell TG, Hafez MJ, Vang R, Ronnett BM. Molecular genotyping of hydatidiform moles: analytic validation of a multiplex short tandem repeat assay. J Mol Diagn 2009;11(6):598-605.
- Repiská V, Shawkatová I, Böhmer D, Hatzibougias D, Šišovský V, Danihel L. Gestational choriocarcinoma analyzed by polymerase chain reaction amplification of polymorphic VNTR and human leukocyte antigen regions. Int J Gynaecol Obstet 2010; 110(2):152-154.
- LeGallo RD, Stelow EB, Ramirez NC, Atkins KA. Diagnosis of hydatidiform moles using p57 immunohistochemistry and HER2 fluorescent in situ hybridization. Am J Clin Pathol 2008;129(5):749-755.
- 22. Kipp BR, Ketterling RP, Oberg TN, Cousin MA, Plagge AM, Wiktor AE, et al. Comparison of Fluorescence In Situ Hybridization, p57 immunostaining, flow cytometry, digital image analysis for diagnosing molar and nonmolar products of conception. Am J Clin Pathol 2010;133(2):196-204.
- Soper JT, Mutch DG, Schink JC, American College of Obstetricians and Gynecologists. Diagnosis and treatment of gestational trophoblastic disease: ACOG Practice Bulletin No. 53. Gynecol Oncol 2004;93(3):575-585.
- Sebire NJ, Seckl MJ. Gestational trophoblastic disease: current management of hydatidiform mole. BMJ 2008;337:453-458.
- 25. Mukherjee M, Srivastava A, Kesari A, Mittal B. Analysis of VNTR loci, ApoB 3' HVR and D1S80

Avicenna Journal of Medical Biotechnology, Vol. 6, No. 4, October-December 2014

in North Indians. Indian J Biotechnol 2005;4:358-362.

- 26. Mahdieh N, Tafsiri E, Karimipour M, Akbari MT. Heterozygosity and allele frequencies of the two VNTRs (ApoB and D1S80) in Iranian population. Indian J Hum Genet 2005;11(1):31-34.
- Jeffreys AJ, Wilson V, Thein SL. Hypervariable minisatellite regions in human DNA. Nature 1985; 314(6006):67-73.
- 28. Mahianeh A, Bayat B, Banoei M, Sanati M, Jahangir Soltani N, Hashemzadeh Chaleshtari M. Allele distribution of apolipoprotein B 3'hyper variable region polymorphism between Lurs and Sistani-va-Baluchi two ethnic group from Iran. 4th national biotechnology congress, Islamic republic of Iran, Kerman. August 2005.
- 29. Van Meurs JBJ, Arp P, Fang Y, Slagboom PE, Meulenbelt I, Van Leeuwen JP, et al. Detection of sequence variability of the collagen type II^α1 3 var-

iable number of tandem repeat. Electrophoresis 2000;21(17):3571-3577.

- 30. Kloosterman A, Budowle B, Daselaar P. PCR-amplification and detection of the human D1S80 VNTR locus amplification conditions, population genetics and application in forensic analysis. Int J Legal Med 1993;105(5):257-264.
- 31. Boerwinkle E, Xiong W, Fourest E, Chan L. Rapid typing of tandemly repeated hypervariable loci by the polymerase chain reaction: Application to the apolipoprotein B 3' hypervariable region. Proc Natl Acad Sci USA 1989;86(1):212-216.
- 32. Cheung AN, Khoo US, Lai CY, Chan KY, Xue WC, Cheng DK, et al. Metastatic trophoblastic disease after an initial diagnosis of partial hydatidiform mole: genotyping and chromosome in situ hybridization analysis. Cancer 2004;100(7):1411-1417.

253