

Distribution and Diversity of *hmw1A* Among Invasive Nontypeable *Haemophilus influenzae* Isolates in Iran

Milad Shahini Shams abadi, Seyed Davar Siadat *, Farzam Vaziri, Mehdi Davari, Abolfazl Fateh, Shahin Pourazar, Farid Abdolrahimi, and Morteza Ghazanfari

Department of Mycobacteriology and Pulmonary Research, Pasteur Institute of Iran, Tehran, Iran

Abstract

Background: The pathogenesis of nontypeable *Haemophilus influenzae* (NTHi) begins with adhesion to the rhinopharyngeal mucosa. Almost 38-80% of NTHi clinical isolates produce proteins that belong to the High Molecular Weight (HMW) family of adhesins, which are believed to facilitate colonization.

Methods: In the present study, the prevalence of *hmwA*, which encodes the HMW adhesin, was determined for a collection of 32 NTHi isolates. Restriction Fragment Length Polymorphism (RFLP) was performed to advance our understanding of *hmwA* binding sequence diversity.

Results: The results demonstrated that *hmwA* was detected in 61% of NTHi isolates. According to RFLP, isolates were divided into three groups.

Conclusion: Based on these observations, it is hypothesized that some strains of nontypeable *Haemophilus influenzae* infect some specific areas more than other parts.

Avicenna J Med Biotech 2016; 8(2): 99-102

Keywords: Adhesins, *Haemophilus influenzae*, HMW1 protein

* Corresponding author:
Seyed Davar Siadat, Ph.D.,
Department of Mycobacteriology
and Pulmonary Research,
Pasteur Institute of Iran,
Tehran, Iran
Tel: +98 21 66953311
E-mail:
d.siadat@gmail.com
Received: 19 Aug 2015
Accepted: 15 Oct 2015

Introduction

Nontypeable *Haemophilus influenzae* (NTHi) is a small, fastidious gram-negative coccobacillus that colonizes the human pharynx, the only known natural reservoir. Although generally considered a commensal, NTHi is capable of inducing localized infections of the upper and lower respiratory tracts, such as acute otitis media, sinusitis, and bronchitis, as well as, more rarely, severe invasive infections such as pneumonia, bacteremia and meningitis¹⁻³.

The initial step in the pathogenesis of NTHi involves establishment of bacteria on the rhinopharyngeal respiratory mucosa followed by contiguous spreading within the respiratory tract and, occasionally, to sterile sites⁴.

High Molecular Weight (HMW) adhesins, which have been reported in 38-80% of NTHi isolates, are the major adhesins responsible for attachment to human epithelial cells⁵. The HMW adhesins are a family of paralogous proteins encoded by the *hmw* locus, which is present in two complete copies, *hmw1* and *hmw2*, located discontinuously on the NTHi chromosome⁶. Each *hmw* locus encodes three proteins, *hmwA*, *hmwB*, and *hmwC* and both the gene content and the chromosomal locations of the *hmw* loci are conserved across isolates^{7,8}.

The *hmwA* encodes the functional HMW adhesins while *hmwB* and *hmwC* encode proteins required for

maturation, glycosylation and secretion of mature HMW adhesions^{9,10}. In particular, the 124 amino acids between residues 114 and 237 in mature HMW1 have been found to be essential for full-level adhesive activity¹¹. These essential regions have been previously referred to as HMW1 core-binding domains and the encoding sequences as the *hmw1A* core-binding domain sequences^{12,13}. Amino acid diversity within the HMW binding domain likely serves the diversity of *hmwA* locus in various strains¹⁴.

The research presented in this study identified diversity of the *hmw1A* locus in different clinical strains isolated from nasopharynx, middle ear, pleura and conjunctiva.

Materials and Methods

Bacterial strains and growth

NTHi strains were obtained from four infection sites: nasopharyngeal or throat isolates from healthy children, strains cultured from middle ear aspirates from children with otitis media and pleural fluid and conjunctiva isolates obtained from children with invasive diseases. Strains were isolated between 2011 and 2013, at different locations in Iran. NTHi isolates were grown overnight on 7% chocolate agar plates supplemented with Vitox (Oxoid Ltd., Basingstoke, Hampshire) at 37°C in 5% CO₂ and were identified by their

Table 1. Polymerase Chain Reaction (PCR) primers used in this study

Primer name	PCR primer sequence (5' to 3')	References
hmwA_leader_F	CCATGGTTGATGTTTCATAAAAAT	This study
hmwA_leader_R	TCGAGTACATTAAGTGAATTT	This study

requirements for X and V factors and by their lack of reaction with antisera against capsular (Phadebact; KaroBio Diagnostics AB). For DNA extraction, bacterial strains were grown in Haemophilus Test Medium (HTM) broth, consisting of Muller-Hinton broth (Oxoid Ltd.) supplemented with 0.5% yeast extract and HTM supplement (Oxoid Ltd.), and incubated under the same conditions.

DNA amplification

Oligonucleotide primers were designed (Table 1) on the basis of the sequence of the *hmw1A* gene. Regions with low variability were chosen. The sequence of each primer was checked for homology to other sequences that may also be amplified by them, in the GenBank and EMBL databases. The PCR product was a 1.045-bp nucleotide fragment containing regions with low variability of the *hmw1A* gene.

Restriction fragment length analysis

The amplified DNA fragments were digested with one restriction endonuclease (Taq1) (5'TCGA-3'AGCT), which was selected on the basis of the nucleotide sequence of the *hmw1A* gene of NTHi.

Nucleotide sequencing

A PCR product of each restriction fragment pattern (rfp) was randomly chosen and was sequenced.

Sequence analysis

Sequence analysis was performed with the BioEdit and CLC Main Workbench 5 software.

Results

The strains containing *hmw1A* gene were separated with PCR of *hmw1a* gene. Only 20 of the 32 isolates contained the *hmw1a* gene. Polymorphism in the *hmw1A* gene was found in *Haemophilus influenzae* strains isolated from the nasopharynx, middle ear, pleural fluid and conjunctiva in Iran. Figure 1 shows the RFLP patterns of *hmw1A* gene in nontypeable *Haemophilus influenzae* strains isolated from the nasopharynx, middle ear, pleural fluid and conjunctiva by Taq1 digested PCR products.

This restriction enzyme (Taq1) gave three different patterns with all isolated strains, isolated strains from pleural fluid and conjunctiva contained restriction fragment pattern (rfp1) and isolated strains from nasopharynx contained rfp1 and rfp2 (Figure 2).

A sample was selected randomly from each pattern for sequence. The PCR product of the numbers (Ppn) 3, 9 and 13 were representative of the restriction fragment pattern of 1, 2 and 3, respectively. CLC Main Workbench 5 software revealed that sample numbers 1, 2 and 3 were similar to strains of nontypeable *Haemo-*




Figure 1. Restriction fragment pattern (rfp) of *hmw1A* gene fragment amplified by PCR. RFLP patterns were obtained after digestion of PCR products with Taq1. Number 0, PCR products of *hmw1A* gene without restriction enzyme; numbers 1-3, digestion products of *hmw1A* gene fragments amplified (three different patterns); the last lane, molecular weight marker.




Figure 2. This chart describes the number and type of pattern of the strain isolated from each section.



Figure 3. Strains with a coefficient of similarity value $\geq 80\%$ were considered to belong to the same cluster. On the right, the strain code number and the different clusters (clusters 1 through 3) are reported.

philus influenzae with GenBank accession numbers (Gan), AY497551, HIU08876 and AY497554, respectively (Figure 3).

Discussion

Nontypeable *Haemophilus influenzae* is a major cause of localized respiratory tract diseases, including otitis media, sinusitis, conjunctivitis, bronchitis and pneumonia and strains of bacteria are various ^{5,15,16}. For prevention of infection or immunomodulation of chronically infected individuals for appropriate humoral and cellular immune mechanisms, it is necessary to recognize the available strains in different regions.

In this study, an attempt was made to determine the prevalence of *hmw1A* in a collection of 32 NTHi isolates collected between 2011 and 2013 from the throat isolates of healthy children, the middle ears of children

with otitis media and pleural fluid and conjunctiva isolates obtained from children with invasive diseases. The *hmwA* prevalence (63%) in this isolate collection was consistent with previous studies, which report prevalences ranging from 38 to 80%^{5,8,17}. In previous studies, to determine the type of strains in a region, some techniques such as MLST were used, or genome of all strains had been sequenced¹².

In this study, it was revealed that *hmwIA* genes of nontypeable *Haemophilus influenzae* in Iran are similar to strains of nontypeable *Haemophilus influenzae* with GenBank accession numbers, AY497551, HIU08876 and AY497554. Several studies have demonstrated that phase variable HMW adhesin expression has important implications for NTHi pathogenesis¹⁸⁻²⁰. In this study, it was found that the restriction fragment pattern 3 (rfp 3) that is similar to GenBank accession number HIU-08876 is not indeed in throat isolates but is present in some otitis isolates and rfp 1 that is similar to GenBank accession number AY497551 is present in pleural fluid, throat, conjunctiva and otitis isolates. Efficient adherence of nontypeable *Haemophilus influenzae* to a variety of mammalian cells is dependent on bacterial expression of HMW proteins²¹ and based on these observations, it is hypothesized that some strains of nontypeable *Haemophilus influenzae* infect some specific areas more than other parts or in other words, the strains on those areas are more connected and they stimulate colonization. However, more studies are needed to prove this hypothesis. To obtain and submit a comprehensive report, it is required to collect more samples and apply newer molecular methods.

Finally, this work has important public health implications since the HMW adhesins have been proposed as potential components of the NTHi targeted vaccine. Phylogenetic analysis can be used to inform the selection of HMW adhesins for inclusion in a vaccine, e.g., selecting representative alleles from each phylogenetic cluster. In previous studies, the extensive amino acid diversity and the lack of disease-specific *hmwA* sequence clusters underscore the challenges of targeting HMW adhesins as vaccine components² but in this study, specific sequences of *hmwA* in the specific sites were observed.

Conclusion

The prevalence of *hmwIA* among invasive nontypeable *Haemophilus influenzae* isolates in Iran is almost similar to previous studies reported in other countries, and based on our observations, strains with specific sequence of *hmwIA* in some parts of the body were found, whereas in other parts of the body other sequences of the *hmwIA* were observed.

References

1. Barenkamp SJ, St Geme JW 3rd. Identification of a second family of high-molecular-weight adhesion pro-

- teins expressed by non-typable *Haemophilus influenzae*. *Mol Microbiol* 1996;19(6):1215-1223.
2. Davis GS, Patel M, Hammond J, Zhang L, Dawid S, Marrs CF, et al. Prevalence, distribution, and sequence diversity of *hmwA* among commensal and otitis media non-typeable *Haemophilus influenzae*. *Infect Genet Evol* 2014;28:223-232.
3. Davis GS. Nontypeable *Haemophilus influenzae* high molecular weight adhesins: molecular epidemiology, evolution, and within-host population dynamics [dissertation]. [USA]: University of Michigan; 2013. 203 p.
4. Rao VK, Krasan GP, Hendrixson DR, Dawid S, St Geme JW 3rd. Molecular determinants of the pathogenesis of disease due to non-typable *Haemophilus influenzae*. *FEMS Microbiol Rev* 1999;23(2):99-129.
5. Erwin AL, Sandstedt SA, Bonthuis PJ, Geelhood JL, Nelson KL, Unrath WC, et al. Analysis of genetic relatedness of *Haemophilus influenzae* isolates by multilocus sequence typing. *J Bacteriol* 2008;190(4):1473-1483.
6. Buscher AZ, Burmeister K, Barenkamp SJ, St Geme JW 3rd. Evolutionary and functional relationships among the nontypeable *Haemophilus influenzae* HMW family of adhesins. *J Bacteriol* 2004;186(13):4209-4217.
7. Barenkamp SJ, St Geme JW 3rd. Genes encoding high-molecular-weight adhesion proteins of nontypeable *Haemophilus influenzae* are part of gene clusters. *Infect Immun* 1994;62(8):3320-3328.
8. Ecevit IZ, McCrea KW, Marrs CF, Gilsdorf JR. Identification of new *hmwA* alleles from nontypeable *Haemophilus influenzae*. *Infect Immun* 2005;73(2):1221-1225.
9. Grass S, St Geme JW 3rd. Maturation and secretion of the non-typable *Haemophilus influenzae* HMW1 adhesin: roles of the N-terminal and C-terminal domains. *Mol Microbiol* 2000;36(1):55-67.
10. Xie J, Juliao PC, Gilsdorf JR, Ghosh D, Patel M, Marrs CF. Identification of new genetic regions more prevalent in nontypeable *Haemophilus influenzae* otitis media strains than in throat strains. *J Clin Microbiol* 2006;44(12):4316-4325.
11. Dawid S, Grass S, St Geme JW 3rd. Mapping of binding domains of nontypeable *Haemophilus influenzae* HMW1 and HMW2 adhesins. *Infect Immun* 2001;69(1):307-314.
12. Giufrè M, Muscillo M, Spigaglia P, Cardines R, Mastantonio P, Cerquetti M. Conservation and diversity of HMW1 and HMW2 adhesin binding domains among invasive nontypeable *Haemophilus influenzae* isolates. *Infect Immun* 2006;74(2):1161-1170.
13. St Geme JW 3rd. The pathogenesis of nontypable *Haemophilus influenzae* otitis media. *Vaccine* 2000;19 Suppl 1:S41-50.
14. van Schilfgaarde M, van Alphen L, Eijk P, Everts V, Dankert J. Paracytosis of *Haemophilus influenzae* through cell layers of NCI-H292 lung epithelial cells. *Infect Immun* 1995;63(12):4729-4737.
15. Hardy GG, Tudor SM, St Geme JW 3rd. The pathogenesis of disease due to nontypeable *Haemophilus influenzae*. *Methods Mol Med* 2003;71:1-28.

16. Erwin AL, Smith AL. Nontypeable *Haemophilus influenzae*: understanding virulence and commensal behavior. *Trends Microbiol* 2007;15(8):355-362.
17. Barenkamp SJ, Leininger E. Cloning, expression, and DNA sequence analysis of genes encoding nontypeable *Haemophilus influenzae* high-molecular-weight surface-exposed proteins related to filamentous hemagglutinin of *Bordetella pertussis*. *Infect Immun* 1992;60(4):1302-1313.
18. Cholon DM, Cutter D, Richardson SK, Sethi S, Murphy TF, Look DC, et al. Serial isolates of persistent *Haemophilus influenzae* in patients with chronic obstructive pulmonary disease express diminishing quantities of the HMW1 and HMW2 adhesins. *Infect Immun* 2008;76(10):4463-4468.
19. Davis GS, Marino S, Marrs CF, Gilsdorf JR, Dawid S, Kirschner DE. Phase variation and host immunity against high molecular weight (HMW) adhesins shape population dynamics of nontypeable *Haemophilus influenzae* within human hosts. *J Theor Biol* 2014;355:208-218.
20. Dawid S, Barenkamp SJ, St Geme JW 3rd. Variation in expression of the *Haemophilus influenzae* HMW adhesins: a prokaryotic system reminiscent of eukaryotes. *Proc Natl Acad Sci USA* 1999;96(3):1077-1082.
21. Noel GJ, Love DC, Mosser DM. High-molecular-weight proteins of nontypeable *Haemophilus influenzae* mediate bacterial adhesion to cellular proteoglycans. *Infect Immun* 1994;62(9):4028-4033.

Archive of SID