

A Single Tube Overlap Extension PCR Method for Splicing of Multiple DNA Fragments

Farzaneh Zarghampoor¹, Abbas Behzad-Behbahani^{2*}, Negar Azarpira^{3*}, Saeed Reza Khatami¹, Maryam Fanian³, Mahdokht Hossein Aghdaie³, and Gholamreza Rafiei Dehbidi²

1. Department of Biology, Faculty of Science, Shahid Chamran University of Ahvaz, Ahvaz, Iran

2. Diagnostic Laboratory Sciences and Technology Research Centre, Faculty of Paramedical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

3. Transplant Research Centre, Shiraz University of Medical Science, Shiraz, Iran

Abstract

Background: Despite the ease of conventional splicing by overlap-extension (SOEing) PCR technique in theory, when splicing more than two fragments, and especially if one of the complementary sequences is A-T rich, the attachment of the fragments would be challenging. A new rapid and highly efficient SOEing PCR assay was developed for simultaneous splicing of multiple DNA fragments and induction of site-directed mutagenesis in a single tube.

Methods: The method was adapted for splicing human beta-globin UTRs to OCT4, SOX2, KLF4, C-MYC, LIN28A, and destabilized GFP for the construction of chimeric DNA fragments for *in vitro* transcription. In addition, the native Kozak sequence of beta-globin (K1) was replaced by the strongest Kozak sequence (K2) using site-directed mutagenesis to enhance the expression of target genes.

Results: ChimericGFPd2/K1, GFPd2/K2, OCT4, and KLF4 were created by the optimized conventional SOEing PCR. The single tube method was able to create the chimeric SOX2, C-MYC, and LIN28A in high quality and quantity in comparison with the conventional SOEing PCR. Moreover, using single tube SOEing PCR, the reaction time and materials that are required in the conventional SOEing PCR were significantly reduced. Fluorescent microscopy and flow cytometry examinations indicated highly efficient translation of K2 sequence in comparison with the K1sequence.

Conclusion: Single tube SOEing PCR is a valuable method to construct more multiple fragments with high yield. The method can successfully be applied for construction of various kinds of complex chimeric genes.

Avicenna J Med Biotech 2020; 12(1): 37-43

Keywords: Beta-globins, Mutagenesis, Polymerase chain reaction, Site-directed, Untranslated regions

Introduction

The Splicing by overlap-extension/Splicing by overlap-extension PCR (SOEing PCR) is a type of PCR which is used to insert specific mutations at specific points in a sequence^{1,2} or splice smaller DNA fragments to construct chimeric gene fragment with no dependence on the restriction site or ligase³. Any overlap sequences or mismatches can be incorporated into the 5'end of primers, so that a DNA fragment with incorporate new sequences which did not exist in the first template is created. This idea was first introduced under the title of mispriming^{4,5}. Using simple mispriming can create site-directing mutagenesis only at the end of PCR products⁶, whereas overlap-extension can generate mutations in the center of PCR-products².

To create chimeric DNA fragment by the conventional SOEing PCR, two separate PCR-generated products should have been a short overlap of complement-

ary sequence (typically 30-60 bp). The overlap can be formed in the PCR products by addition of bases at the 5' ends of primers. Subsequently, the fragments are combined in equal amounts of molecules of all fragments to create a longer fragment sequence. After splicing two fragments at the complementary sequence, the 3' end of each fragment plays the role of primer and continues the extension. The resulting product is further amplified by PCR⁶⁻⁸.

Despite the ease of the SOEing PCR technique in theory and its advantages compared with the other techniques such as restriction method, its complexity limits its application^{7,8}. When splicing more than two fragments, especially if the complementary sequence is AT-rich⁹, splicing of fragments would be challenging. In addition, in the conventional SOEing PCR, the smear or multiple bands are often seen on the agarose

*** Corresponding authors:**

Abbas Behzad-Behbahani, Ph.D., Diagnostic Laboratory Sciences and Technology Research Centre, Faculty of Paramedical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

Negar Azarpira, Ph.D., Transplant Research Centre, Shiraz University of Medical Science, Shiraz, Iran

Tel: +98 71 32270301

Fax: +98 71 32270301

E-mail: behzadba@sums.ac.ir,

behzadba@gmail.com,

negarazarpira@gmail.com

Received: 12 Jan 2019

Accepted: 4 May 2019

gel electrophoresis and occasionally the main band is very weak. On the other hand, constructing a chimeric DNA fragment from multiple small fragments requires several PCR reactions, thus the conventional SOEing PCR is tedious, time-consuming, and not cost-effective. Furthermore, random error may be increased by the polymerase during several PCR reactions. Although some researchers have successfully assembled up to four DNA fragments simultaneously^{10,11}, multiple DNA splicing based on overlap extension PCR remains a challenge which needs to be improved.

In the present study, a single tube overlap extension PCR method was developed for the simultaneous fusion of 5' and 3' untranslated regions of human beta globin into classical reprogramming genes including SOX2, C-MYC, and LIN28A. Furthermore, to obtain the maximum production of chimeric DNA constructs including destabilized GFP, OCT4, and KLF4, the conventional SOEing PCR method was modified by improving parameters such as annealing temperature of the complementary sequences, reaction conditions, and elongation time. The chimeric DNA templates were then used for *in vitro* transcription and production of stable synthetic mRNA after transfection in the eukaryotic cells. In addition, the SOEing PCR technique was adapted to replace the native Kozak sequence (K1) of beta-globin by the strongest Kozak sequence (K2) to enhance the translation efficiency of the mRNA.

Materials and Methods

In the present study, for splicing the UTRs of beta-globin gene to Reprogramming Factors (RFs) and GFPd2, the conventional and developed overlap extension PCR (single tube) were used.

Primer designing for splicing of DNA fragments

The GeneRunner software (Version 6.5.51×64 Beta) was used to design the primers for the PCR amplification of different DNA fragments. The sequences of the nineteen set of primers used for this recombination are listed in table 1. The internal primers (the primer with a short overlap of complementary sequence) were designed between 20 and 33 bp for splicing the fragments together and generating site-directed mutation in the K1 sequence in order to create the K2 sequence. The primer specificity was confirmed by Primer-BLAST (www.ncbi.nlm.nih.gov/tools/primer-blast). The internal primers containing the K1 and K2 sequences were just designed for chimeric GFPd2 construct and the internal primers of RF contained the K2 sequence.

Plasmids

PUC57 vector containing 5' and 3' UTR of human beta-globin, T7 promoter for *in vitro* transcription and the native Kozak sequence (K1) of human beta-globin were synthesized (Biomatik Corporation, Canada). The plasmids of pMXs-hOCT4¹², pMXs-hSOX2¹², pMXs-hKLF4¹², pMXs-hcMYC¹², pMXs-hLIN28A¹³, and pCAG-GFPd2¹⁴ were purchased from Addgene, a US

non-profit organization. The pcDNA 3.1+ plasmid (Invitrogen) contains a CMV promoter serving as plasmid backbone for cloning of the chimeric DNA fragments.

The simple PCR reaction conditions

The UTRs and gene fragments were separately amplified by the simple PCR method, using Q5® Hot Start High-Fidelity 2X Master Mix (New England Biolabs, U.K). The 5'UTR containing the K2 sequence was created by site-directed mutagenesis SOEing PCR. The PCR products were determined on agarose gel electrophoresis and then recovered from the gel by AccuPrep® Gel Purification Kit (Bioneer, South Korea) according to the manufacturer's protocol. The simple PCR reactions were performed in a final volume of 20 µL of mixture containing 10 µL of master mix, 1 µL of the plasmid DNA, and 0.5 µM of each primer. The PCR reactions were performed in the following cycling conditions: 98°C for 2 min, 30 cycles at 98°C for 10 s, X°C for the 30 s, 72°C for 1 min, and a final elongation step at 72°C for 4 min. The annealing temperatures (X°C) for the PCR reactions were estimated using NEB Tm calculator (<https://tmcalculator.neb.com/#/main>).

Splicing by overlap extension (SOEing) polymerase chain reaction

In order to splice DNA fragments together, the annealing temperature of complementary sequence I and II (I: 5'UTR-gene; II: gene-3'UTR) was initially estimated. The chimeric DNA fragments including GFPd2, OCT4 and KLF4 were created by the conventional SOEing PCR and chimeric reprogramming factors including SOX2, C-MYC, and LIN28A created by the single tube SOEing PCR.

The conventional SOEing PCR reaction conditions

To construct chimeric DNA fragments using the conventional SOEing PCR, the 3'UTR was initially ligated to RFs or GFPd2 (in the SOE 1 reaction) and then 5'UTR was ligated (in the SOE 2 reaction). To achieve this, the fragments were combined in equal amounts of molecules of two fragments without primers and the PCR mixture was subjected to PCR with the following cycling conditions, as illustrated in table 2. The annealing temperature for the ligation of the two fragments was determined with the complementary sequence of two fragments and using NEB Tm calculator. To carry out the steps 1 and 2 of the conventional SOE 1 reaction, the final volume of reaction mixture was 18.6 µL containing 2.5 µL 10X PFU buffers, 2.5-3 U PFU DNA polymerase, 0.1 mM dNTPs, and 2.5 µL of the DNA fragment mixture (372 ng/2 µL of GFPd2 + 67.64 ng/1.78 µL of 3'UTR). In step 3, 0.5 µM of primers was added into the mixture and the steps 4 and 5 of PCR were performed. To confirm the ligation of the two DNA fragments, PCR products were run on agarose gel electrophoresis and were then purified from agarose gel. Using the SOE 2 reaction, 5'UTR fragment was ligated to the gene-3'UTR fragment with the same procedures. The 5'UTR/K2-GFPd2-3'UTR was

Table 1. Primers designed for amplifying fragments, the OE- PCR, and generation of the K2 90 sequence

DNA fragment name	Primer name	Primer sequence	Product length (bp)
5'UTR	5'UF 5'UR1	5' tcaaggatcc GATCAATAATACGACTCACTATAG3' 5' ctcacc ATGGTGTCTGTTTGAGGTTG 3'	176
	5'UF 5'UR2	5' tcaaggatcc GATCAATAATACGACTCACTATAG3' 5' tcccccat GGTGGCGGTTTGAG 3'	176
	5'UF 5'UR3	5' tcaaggatcc GATCAATAATACGACTCACTATAG3' 5' tggtgtacat GGTGGCGGTTTGAGG 3'	177
	5'UF 5'UR4	5' tcaaggatcc GATCAATAATACGACTCACTATAG3' 5' gacagccat GGTGGCGGTTTGAG 3'	176
	5'UF 5'UR5	5' tcaaggatcc GATCAATAATACGACTCACTATAG3' 5' aggggcat GGTGGCGGTTTGAGG 3'	175
	5'UF 5'UR6	5' tcaaggatcc GATCAATAATACGACTCACTATAG3' 5' agcccat GGTGGCGGTTTGAG 3'	174
3'UTR	3'UF1 3'UR	5' caatgtgtag GCTCGTTTCTTGCTGTCC 3' 5' cacagaattc GCTTCTTTTGCAATG3'	166
	3'UF2 3'UR	5' cattcaaaactga GCTCGTTTCTTGCTGTCC3' 5' cacagaattc GCTTCTTTTGCAATG3'	166
	3'UF3 3'UR	5' catatgta GCTCGTTTCTTGCTGTCC3' 5' cacagaattc GCTTCTTTTGCAATG3'	163
	3'UF4 3'UR	5' gacatttttaa GCTCGTTTCTTGCTGTCC3' 5' cacagaattc GCTTCTTTTGCAATG3'	165
	3'UF5 3'UR	5' ttgtgcgtga GCTCGTTTCTTGCTGTCC 3' 5' cacagaattc GCTTCTTTTGCAATG3'	164
	3'UF6 3'UR	5' acagaattga GCTCGTTTCTTGCTGTCC 3' 5' cacagaattc GCTTCTTTTGCAATG3'	164
GFPd2	GK1F GK1R	5' gacacc ATGGTGAGCAAGGGCGAG 3' 5' gaaagcgagc CTACACATTGATCCTAGCAGAAG 3'	853
Modified Kozak	5'UK2F 5'UK2R	5' tcaaggatcc GATCAATAATACGACTCACTATAG 3' 5' cttgetccatc GGTGGCGGGTTGAGGTTG 3'	176
GFPd2K2	GK2F GK2R	5' aacagacacc ATGGTGAGCAAGGGCGAG 3' 5' gaaagcgagc CTACACATTGATCCTAGCAGAAG	858
OCT4	OF OR	5' gccacc ATGGCGGGACACCTG 3' 5' gaaagcgagc TCAGTTTGAATGCATGG 3'	1099
SOX2	SF SR	5' aaccgccacc ATGTACAACATGATG 3' 5' gaaagcgagc TCACATGTGTGAGAGG 3'	974
KLF4	KF KR	5' cgccacc ATGGCTGTCAGTG 3' 5' gaaagcgagc TTAAAAATGTCTTTCATGTG 3'	1431
C-MYC	MF MR	5' cgccacc ATGCCCTCAACGTTAG 3' 5' gaaagcgagc TCACGCACAAGAGTTC 3'	1337
LIN28A	LF LR	5' cgccacc ATGGGCTCCGTGTC 3' 5' aaagcgagc TCAATTCTGTGCCTCC 3'	646

Key: Overlap of complementary sequence are shown in bold lowercase letters.

also constructed using the conventional SOEing PCR method.

The single tube SOEing PCR reaction conditions

In a single tube SOEing PCR, the fragments were combined in equal amounts of molecules of three fragments without primers and the mixture was subjected to PCR with the following cycling conditions illustrated in table 3. The steps 1 and 2 of the single tube SOEing PCR reactions were carried out in a final volume of 28 µL of the mixture containing 5 µL 10X PFU buffers, 6 U PFU DNA polymerase, 0.2 mM dNTPs and 4 µL of the DNA fragment mixture. In step 3, 0.3 µM of external primers (5'UF and 3'UR) were added into the mixture and the steps 4 and 5 of PCR were performed.

Table 2. The OE 1 reaction for construction GFPd2-3'UTR fragment

Step	Cycle	Denaturation	Annealing	Extension
1	1	95°C for 3 min	--	--
2	12	95°C for 30 s	55°C for 45 s	72°C for 2 min
3	1	50°C for 6 min	--	--
4	30	95°C for 30 s	55°C for 45 s	72°C for 2 min
5	1	--	--	72°C 5 min

Cloning of the chimeric fragments

The full-length chimeric DNA constructs were excised and purified from the agarose gel and then were double digested using EcoRI (New England Biolabs, U.K) and BamHI (New England Biolabs, U.K) re-

Table 3. The single tube OE- PCR reaction for construction of 5'UTR+K2-SOX2/C-MYC/LIN28A-3'UTR

Step	Cycle	Denaturation	Annealing	Extension
1	1	95°C for 4 min	--	--
2	13	95°C for 30 s	X°C for 55 s	72°C for 3 min
3	1	50°C for 6 min	--	--
4	32	95°C for 30 s	55°C for 50 s	72°C for 3 min
5	1	--	--	72°C 5 min

striction enzymes. The constructs were then inserted into pcDNATM3.1 (+) mammalian expression vector (Invitrogen) using calcium chloride method and the recombinant plasmids were transformed into the *Escherichia coli* (*E. coli*) strain DH5 α . To screen the plasmids containing the desired inserted DNA, colony PCR was performed. For further confirmation, DNA sequencing was performed on all plasmids containing chimeric genes.

Cell culture and plasmid DNA transfection

The HEK293T cell line was cultured in DMEM supplemented with 10% FBS (Gibco, USA) and incubated at 37°C in 5% CO₂. For transfection, the cells were in the 3rd passage. The cells were seeded (5x10⁴ HEK293T cells/well) in 24-well plates a day prior to the experiment. The chimeric GFPd2/K1 or K2 plasmid transfections were performed with Lipofectamine 3000 (Invitrogen). Hence, the culture media was changed to Opti-MEM I Reduced Serum Media (Thermo Fisher Scientific) 2 hr before transfection. The transfection was performed according to the manufacturer's protocol and then the mixture was added to the culture media and incubated at 37°C in 5% CO₂. Four hours after transfection, the culture media was changed to DMEM supplemented with 10% FBS. A day post-transfection, the cells were monitored under a fluorescence microscope.

Flow cytometry assay

The HEK293T cell line was trypsinized 24 hr post-transfection and washed twice with PBS. Then, the cells were suspended in PBS and analyzed by flow cytometry method (BD FACS Calibur).

Results

Construction of the chimeric DNA fragments

In order to splice DNA fragments, the GC content and temperature of the complementary sequences were initially evaluated. The annealing temperature of complementary sequence I and II of SOX2, C-MYC, and LIN28A was the same while the annealing temperature of complementary sequence I and II of GFPd2, OCT4 and KLF4 was different. Thus, the chimeric DNA fragments including GFPd2, OCT4 and KLF4 were created by the conventional SOEing PCR (Figure 1) and chimeric reprogramming factors including SOX2, C-MYC, and LIN28A were created by the single tube SOEing PCR (Figure 2). In addition, the K1 sequence (ACAG-

ACACCATG) was successfully replaced by K2 sequence (GCCGCCACCATG) using SOEing PCR to enhance translation efficiency. Surprisingly, the concentration of purified chimeric SOX2, C-MYC, and LIN28A was higher than GFPd2, OCT4, and KLF4 (for example, SOX2: 358 ng/ μ L vs. GFPd2: 52 ng/ μ L). Figure 3 shows the PCR products of simple PCRs and the SOEing PCRs of reprogramming factors and GFPd2 on agarose gel electrophoresis.

Plasmid DNA transfection and flow cytometry assay

To evaluate the performance of Kozak sequences in the efficiency of translation, both chimeric GFPd2 con-




Figure 1. Construction of chimeric GFPd2 fragments using the conventional OE- PCR. In the first stage, the 5'UTR-K1 (A), 5'UTR-K2 (A'), 3'UTR (B), GFPd2 with K1 sequence (C), and GFPd2 with K2 sequence (C') fragments were amplified using the simple PCR. The fragments had a complementary overlapping end to ensure splicing fragments together. The complementary sequence II was AT-rich; thus, the OE 1 reaction was initially performed for ligation of 3'UTR to GFPd2. In the OE 1 reaction, fragment (B) spliced to fragments (C and C') and created the new chimeric fragments (CB and C'B). The 3'end of each fragment plays the role of primer and continues extension. The resulting products were amplified further by PCR. Following, the fragments (A and A') spliced to fragments (CB and C'B) using the OE 2 reaction and created the new chimeric fragments (ACB and A'C'B) as well. Finally, the fragments (ACB and A'C'B) were amplified with outside primers. The final chimeric constructs double digested using EcoRI and BamHI restriction enzymes; then, the constructs were inserted into pcDNATM3.1 (+) Vector by Calcium chloride method.






Figure 2. The principle of construction of the chimeric SOX2 fragment using the single tube OE- PCR. The complementary sequence I and II of SOX2 had the same annealing temperature; therefore, the chimeric DNA fragment was constructed using the single tube OE-PCR method. The 3'end of each fragment plays the role of primer and continues extension. The resulting products were amplified with the outside primer.




Figure 3. Construction of chimeric reprogramming factors and GFPd2. OCT4: 100 bp-5 Kb DNA ladder (L), 5'UTR (176 bp), 3'UTR (166 bp), OCT4 fragment (1099 bp), OE 1 (1260 bp), OE 2 (1404 bp). SOX2: 100 bp-5 Kb DNA ladder (L), 5'UTR (177 bp), 3'UTR (163 bp), SOX2 fragment (974 bp), single tube OE- PCR (1275 bp). KLF4: 100 bp-5 Kb DNA ladder (L), 5'UTR (176 bp), 3'UTR (165 bp), KLF4 fragment (1431 bp), OE 1 (1590 bp), OE 2 (1734 bp). C-MYC: 100 bp-5 Kb DNA ladder (L), 5'UTR (175 bp), 3'UTR (164 bp), C-MYC fragment (1337 bp), single tube OE- PCR (1641 bp). LIN28A: 100 bp-5 Kb DNA ladder (L), 5'UTR (174 bp), 3'UTR (164 bp), LIN28A fragment (646 bp), single tube OE- PCR (951 bp). GFPd2: 100 bp-5 Kb DNA ladder (L), 5'UTR-k1 (176 bp), 5'UTR-k2 (176 bp), 3'UTR (166 bp), GFPd2 fragment (853 bp), OE 1 (1011 bp), OE 2 (1147 bp).

structs were transfected into HEK293T cells using Lipofectamine 3000. The results obtained from fluorescent microscope 24 hr post-transfection showed that GFP production from the K2 plasmid was significantly more than the K1 plasmid. On the other hand, the transfected cells with the K2 plasmid initiated GFP expression earlier than the K1 plasmid. The GFP expression intensity of both plasmids was evaluated by




Figure 4. Transfection using Lipofectamine 3000. (A): The transfection of chimeric GFPd2 with the K2 and K1 sequence in HEK293T cell line after 24 hr illustrated by fluorescent microscope. (B): The mean fluorescent intensity (MFI) of GFP expression from chimeric GFPd2-K2 or K1 plasmid 24 hr after transfection.

flow cytometry assay as well. Overall, the mean fluorescence intensity of the K2 plasmid was higher than the K1 plasmid. Hence, other chimeric genes were constructed with the K2 sequence (Figure 4).

Discussion

There are three procedures to construct a chimeric DNA fragment, including gene synthesis, the restriction method and the SOEing PCR strategy. Compared with the SOEing PCR approach, the restriction method needs the integration of mutations to construct unique restriction sites. Thus, the restriction method cannot be made at any position within the nucleotide sequence. On the other hand, construction of the unique restriction site requires introduction of an unwanted sequence into the junction sites. Nevertheless, the SOEing PCR successfully achieved gene ligation, using the polymerase chain reaction at any chosen location devoid of unwanted sequence¹⁵. Another method to construct a chimeric DNA fragment is gene synthesis but for construction, the large chimeric DNA fragment is not cost-effective.

The conventional SOEing PCR method included several PCR reactions and requires laborious steps to purify the chimeric intermediate fragments to construct the final recombinant fragments. Thus, this method is tedious, time-consuming, expensive, and may increase mutations in the sequence of DNA fragments. Despite success to assemble more than three DNA fragments simultaneously^{16,17}, multiple DNA splicing in a single tube remains a challenge for researchers. A single tube SOEing PCR method was successfully developed in producing the desired chimeric fragments and induction of site-directed mutation. This OE-PCR method is distinct from previous gene assembly approaches^{18,19} in that the parameters such as annealing temperature of the complementary sequences, reaction conditions, and elongation time were improved.

The induced Pluripotent Stem Cells (iPSCs) hold promise in the field of regenerative medicine, tissue bioengineering, disease modeling, autologous cell therapy, and basic research. There are several procedures can be successfully generated iPSCs¹⁹. Of those, IVT mRNA-reprogramming has several advantages compared with the other procedures²⁰⁻²². However, IVT mRNA has several limitations to be overcome. The main limitation of IVT mRNA is short half-life mRNA and mRNA-mediated translation. One approach for increasing IVT mRNA stability and protein translation is used 5' and 3' UTR of genes that have long half-life to construct the chimeric reprogramming factors. Therefore, in the present study spliced the 5' and 3'UTRs of human beta-globin to GFPd2 and RFs to construct *in vitro* transcription DNA template. Additionally, to further enhance protein production, the Kozak sequence (K1) of the 5'UTR beta-globin gene replaced by the strongest Kozak sequence (K2) by the site-directed mutagenesis OE-PCR as well.

Concentration, ratio, and purity of initial fragments are the key factors influencing the efficacy of the SOEing PCR reaction. The fragment concentration ratio is the inverse of the fragment size ratio. Consequently, in the present study, the conventional SOEing PCR operational conditions such as regulation of PCR conditions (including T_m, time of the extension, ratio, and concentration of the initial DNA template) were optimized and a single tube SOEing PCR was introduced.

In the conventional SOEing PCR (For example, to construct chimeric 5'UTR-GFPd2-3'UTR), three fragments were amplified which had a short overlap of the complementary sequence. Subsequently, SOE 1 and 2 reaction was performed for generation of GFPd2-3'UTR and 5'UTR-GFPd2-3'UTR, respectively. It was found that for ligation of different fragments, T_m and GC% of the complementary sequence should first be calculated. The annealing temperature of the complementary sequence of the two fragments was estimated, using NEB T_m calculator or according to the formula $T_d = 4(C+G) + 2(A+T)$ in Celsius degrees. In the SOE reaction, the fragments that had low T_m or GC% complementary sequence were preferentially spliced. It is critical to use PFU DNA polymerase for the SOEing PCR because Taq DNA polymerase may create mutations in the sequence, and especially in the case of *in vitro* transcription, the DNA template should be error-free. Thus, Q5 Hot Start High-Fidelity 2X Master Mix was used for amplification. However, it was found that it would be better to use PFU DNA polymerase in the ligation steps and increase the extension time up to 2-3 min. Interestingly, it was found that if the annealing temperature of complementary sequence I and II is the same, SOEing PCR can be used in the single tube reaction. Therefore, ligation of UTRs to SOX2, C-MYC, and Lin28A was performed in the single tube SOEing PCR reaction.

In cases of GFPd2, OCT4 and KLF4, the annealing temperature of complementary sequence I and II was different. Hence, these chimeric fragments were constructed using the optimized conventional SOEing PCR. In each case, the chimeric DNA fragments were successfully generated. However, the concentration of purified chimeric DNA fragments from the single tube SOEing PCR was significantly higher than the conventional SOEing PCR procedures. The method provided a maximum efficiency yield of chimeric DNA fragments in minimum time, reduced the steps in the procedure, and was cost effective compared with the conventional method. Thus, in designing the internal primers (primer with overlap sequence), the annealing temperature of complementary sequences should be the same or similar so that the SOEing PCR can be performed in single tube procedures. However, in order to construct complex chimeric DNA fragments with different annealing temperatures of complementary sequence, it is suggested to perform touch-up SOEing PCR method.

Conclusion

To sum up, the current study described a very reliable and efficient SOEing PCR procedure that allows a quick construction of chimeric DNA fragments. The procedure allows to combine more than two DNA fragments simultaneously in a single reaction tube. The method prevents the risk of undesirable mutation which usually occurs when DNA polymerase is used. The efficiency and simplicity of this procedure makes it a valuable approach for generating chimeric fragments in genetic engineering studies.

Acknowledgement

The present article was extracted from a Ph.D. thesis and supported by a National Institute for Medical Research Development (NIMAD) under a grant number 977205 and Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. The study team would like to gratefully acknowledge the assistance and sincere cooperation of staff at Diagnostic Laboratory Sciences and Technology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. We would like to thank Fereshteh Jafaripour for her expert advice and brilliant help in the case of designing the schematic figures of SOEing PCR as well.

References

1. Higuchi R, Krummel B, Saiki R. A general method of *in vitro* preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions. *Nucleic Acids Res* 1988;16(15):7351-7367.
2. Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* 1989;77(1):51-59.
3. Horton RM, Hunt HD, Ho SN, Pullen JK, Pease LR. Engineering hybrid genes without the use of restriction en-

- zymes: gene splicing by overlap extension. *Gene* 1989; 77(1):61-68.
4. Mullis KB, Faloona FA. Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol* 1987;155:335-350.
 5. Mullis K, Faloona F, Scharf S, Saiki R, Horn G, editors. Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. Cold Spring Harbor symposia on quantitative biology; 1986: Cold Spring Harbor Laboratory Press.
 6. Charlier N, Molenkamp R, Leyssen P, Vandamme AM, De Clercq E, Bredenbeek P, et al. A rapid and convenient variant of fusion-PCR to construct chimeric flaviviruses. *J Virol Methods* 2003;108(1):67-74.
 7. Goh KM, Liew KJ, Chai KP, Ilias RM. Use of megaprimer and overlapping extension PCR (OE-PCR) to mutagenize and enhance cyclodextrin glucosyltransferase (CGTase) function. *Methods Mol Biol* 2017;498:385-396.
 8. You BJ, Lee MH, Chung KR. Gene-specific disruption in the filamentous fungus *Cercospora nicotianae* using a split-marker approach. *Arch Microbiol* 2009;191(7):615-622.
 9. Nakamura M, Suzuki A, Hoshida H, Akada R. Minimum GC-rich sequences for overlap extension PCR and primer annealing. *Methods Mol Biol* 2014;1116:165-181.
 10. Shevchuk NA, Bryksin AV, Nusinovich YA, Cabello FC, Sutherland M, Ladisch S. Construction of long DNA molecules using long PCR-based fusion of several fragments simultaneously. *Nucleic Acids Res* 2004;32(2):e19.
 11. Luo WG, Liu HZ, Lin WH, Kabir MH, Su Y. Simultaneous splicing of multiple DNA fragments in one PCR reaction. *Biol Proced Online* 2013;15(1):9.
 12. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;131(5):861-872.
 13. Tanabe K, Nakamura M, Narita M, Takahashi K, Yamana S. Maturation, not initiation, is the major roadblock during reprogramming toward pluripotency from human fibroblasts. *Proc Natl Acad Sci USA* 2013;110(30):12172-12179.
 14. Matsuda T, Cepko CL. Controlled expression of transgenes introduced by in vivo electroporation. *Proc Natl Acad Sci USA* 2007;104(3):1027-1032.
 15. Yon J, Fried M. Precise gene fusion by PCR. *Nucleic Acids Res* 1989;17(12):4895.
 16. Geu-Flores F, Nour-Eldin HH, Nielsen MT, Halkier BA. USER fusion: a rapid and efficient method for simultaneous fusion and cloning of multiple PCR products. *Nucleic Acids Res* 2007;35(7):e55.
 17. Stein V, Hollfelder F. An efficient method to assemble linear DNA templates for in vitro screening and selection systems. *Nucleic Acids Res* 2009;37(18):e122.
 18. Ge L, Rudolph P. Simultaneous introduction of multiple mutations using overlap extension PCR. *Biotechniques* 1997;22(1):28-30.
 19. Kim YG, Maas S. Multiple site mutagenesis with high targeting efficiency in one cloning step. *Biotechniques* 2000;28(2):196-198.
 20. Smith ZD, Sindhu C, Meissner A. Molecular features of cellular reprogramming and development. *Nat Rev Mol Cell Biol* 2016;17(3):139-154.
 21. Zou S, Scarfo K, Nantz MH, Hecker JG. Lipid-mediated delivery of RNA is more efficient than delivery of DNA in non-dividing cells. *Int J Pharm* 2010;389(1-2):232-243.
 22. Bernal JA. RNA-based tools for nuclear reprogramming and lineage-conversion: towards clinical applications. *J Cardiovasc Transl Res* 2013;6(6):956-968.