

The effects of extraction technique on phenolic compounds extracted from fig (*Ficus carica*) pulp and skin

E. Maghsoudlou¹, R. Esmailzade Kenari^{2*}, Z. Raftani Amiri³

Received: 2015.11.28

Accepted: 2016.06.20

Abstract

Recently, Subcritical Water Extraction (SWE) has been well known as a green technology for extraction of bioactive compounds from plants. In this study, Subcritical water extraction, ultrasound assisted extraction (UAE) and shaker solvent extraction (SSE) were compared for extraction of phenolic compounds from fig (*Ficus carica*) pulp and skin. Antioxidant activity of the extracts was evaluated using DPPH radical scavenging, reducing power and rancimat tests. Subcritical water had the highest ability for extraction of total phenolic content (65.89 ± 0.21 and 80.79 ± 0.09 mg of gallic acid equivalents per gram of extract respectively) and flavonoid compounds (7.51 ± 0.33 and 10.1 ± 1.02 mg of quercetin equivalents per gram of extract, respectively) from both pulp and skin. The lowest IC₅₀ in DPPH radical scavenging and reducing power tests were related to SWE of skin extract of fig. Furthermore, in extraction of total phenol and flavonoid compounds, subcritical water extraction showed to be a more suitable method than other solvent extraction methods, both in pulp and skin.

Keywords: Antioxidant activity; Fig; Extraction; Ultrasound; Subcritical water extraction

Introduction

The Common Fig (*Ficus carica* L.) is a tree native to the Middle East and the Mediterranean region, which belongs to botanical family Moraceae. The Common (Flaishman *et al.*, 2008; Oliveira *et al.*, 2009) fig is an excellent source of minerals, vitamins and dietary fiber; fat and cholesterol-free and contain a high number of amino acids (Caliskan, 2015; Viuda-Martos *et al.*, 2015). In addition of several health benefits which were previously reported (Lansky *et al.*, 2008; Viuda-Martos *et al.*, 2015; Weli *et al.*, 2015), figs are an excellent source of phenolic compounds.

Polyphenols are natural antioxidants and the most abundant secondary metabolites of plants that possess interesting properties, such as free-radical scavenging and inhibition of various oxidative stress in the body (Dai and Mumper, 2010). Plant-derived antioxidants are

molecules, which donate electrons or hydrogen atoms. These compounds are able to form less reactive antioxidant-derived radicals, which are efficiently quenched by other electrons or hydrogen sources to prevent cellular damage therefore, they may delay and inhibit lipid oxidation, protect human cells against oxidative damage, lead to a reduced risk of several oxidative-stress associated degenerative diseases, such as cancer, cardiovascular or neurodegenerative diseases (Scalbert *et al.*, 2005). When added to foods, these antioxidant compounds tend to minimize rancidity, retard the formation of toxic oxidation products, help to maintain the nutritional quality and increase their shelf life (Fukumoto and Mazza, 2000).

Phenolic compounds from plants has been traditionally extracted using solvent extraction or steam distillation techniques. Traditional methods of extraction are cost and time consuming protocols, and require large volumes of solvents (Teixeira *et al.*, 2006). Recently several new methods have been applied for plant phenolic extraction such as supercritical fluid extraction (SFE) (Herrero *et al.*, 2010; McHugh and Krukonis, 2013; Pereira and Meireles, 2010), pulse electric

1, 2 and 3: Former MSc student, Assistant and Associate Professors, Department of Food Science and Technology, Sari Agricultural Sciences and Natural Resources University, Sari, Mazandaran, Iran, respectively.

(* - Corresponding Author Email: reza_kenari@yahoo.com)

field assisted extraction (PEFE)(Vorobiev et al., 2005), high pressure assisted extraction (HPE)(Corrales et al., 2008) and matrix solid-phase dispersion (MSPD) (Capriotti et al., 2010; Dawidowicz and Rado, 2010), which are less labor intensive and more environmentally friendly. Despite the use of this new extraction techniques solid-liquid extraction (SLE) is still commonly used.

Subcritical water extraction (SWE), also called pressurized polarity water extraction or superheated water extraction, is a technique using water as extraction solvent at temperatures between 100 and 374°C with a pressure high enough to maintain water in liquid state (Ma et al., 2015). Previous studies have proved that it is a promising technique because of its short extraction time, decrease the organic solvent consumption, less volatile oil loss and increase the quality of extracts (Jayawardena and Smith, 2010; Kumar et al., 2011; Rodríguez-Meizoso et al., 2010).

In recent years, ultrasound-assisted extraction method has become an effective method for edible oils and fats from natural product extraction UAE is an inexpensive, simple and efficient alternative to conventional extraction techniques (Wang and Weller, 2006). The mechanism of UAE is attributed to mechanical and cavitation efficacies which can result in disruption of cell wall, particle size reduction, and enhanced mass transfer across cell membrane (Wang *et al.*, 2013). The objective of this study is to compare antioxidant activity of the phenols extracted from the pulp and skin of fig by ultrasound-assisted aqueous extraction, Subcritical water extraction and solvent extraction

Materials and methods

Material

Fig fruit (*F. carica L.*) from Siyah variety collected from Gorgan city on September 2014. Canola oil was purchased from Alia Golestan company (Kordkooy,iran) All other chemicals used in this study were of analytical grade and were purchased from chemical suppliers.

Preparation of extracts

The figs (Siyah variety) were weighed and immediately peeled. The pulp was cut and made into flat sheets. Thereafter, the pulp and skin of fruit were shade-dried for 5 days followed by drying at 60°C in an oven for 24 hours to ensure complete drying (Memmert 100-800, Germany). The samples were then milled and sieved through No. 67. Samples obtained were kept in polyethylene bags

Solvent extraction

Ten grams of each sample was mixed with water-ethanol (70%) in a ratio of 1 to 10 and gently stirred. For a better extraction, the mixture was shaken (120 rpm) at dark at 25 °C for 24 hours on a shaker, then the supernatant was filtered by Buchner funnel and Whatman filter paper No. 1. The extracts containing solvent was poured in a glass plate and placed in oven at 40°C for 24 hours. After evaporation of the solvent, the extracts were placed in desiccator until constant weight and kept at -18°C for further analyses (Esmailzadeh Kenari *et al.*, 2014).

Ultrasound assistant extraction

Dried powders of sample (10 g) were mixed with ethanol (1:10). The mixture was sonicated for 20 min at 40 °C in an ultrasonic bath (Elma s 30 H model, total power consumption: 280 W, heating power: 200 W, operating at 37 KHz frequency and internal dimensions: 198 × 106 × 50 cm). The temperature was controlled and maintained at 40 °C by circulating water. The extract was filtered and subsequently evaporated using a rotary evaporator. The concentrated extracts were stored at -18°C until further analyses (Esmailzadeh Kenari *et al.*, 2014).

Subcritical water extraction

Subcritical water extraction was carried out by a system consists of a distilled water tank, a pump (Comet type:MTP AX 2/70 m) providing pressure up to 170±5 bar, an extraction cell with 140 ml capacity, a heating coil, a pressure gauge and a temperature

control device. The powdered sample was loaded into the cell. Extraction was carried out at temperature of 160°C for 30 minutes (Hassas-Roudsari et al., 2009). The impurities of liquid (extract) was removed using filter paper (Whatman paper No. 4) under vacuum condition. The filtered extracts were cooled and stored at -18°C in dark polypropylene bags until used in the analysis (Shaddel *et al.*, 2014; Sharifi *et al.*, 2013).

Determination of total phenolic content

Total phenolic content of each extract was determined by the Folin–Ciocalteu micro-method according to revised methods of Javanmardiet al. (2003). Briefly, in a 50 ml volumetric flask, 1 ml of a standard solution of gallic acid, 6 ml of methanol, 2.5 ml of the Folin-Ciocalteu reagent, and 5 ml of 7.5% Na₂CO₃ were added. The final volume was achieved by addition of distilled water. The solutions were stored overnight and the spectrophotometric analysis was performed at $\lambda=765$ nm. (PG-instrument, USA). The total phenol content of samples and canola oil were expressed as gallic acid equivalents per g of extract using the following linear equation based on the calibration curve:

$$y = 0.0008x + 0.029 \quad R^2 = 0.094$$

Where y is absorbance at 765 nm and x is concentrations of gallic acid equivalents (mg/g) (Capannesi *et al.*, 2000; Javanmardi *et al.*, 2003).

Determination of total flavonoid content

Colorimetric aluminum chloride method was used for flavonoid determination. Briefly, 0.5 ml solution of each plant extracts in methanol were separately mixed with 1.5 ml of methanol, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate, and 2.8 ml of distilled water, and left at room temperature for 30 minutes. The absorbance of the reaction mixture was measured at 415 nm with a double beam Perkin Elmer UV/Visible spectrophotometer (PG-instrument, USA). The calibration curve was prepared by preparing pure quercetin solutions at concentrations 12.5

to 100 mg/ml in methanol. Total flavonoid content was calculated as quercetin per g of extract using the following linear equation based on the calibration curve:

$$y = 0.0064x + 0.0124 \quad R^2 = 0.9982$$

Where y is absorbance at 415 nm and x is concentrations of flavonoids compounds (mg quercetin/g extract) (Nabaviet al., 2012).

DPPH Radical-Scavenging Activity

Stable 2, 2'-diphenyl-1-picryl hydrazyl radical (DPPH) was used for determination of free radical-scavenging activity of the extracts. Different concentrations of each extracts were added, at an equal volume, to methanol solution of DPPH (100 μ M). The samples were kept at room temperature in darkness and after 15 min the absorbance of each sample was measured at 517 nm and the percentage of scavenging activity was calculated from the equation 1. The experiment was repeated for three times. IC₅₀ values denote the concentration of sample, which is required to scavenge 50% of DPPH free radicals (Aksoy *et al.*, 2013).

$$\text{DPPH scavenging activity (\%)} = \frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \quad (1)$$

Reducing power

Ability of extracts to reduce iron (III) was evaluated using the method of Yildirim *et al.* (2001). Samples (2.5 ml) were mixed with 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide (K₃Fe (CN)₆; 10 g/L) and incubated for 30 min at 50°C. Then 2.5 ml of trichloroacetic acid (100 /L) were added to the solution and centrifuged for 10 min. Finally, 2.5 ml of supernatant was combined with 2.5 ml of distilled water and 0.5 ml FeCl₃ (1 g/L). The absorbance of samples was measured at 700 nm. Higher absorbance means higher reducing power (Yildirim *et al.*, 2001).

Determination of the thermal oxidative stability index (Rancimat test)

One mg/ml of each sample was mixed with 100 ml pure canola oil (without antioxidant) and then oxidative stability measured by Rancimat (Metrohm, 743, Switzerland) test based on the AOCS method (2007). The air flow rate and the temperature was set at 20 m³/h and at 110 °C, respectively.

Statistical analysis

Each experiment was carried out at least in duplicate and measurement performed at least in triplicate. Statistical analysis of data was performed using Microsoft Excel. Analysis of variance was calculated using the SPSS program with a confidence level of 0.05, to find any significant difference between treatments. Duncan multiple range test (MRT)

was used for mean separation at $P < 0.05$ where treatment effect was significant

Result and discussion

The effect of extraction on phenolic and flavonoid compounds

Analysis of variance (ANOVA) showed that the techniques used for extraction of phenolic compounds in skin and pulp of the fig were significant different ($P < 0.05$). Subcritical water method in both skin and pulp samples showed the highest amount of total phenolic (80.79±0.09 and 65.89±0.21 mg gallic acid/g of extract, respectively) and flavonoid (10.1±1.02 and 7.51±0.33 mg quercetin/g of extract, respectively) contents whereas there was no difference between UAE and SSE (Table 1).

Table 1- Total phenol and flavonoids of pulp and skin of fig extract with different extraction methods

Extraction technique	Total Phenol mg gallic acid/g of extract		Total Flavonoid mg quercetin/g of extract	
	pulp	skin	pulp	skin
SSE	58.35±0.05 ^b	68.59±1.02 ^b	2.04±0.13 ^b	4.29±0.12 ^b
UAE	58.84±0.13 ^b	70.09±0.32 ^b	2.06±0.23 ^b	4.32±0.12 ^b
SWE	65.89±0.21 ^a	80.79±0.09 ^a	7.51±0.33 ^a	10.1±1.02 ^a

Means with different letters within column indicate significance difference at $P < 0.05$.

SSE: Shaker Solvent Extraction; UAE: Ultrasonic Assisted- Extraction; SWE: Subcritical Water Extraction

Flavonoids are strong inhibitors of hydroxyl and peroxide radicals. These compounds can influence on free radicals even when they form complexes with metal ions (Koda *et al.*, 2008). Flavonoid compounds have pharmaceutical, antibacterial and anti-oxidation properties. Quercetin is a flavonol which is one of the most powerful natural antioxidants and a major component known in figs (Solomon *et al.*, 2006). Antioxidant properties of flavonoids, especially quercetin, are rendered through metal ion chelating, radical scavenging, and stimulating the expression of protective enzymes (Baghel *et al.*, 2012). Therefore, measurement of these compounds in figs deems very important.

Higher efficiency of SWE in the extraction of flavonoids also reported by earlier studies (Ko *et al.*, 2011; Liang and Fan, 2013; Rangsrivong *et al.*, 2009). Extraction of a

wide range of flavonoids by common extraction methods is limited due to poor solubility in water. By application of subcritical water, the polarity, viscosity, surface tension, and disassociation constant of subcritical water are significantly lowered compared to water at ambient temperature and pressure, with more similar chemical properties to those of organic solvents. The temperature and pressure of subcritical water extraction provide higher penetration rate that improves its efficiency (Rangsrivong *et al.*, 2009). Higher efficiency of SWE also could possibly be related to the effect of hydrolysis reaction caused by the increase in the ionization constant (K_w) of water at subcritical conditions.

Extraction time considerably decreased in ultrasonic process (from 24 h in SSE to 20 min in UAE) due to the positive effect of cavitation

although no significant difference was detected between the methods in terms of amount of phenolic and flavonoid compounds. Solid-liquid extraction is assisted by ultrasounds leading to formation of cavitation bubbles. The collapse of these bubbles near the cell walls is expected to cause cell disruption along with good penetration of the solvent into the cells through the ultrasonic jet action. This process leads to an intensification of mass transfer and improved solvent penetration into the plant tissue (Da Porto *et al.*, 2013). Nevertheless, extraction conditions (e.g. time, temperature, solvent) greatly influence on the efficiency of extraction of phenolic compounds and flavonoids in plant tissue. Similar results which compared ultrasound and solvent methods, were in consistent with our results (Chemat *et al.*, 2004; Da Porto *et al.*, 2013; Kimbaris *et al.*, 2006; Milić *et al.*, 2013; Vinatoru *et al.*, 1997).

The effect of extraction on DPPH Radical-Scavenging Activity

DPPH scavenging activity assay is widely used to evaluate the ability of compounds to scavenge free radicals or donate hydrogen, and determine the antioxidant activity in foods (Bidchol *et al.*, 2011). Subcritical water extraction of fig pulp at concentration of 1 mg/ml showed the highest radical scavenging activity ($65.66 \pm 1.54\%$) while a sharp decline in antioxidant activity was observed with higher concentrations (e.g. 1.5 and 2 mg/ml) (Figure 1). This extracts showed the highest inhibition at lower concentrations. While the other extracts with increasing concentrations, increased antioxidant activity. Subcritical water extraction of fig skin at concentrations of 0.5, 1, 1.5 and 2 mg/ml and UAE and SSE extract at concentrations of 2.5 and 3 mg/ml had the highest percentage of inhibition. Radical scavenging activity in the UAE- and SSE-extracts from both skin and pulp of fig was dependent on concentration where a higher radical scavenging was achieved with higher concentrations of phenolic and flavonoid compounds; this effect might be owing to higher amount of hydroxyl groups and consequently, increased probability of

hydrogen donation to free radicals (Sanchez-Moreno *et al.*, 1999). However, radical scavenging activity of subcritical water extracts decreased with higher extract concentrations. A logic behind this observation is the reaction between released chemicals in solvent leading to formation of new compounds and interference in identification of target compounds in higher concentration (Plaza *et al.*, 2010).

IC₅₀ is defined as a concentration of the extract required to scavenge 50% of DPPH radicals. Subcritical water extract of skin revealed the lowest IC₅₀ (0.45 ± 0.02 mg/ml); in other word, it is the best extract to scavenge free radicals. On the contrary, subcritical water extract of pulp had significantly higher IC₅₀ (0.65 ± 0.09 mg/ml) values than other treatments (figure 2).

The effect of extraction on reducing power

Reducing property is generally defined as the ability of donating a hydrogen atom and thereby breaking a radical chain. Furthermore, reluctant react with peroxide precursors and prevent the formation of peroxides [45]. Thus, samples with higher reducing powers are more able to donate electrons. The reducing powers of different samples are shown in Fig. 3 (a,b). In all concentrations, SWE of fig pulp and skin had the highest reducing power, UAE and SSE had no significant difference. The results showed that the compounds in fig skin extracts were good electron or hydrogen donors and could successfully terminate radical chain reactions. So, it can be considered a good alternative to synthetic antioxidants in the diet. These results were consistent with Gou *et al.* (2003) who evaluated the antioxidant activity of the skin and the pulp and kernel of twenty-eight common fruit in China via reducing power analysis and reported that fruit peels and kernels had higher antioxidant activity than pulps (Guo *et al.*, 2003). The results also showed the advantages of SWE because with higher temperature, the dielectric constant of water reduces leading to lower polarity of water. Therefore, compounds with different polarities including a variety of

flavonoids can be extracted via SWE, which increases the reducing power of extracts. Increasing concentrations lead to increase in the reducing power of extracts due to the increased amount of phenolic compounds present in the extract. These data largely confirm the DPPH test. The only difference is that a better distinction of antioxidant activities of extracts with varying concentrations is

achieved through reducing power assay. Since this method is usually used to measure the antioxidant capacity of hydrophilic compounds (Pérez-Jiménez et al., 2008) and also because of the nature of hydrophilic compounds in the figs extracts (due to the presence of anthocyanins), this method seems more accurate to measure the antioxidant activity fig extract.




Fig. 1. DPPH scavenging activity of fig extract with different extraction methods: (a) fig pulp and (b) fig skin (SSE: Shaker Solvent Extraction; UAE: Ultrasonic Assisted Extraction; SWE: Subcritical Water Extraction)




Fig. 2. IC₅₀ of extracts in DPPH assay with different extraction methods (SSE: Shaker Solvent Extraction; UAE: Ultrasonic Assisted Extraction; SWE: Subcritical Water Extraction)




Fig. 3. Reducing power of fig extract with different extraction methods: (a) pulp and (b) skin (SSE: Shaker Solvent Extraction; UAE: Ultrasonic Assisted Extraction; SWE: Subcritical Water Extraction)

Figure 4 shows IC_{50} in reducing power assay (mg/ml). In the case of reducing power, the IC_{50} value presents the concentration at which the absorbance is 0.5. Similar to DPPH assay, Subcritical water extracts in both skin and pulp samples showed significantly the lowest IC_{50} (0.90 ± 0.1 and 0.96 ± 0.05 mg/ml) and the highest reducing power.

Antioxidant activity of the extracts in edible oil was measured based on the electrical conductivity of water in addition to the accumulation of volatile compounds from oxidation, especially carboxylic acids under accelerated oxidation. Oxidative stability index (OSI) is defined as oil stability time at a given temperature. The results has shown that the effect of the extraction method was significant on oxidative stability and thus antioxidant activity ($P < 0.05$). In this study, UAE showed higher stability among other extracts (Figure 5). This can be due to differences in methods of extraction and the resulting difference is in the type of phenolic compounds in the extracts. These factors are important in solubility of the extracts in oil and hence in the oxidative stability of the oil.

Conclusion

The result presented in this study show that the subcritical water extraction method is an alternative for extraction of several phenolic and flavonoid compounds from skin and pulp of fig, and more efficient than both solvent extraction method and ultrasound. Subcritical water extracts of skin and pulp of fig showed higher antioxidant activities in the DPPH radical scavenging and reducing power assays, while in rancimat test ultrasound method was found to yield more antioxidant extracts. This difference can be justified by varying nature of the methods of measuring antioxidant activity. Additional research and development studies are necessary to thoroughly compare subcritical extraction with other recent methods.

Acknowledgment

The authors acknowledge department of food science and technology of Sari University of Agricultural Sciences and Natural Resources for providing financial support to this research work.




Fig. 4. IC_{50} of extracts in reducing power assay with different extraction methods (SSE: Shaker Solvent Extraction; UAE: Ultrasonic Assisted Extraction; SWE: Subcritical Water Extraction)




Fig 5- Oxidative stability index of pulp and skin extract with different extraction (SSE: Shaker Solvent Extraction; UAE: Ultrasonic Assisted Extraction; SWE: Subcritical Water Extraction)

References

- Aksoy, L., Kolay, E., Ağılonü, Y., Aslan, Z., Kargioğlu, M.(2013). Free radical scavenging activity, total phenolic content, total antioxidant status, and total oxidant status of endemic *Thermopsis turcica*. *Saudi Journal of Biological Sciences* 20(3), 235-239.
- AOCS (2007). Official methods and recommended practices of the American Oil Chemist's Society (7th ed). Champaign: *American Oil Chemists-Society*.
- Baghel, S.S., Shrivastava, N., Baghel, R.S., Agrawal, P., Rajput, S.(2012). A review of quercetin: antioxidant and anticancer properties. *World J Pharm Pharmaceut Sci* 1, 146-160.
- Bidchol, A.M., Wilfred, A., Abhijna, P., Harish, R.(2011). Free radical scavenging activity of aqueous and ethanolic extract of *Brassica oleracea L. var. italica*. *Food and Bioprocess Technology* 4(7), 1137-1143.
- Caliskan, O. (2015). Chapter 56 - Mediterranean Figs (*Ficus carica L.*) Functional Food Properties, in: Watson, V.R.P.R. (Ed.), *The Mediterranean Diet*. Academic Press, San Diego, pp. 629-637.
- Capannesi, C., Palchetti, I., Mascini, M., Parenti, A.(2000). Electrochemical sensor and biosensor for polyphenols detection in olive oils. *Food Chemistry* 71(4), 553-562.
- Capriotti, A.L., Cavaliere, C., Giansanti, P., Gubbiotti, R., Samperi, R., Laganà, A.(2010). Recent developments in matrix solid-phase dispersion extraction. *Journal of Chromatography A* 1217(16), 2521-2532.
- Chemat, S., Lagha, A., AitAmar, H., Bartels, P.V., Chemat, F. (2004). Comparison of conventional and ultrasound-assisted extraction of carvone and limonene from caraway seeds. *Flavour and Fragrance Journal* 19(3), 188-195.
- Corrales, M., Toepfl, S., Butz, P., Knorr, D., Tauscher, B.(2008). Extraction of anthocyanins from grape by-products assisted by ultrasonics, high hydrostatic pressure or pulsed electric fields: a comparison. *Innovative Food Science & Emerging Technologies* 9(1), 85-91.
- Da Porto, C., Porretto, E., Decorti, D.(2013). Comparison of ultrasound-assisted extraction with conventional extraction methods of oil and polyphenols from grape (*Vitis vinifera L.*) seeds. *Ultrasonics sonochemistry* 20(4), 1076-1080.
- Dai, J., & Mumper, R. J. (2010). Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313-7352.
- Dawidowicz, A.L., Rado, E. (2010). Matrix solid-phase dispersion (MSPD) in chromatographic analysis of essential oils in herbs. *Journal of pharmaceutical and biomedical analysis* 52(1), 79-85.
- Esmailzadeh Kenari, R., Mohsenzadeh, F., Amiri, Z.R.(2014). Antioxidant activity and total phenolic compounds of Dezful sesame cake extracts obtained by classical and ultrasound-assisted extraction

- methods. *Food Science & Nutrition* 2(4), 426-435.
- Flaishman, M.A., Rodov, V., Stover, E.(2008). The fig: botany, horticulture, and breeding. *Horticultural Reviews-Westport Then New York-* 34, 113.
- Fukumoto, L., Mazza, G.(2000). Assessing antioxidant and prooxidant activities of phenolic compounds. *Journal of Agricultural and Food Chemistry* 48(8), 3597-3604.
- Guo, C., Yang, J., Wei, J., Li, Y., Xu, J., Jiang, Y. (2003). Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutrition Research* 23(12), 1719-1726.
- Hassas-Roudsari, M., Chang, P. R., Pegg, R. B., Tyler, R.T. (2009). Analytical Methods-Antioxidant capacity of bioactives extracted from canola meal by subcritical water, ethanolic and hot water extraction, *Food Chemistry*, 114: 717-726.
- Herrero, M., Mendiola, J.A., Cifuentes, A., Ibáñez, E.(2010). Supercritical fluid extraction: Recent advances and applications. *Journal of Chromatography A* 1217(16), 2495-2511.
- Javanmardi, J., Stushnoff, C., Locke, E., Vivanco, J. M. (2003). Antioxidant activity and total phenolic content of Iranian *Ocimum* accessions. *Food Chemistry* 83(4), 547-550.
- Jayawardena, B., Smith, R. M. (2010). Superheated water extraction of essential oils from *Cinnamomum zeylanicum* (L.). *Phytochemical Analysis* 21(5), 470-472.
- Kimbaris, A.C., Siatis, N.G., Daferera, D.J., Tarantilis, P.A., Pappas, C. S., Polissiou, M. G.(2006). Comparison of distillation and ultrasound-assisted extraction methods for the isolation of sensitive aroma compounds from garlic (*Allium sativum*). *Ultrasonics sonochemistry* 13(1), 54-60.
- Ko, M. J., Cheigh, C. I., Cho, S. W., Chung, M. S. (2011). Subcritical water extraction of flavonol quercetin from onion skin. *Journal of Food Engineering* 102(4), 327-333.
- Koda, T., Kuroda, Y., Imai, H.(2008). Protective effect of rutin against spatial memory impairment induced by trimethyltin in rats. *Nutrition Research* 28(9), 629-634.
- Kumar, M.Y., Dutta, R., Prasad, D., Misra, K. (2011). Subcritical water extraction of antioxidant compounds from Seabuckthorn (*Hippophae rhamnoides*) leaves for the comparative evaluation of antioxidant activity. *Food Chemistry* 127(3), 1309-1316.
- Lansky, E.P., Paavilainen, H. M., Pawlus, A. D., Newman, R. A.(2008). *Ficus* spp.(fig): Ethnobotany and potential as anticancer and anti-inflammatory agents. *Journal of Ethnopharmacology* 119(2), 195-213.
- Liang, X., Fan, Q.(2013). Application of sub-critical water extraction in pharmaceutical industry. *Journal of Materials Science and Chemical Engineering* 1(05), 1.
- Ma, Q., Fan, X. D., Liu, X. C., Qiu, T. Q., Jiang, J. G.(2015). Ultrasound-enhanced subcritical water extraction of essential oils from *Kaempferia galangal* L. and their comparative antioxidant activities. *Separation and Purification Technology* 150, 73-79.
- McHugh, M., Krukonis, V. (2013). *Supercritical fluid extraction: principles and practice*. Elsevier.
- Milić, P.S., Stanojević, L.P., Rajković, K.M., Milić, S.M., Nikolić, V.D., Nikolić, L.B., Veljković, V. B.(2013). Antioxidant activity of *Galium mollugo* L. extracts obtained by different recovery techniques. *Hemjska industrija* 67(1), 89-94.
- Nabavi, S.M., Nabavi, S.F., Ebrahimzadeh, M. A.(2012). Free radical scavenging and antioxidant activities of *Dorema aitchisonii*. *Journal of Food and Drug Analysis* 20(1).
- Oliveira, A.P., Valentão, P., Pereira, J.A., Silva, B.M., Tavares, F., Andrade, P. B.(2009). *Ficus carica* L.: Metabolic and biological screening. *Food and Chemical Toxicology* 47(11), 2841-2846.
- Pereira, C.G., Meireles, M.A. A.(2010). Supercritical fluid extraction of bioactive compounds: fundamentals, applications and economic perspectives. *Food and Bioprocess Technology* 3(3), 340-372.
- Pérez-Jiménez, J., Arranz, S., Taberner, M., Díaz-Rubio, M.E., Serrano, J., Goñi, I., Saura-Calixto, F.(2008). Updated methodology to determine antioxidant capacity in plant foods, oils and beverages: Extraction, measurement and expression of results. *Food Research International* 41(3), 274-285.
- Plaza M, Amigo-Benavent M, Del Castillo MD, Ibanez E, Herrero M. (2010). Facts about the formation of new antioxidants in natural samples after subcritical water extraction. *Food Res. Int.* 43: 2341-2348.
- Rangsriwong, P., Rangkadilok, N., Satayavivad, J., Goto, M., Shotipruk, A.(2009). Subcritical water extraction of polyphenolic compounds from *Terminalia chebula* Retz. fruits. *Separation and Purification Technology* 66(1), 51-56.
- Rodríguez-Meizoso, I., Jaime, L., Santoyo, S., Señoráns, F. J., Cifuentes, A., Ibáñez, E.(2010). Subcritical water extraction and characterization of bioactive compounds from *Haematococcus pluvialis* microalga. *Journal of pharmaceutical and biomedical analysis* 51(2), 456-463.

- Sanchez-Moreno, C., Larrauri, J. A., Saura-Calixto, F. (1999). Free radical scavenging capacity and inhibition of lipid oxidation of wines, grape juices and related polyphenolic constituents. *Food Research International*.32:407-412.
- Scalbert, A., Manach, C., Morand, C., Rémésy, C., Jiménez, L.(2005). Dietary polyphenols and the prevention of diseases. *Critical reviews in food science and nutrition* 45(4), 287-306.
- Shaddel, R., Maskooki, A., Haddad-Khodaparast, M.H., Azadmard-Damirchi, S., Mohamadi, M., Fathi-Achachlouei, B.(2014). Optimization of extraction process of bioactive compounds from Bene hull using subcritical water. *Food Science and Biotechnology* 23(5), 1459-1468.
- Sharifi, A., Mortazavi, S., Maskooki, A., Niakousari, M., Elhamirad, A.(2013). Optimization of Subcritical Water Extraction of Bioactive Compounds from Barberry Fruit (*Berberis vulgaris*) by Using Response Surface Methodology. *International Journal of Agriculture and Crop Sciences* 6(2), 89.
- Solomon, A., Golubowicz, S., Yablowicz, Z., Grossman, S., Bergman, M., Gottlieb, H.E., Altman, A., Kerem, Z., Flaishman, M. A.(2006). Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.). *Journal of Agricultural and Food Chemistry* 54(20), 7717-7723.
- Teixeira, D.M., Patão, R.F., Coelho, A.V., da Costa, C. T.(2006). Comparison between sample disruption methods and solid-liquid extraction (SLE) to extract phenolic compounds from *Ficus carica* leaves. *Journal of Chromatography A* 1103(1), 22-28.
- Vinatoru, M., Toma, M., Radu, O., Filip, P., Lazurca, D., Mason, T. (1997). The use of ultrasound for the extraction of bioactive principles from plant materials. *Ultrasonics sonochemistry* 4(2), 135-139.
- Viuda-Martos, M., Barber, X., Pérez-Álvarez, J.A., Fernández-López, J. (2015). Assessment of chemical, physico-chemical, techno-functional and antioxidant properties of fig (*Ficus carica* L.) powder co-products. *Industrial Crops and Products* 69, 472-479.
- Vorobiev, E., Jemai, A.B., Bouzrara, H., Lebovka, N., Bazhal, M.(2005). Pulsed electric field assisted extraction of juice from food plants. *Novel food processing technologies*, 105-130.
- Wang, L. & Weller, C.L. (2006). Recent advances in extraction of nutraceuticals from plants. *Trends Food Sci. Technol*, 17, 300-12.
- Wang, X., Wu, Y., Chen, G., Yue, W., Liang, Q., Wu, Q.(2013). Optimisation of ultrasound assisted extraction of phenolic compounds from *Sparganii rhizoma* with response surface methodology. *Ultrasonics sonochemistry* 20(3), 846-854.
- Weli, A.M., Al-Blushi, A.A.M., Hossain, M. A.(2015). Evaluation of antioxidant and antimicrobial potential of different leaves crude extracts of Omani *Ficus carica* against food borne pathogenic bacteria. *Asian Pacific Journal of Tropical Disease* 5(1), 13-16.
- Wong, J.Y., Chye, F.Y. (2009). Antioxidant properties of selected tropical wild edible mushrooms. *Journal of Food Composition and Analysis* 22(4), 269-277.
- Yildirim, A., Mavi, A., Kara, A. A.(2001). Determination of antioxidant and antimicrobial activities of *Rumex crispus* L. extracts. *Journal of Agricultural and Food Chemistry* 49(8), 4083-4089.

اثر روش‌های استخراج بر ترکیبات فنولی استخراج‌شده از پوست و پالپ انجیر

الهه مقصدلو^۱ - رضا اسماعیل‌زاده کناری^{۲*} - زینب رفتنی امیری^۳

تاریخ دریافت: ۱۳۹۴/۰۹/۰۷

تاریخ پذیرش: ۱۳۹۵/۰۳/۳۱

چکیده

اخیرا استخراج آب زیر بحرانی (SWE) به عنوان یک تکنولوژی سبز برای استخراج ترکیبات زیست فعال گیاهان شناخته شده است. در این مطالعه، استخراج آب زیر بحرانی، استخراج به کمک اولتراسوند (UAE) و استخراج حلالی همزنی (SSE) جهت استخراج ترکیبات فنولی پوست و پالپ انجیر مقایسه شدند. فعالیت آنتی‌اکسیدانی عصاره‌ها با آزمون‌های مهار رادیکال DPPH، قدرت احیاکنندگی و رنسیت ارزیابی شد. آب زیر بحرانی نشان داد که بالاترین توانایی را در استخراج ترکیبات فنولی کل پالپ و پوست (به ترتیب، $65/89 \pm 0/21$ و $80/79 \pm 0/09$ معادل میلی گرم گالیک اسید در گرم عصاره) و ترکیبات فلاونوئیدی (به ترتیب، $7/51 \pm 0/33$ و $10/11 \pm 1/02$ معادل میلی گرم کوئرستین در گرم عصاره) دارد. همچنین، کمترین IC50 در آزمون‌های مهار رادیکال DPPH و قدرت احیاکنندگی به عصاره‌ی آب زیر بحرانی پوست انجیر مربوط بود. بنابراین، آب زیر بحرانی روش مناسب‌تری برای استخراج ترکیبات فنولی و فلاونوئیدی کل پوست و پالپ انجیر می‌باشد.

واژه‌های کلیدی: فعالیت آنتی‌اکسیدانی، انجیر، استخراج، اولتراسوند، استخراج آب زیر بحرانی

۱، ۲ و ۳. دانش‌آموخته کارشناسی ارشد، ستادیار و دانشیار گروه علوم و صنایع غذایی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ساری، مازندران، ایران.
* - نویسنده مسئول : (Email: reza_kenari@yahoo.com)