



## Thin-layer convective air drying of lemon verbena (*Lippia citriodora*) leaves

E. Naghavi<sup>1\*</sup>, S. Rigi<sup>2</sup>

Received: 2016.12.15

Accepted: 2017.04.22

### Abstract

Lemon verbena leaf is a flavoring food additive as well as a good source of valuable compounds such as essential oils, flavonoids and phenolic acids. However, similar to many other aromatic plants, lemon verbena leaf is perishable due to its high moisture content. The aim of this work was to study the effect of air temperature (45, 55, and 65°C) on the quality attributes of lemon verbena leaves during hot-air drying (HAD). The drying kinetics were also modeled. The results showed that higher drying temperature led to a significant decrease ( $p < 0.05$ ) in the rehydration ratio due to a change in the structural features of the dried leaves. The essential oil content of dried samples was also significantly different ( $p < 0.05$ ) from that of the fresh leaves due to high loss of volatile components and ranged from 0.42 to 0.85. Moreover, a significant increase in the value of effective moisture diffusivity ( $D_{eff}$ ) and color change was observed when the samples were dried at 65°C compared to 45°C. The value of  $D_{eff}$  varied from  $1.140 \times 10^{-10}$  to  $2.280 \times 10^{-9}$  m<sup>2</sup>/s and the activation energy was found to be 31.04 kJ/mol. The greatest  $R^2$  ( $\geq 0.999$ ) and the lowest RMSE and SSE were obtained for the Naghavi *et al.* model (proposed in this research)

**Keywords:** Color change, convective drying, effective moisture diffusivity, essential oil, modeling, lemon verbena leaves, rehydration ratio

### Introduction

Lemon verbena (*Lippia citriodora*) is a type of herb which is widely raised in western South America. It is also cultivated in Iran and mainly consumed as a spice and a medicinal plant (Funes *et al.*, 2009). There is an increasing interest in using lemon verbena leaf in the food industry, because it is generally considered as a flavoring food additive. The leaves of lemon verbena have compounds such as essential oils, flavonoids and phenolic acids, which possess antioxidant activity (Pereira *et al.*, 2007). They are mainly used to make herbal teas and refreshing sorbets as well as creating a lemon flavor in a number of food products such as fish and poultry dishes, jams, salad dressings, puddings, and beverages (Funes *et al.*, 2009). Moreover, the leaves have digestive, sedative,

antispasmodic, stomachic, and antipyretic properties (Pereira *et al.*, 2007; Funes *et al.*, 2009). However, similar to many other aromatic plants and herbs, lemon verbena leaves are perishable to microbial growth, mainly due to their high moisture content (around 84-85% wet basis).

Drying is used to extend the shelf life of fruits, vegetables and aromatic plants as well as to reduce or suppress their enzymatic and microbial activities (Doymaz 2009; Doymaz 2012). Aromatic plants are dried in order to extract their valuable compounds by solvents. Among the drying methods, hot-air drying (HAD) is still the most popular method, which is being employed to decrease the moisture content of foods and plants. Although HAD is time and energy consuming (Erbay and Icier 2010), it has gained considerable attention by researchers due to its low capital cost compared to other drying techniques such as freeze-drying and infrared-drying. For this reason, it is still extensively employed by many researchers for long-term preservation of foods and herbs (Erbay and Icier 2010; Doymaz 2012; Lemus-Mondaca *et al.*, 2015;

1. Young Researchers and Elite Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran.

2. Department of Food Science and Technology, Islamic Azad University, Sabzevar Branch, Sabzevar, Iran.

(Corresponding Author Email: E\_naghavi@tabrizu.ac.ir)

DOI: 10.22067/ifstrj.v12i6.61049

Oberoi and Sogi 2015; Aral and Beşe 2016; Nozad *et al.*, 2016; Roshanak *et al.*, 2016; Salarikia *et al.*, 2016).

Effective moisture diffusivity ( $D_{\text{eff}}$ ) and activation energy ( $E_a$ ) are two important physical properties of dehydrated foods which represent the rate of moisture loss during HAD and the level of energy needed to initiate a chemical reaction and to activate moisture diffusion, respectively (Lemus-Mondaca *et al.*, 2015).  $D_{\text{eff}}$  can highly affect the drying kinetics and consequently the quality of dehydrated foodstuffs. Also, mathematical modeling of mass transfer during HAD requires the values of  $D_{\text{eff}}$ . On the other hand, rehydration ratio (RR) is a key physical characteristic of dried foods which can reflect the degree of textural damage (such as shrinkage and tissue collapse) to foodstuffs during HAD (Doymaz *et al.*, 2015).

Numerous research papers can be found on the determination of  $D_{\text{eff}}$ ,  $E_a$ , and RR during HAD of various aromatic plants and herbs (Doymaz 2012; Tasirin *et al.*, 2014; Lemus-Mondaca *et al.*, 2015; Nozad *et al.*, 2016; Salarikia *et al.*, 2016). However, to the best of our knowledge, no study has been conducted on the determination of  $D_{\text{eff}}$ ,  $E_a$ , RR, and color change for lemon verbena leaves under HAD conditions. The purpose of this research was to investigate the influence of hot-air temperature on the drying kinetics, color change, RR, and essential oil content of lemon verbena leaves under HAD and to calculate  $D_{\text{eff}}$  and  $E_a$ , as well as empirical modeling of the dimensionless moisture ratio as a function of drying time.

## Materials and methods

### Materials

Fresh lemon verbena leaves were collected every morning from a farm located in Tabriz (Iran), and immediately transferred to the laboratory. The leaves were sorted visually based on size, shape, color, and freshness and stored under refrigerated conditions (at 5°C) (Lemus-Mondaca *et al.*, 2015) until use. The initial moisture content of the leaves was

measured using the AOAC method (AOAC 1984) and found to be equal to 84.72% (wet basis).

### Hot-air drying

First, the leaves were removed from the refrigerator and arranged uniformly as a thin layer in a stainless steel basket. Then, they were dried using the HAD technique. The experiments were carried out in a pilot plant hot-air drier (UOP 8 Tray dryer, Armfield, UK, equipped with automatic data recording system and temperature and airflow velocity controller units) at 45, 55, and 65±1°C and the airflow rate of 1 m/s (Doymaz 2012; Lemus-Mondaca *et al.*, 2015). Moisture loss was calculated by measuring the mass loss of the samples at 15 min intervals (based on preliminary experiments) by a precision balance with an accuracy of ±0.01 g. Moisture content data were recorded throughout the drying experiments using a data logger connected to a PC. The experiments were continued until reaching a final moisture content of 10% (wet basis).

### Modeling of drying curves

Eighteen different empirical and semi-empirical models were used to evaluate the kinetics of moisture loss during HAD of lemon verbena leaves (Table 1) (Ertekin and Heybeli 2014). The model parameters or drying constants (a, b, c, g, h, k, and n) were estimated by applying non-linear regression analysis using MATLAB software (Version 8.1.0.604 R2013a, The Math works, Inc., USA). The coefficient of determination ( $R^2$ ), adjusted  $R^2$ , root mean squared error (RMSE), and sum of squared error (SSE) were used to evaluate the goodness of fit in order to select the suitable model(s) to predict the drying kinetics. These statistical criteria are as follows (Lemus-Mondaca *et al.*, 2015):

$$R^2 = 1 - \frac{\sum_{i=1}^N (MR_{\text{exp},i} - MR_{\text{pre},i})^2}{\sum_{i=1}^N (\overline{MR_{\text{exp}}} - MR_{\text{pre},i})^2} \quad (1)$$

$$RMSE = \left( \frac{1}{N} \sum_{i=1}^N (MR_{exp,i} - MR_{pre,i})^2 \right)^{\frac{1}{2}} \quad (2)$$

$$SSE = \frac{1}{N} \sum_{i=1}^N (MR_{exp,i} - MR_{pre,i})^2 \quad (3)$$

Where  $MR_{exp,i}$  is the  $i^{th}$  experimental

moisture ratio,  $MR_{pre,i}$  shows the  $i^{th}$  predicted moisture ratio,  $\overline{MR}_{exp}$  stands for the average experimental moisture ratio, and N denotes the number of observations or the number of data values.

Table 1. Kinetic models used to describe the drying of lemon verbena leaves\*

Model number	Model equation	Model name
1	$MR = \exp(-kt)$	Lewis (Newton)
2	$MR = \exp(-kt^n)$	Page
3	$MR = \exp(-(kt)^n)$	Modified Page-I
4	$MR = a \exp(-kt)$	Henderson & Pabis
5	$MR = a \exp(-kt^n)$	Modified Page-II
6	$MR = a \exp(-kt^n) + bt$	Midilli et al.
7	$MR = a \exp(-kt)^n + b$	Demir et al.
8	$MR = (a - b) \exp(-kt^n)$	Weibull distribution-I
9	$MR = (a - b) \exp(-(kt)^n)$	Weibull distribution-II
10	$MR = \exp\left(\frac{-at}{1+bt}\right)$	Aghlasho
11	$MR = \frac{a}{1 + b \exp(kt)}$	Logistic
12	$MR = a \exp(-kt) + b \exp(-gt)$	Two-term
13	$MR = a \exp(-kt^n) + b \exp(-gt^n)$	Hii et al.
14	$MR = a \exp(-kt) + (1 - a) \exp(-kat)$	Two-term exponential
15	$MR = a \exp(-kt) + (1 - a) \exp(-bt)$	Modified two-term exponential
16	$MR = a \exp(-kt) + (1 - a) \exp(-kbt)$	Diffusion approximation
17	$MR = (1 - a) \exp(-kt) + (1 - b) \exp(-gt^n) + c$	Naghavi et al. (present study)
18	$MR = a \exp(-kt) + b \exp(-gt) + c \exp(-ht)$	Modified Henderson & Pabis

\*All models (except model-17) are available in the paper published by Ertekin and Heybeli (2014). a, b, c, g, h, k, and n are model parameters (empirical constants).

**Determination of the effective moisture diffusivity**

The Fick's law-based model (Eq. 4) is often used to determine the effective moisture diffusivity ( $D_{eff}$ ) of different food materials.

$$\frac{\partial M}{\partial t} = D_{eff} \frac{\partial^2 M}{\partial x^2} \quad (4)$$

The following initial and boundary conditions can be considered (Doymaz 2012):

$$\begin{aligned} t = 0, & \quad 0 < x < L, & M &= M_0 \\ t > 0, & \quad x = L, & M &= M_e \end{aligned}$$

$$t > 0, \quad x = 0, \quad \frac{dM}{dx} = 0$$

In the present study, the analytical solution of Fick's second law for an infinite slab (Eq. 5) was applied to calculate  $D_{eff}$  (Erbay and Icier 2010; Doymaz 2012):

$$MR = \frac{M_t - M_e}{M_0 - M_e} = \frac{8}{\pi^2} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \exp\left(-\frac{(2n+1)^2 \pi^2 D_{eff} t}{4L^2}\right) \quad (5)$$

where M is the moisture content (dry basis),  $D_{eff}$  represents the effective moisture diffusivity ( $m^2/s$ ), L is the half thickness of the

slab (m),  $t$  stands for the time (s),  $MR$  is the moisture ratio (dimensionless),  $M_t$  shows the moisture content at any time (kg water/kg dry solid),  $M_0$  is the initial moisture content (kg water/kg dry solid),  $M_e$  denotes the equilibrium moisture content (kg water/kg dry solid), and  $n$  is the number of the terms taken into consideration.

This method has also been used previously by several researchers (Doymaz 2012; Aral and Beşe 2016) and is based on the assumptions that shrinkage is negligible,  $D_{eff}$  remains constant and moisture loss occurs through the diffusion phenomenon (Crank 1975). For long drying times ( $M_e=0$ ), the use of one-term approximation ( $n=1$ ) to the series summation is reasonable and Eq. 5 reduces to (Doymaz 2012):

$$MR = \frac{M_t}{M_0} = \frac{8}{\pi^2} \exp\left(-\frac{\pi^2 D_{eff} t}{4L^2}\right) \quad (6)$$

By taking the natural logarithm, Eq. 6 can be further simplified to (Doymaz 2012; Oberoi and Sogi 2015):

$$\ln(MR) = \ln\left(\frac{8}{\pi^2}\right) - \left(\frac{\pi^2 D_{eff} t}{4L^2}\right) \quad (7)$$

A plot of the experimental data in terms of  $\ln(MR)$  versus time gives a straight line with a slope of  $k_0$  (Oberoi and Sogi 2015):

$$k_0 = \frac{-\pi^2 D_{eff}}{4L^2} \quad (8)$$

#### Determination of the activation energy

The Arrhenius model describes the relationship between  $D_{eff}$ , drying temperature ( $T$ ), and the activation energy ( $E_a$ ). Therefore, for quantifying  $E_a$  and investigating the effect of temperature on  $D_{eff}$ , the Arrhenius type equation (Eq. 9) was employed (Doymaz 2012):

$$D_{eff} = D_0 \exp\left(-\frac{E_a}{RT}\right) \quad (9)$$

Where  $E_a$  denotes the activation energy (kJ/mol),  $R$  is the universal gas constant [8.314 J/(mol K)],  $T$  is the absolute air temperature (K), and  $D_0$  is the pre-exponential factor (constant) ( $m^2/s$ ). This approach has been used

by a number of researchers (Doymaz 2012; Aral and Beşe 2016).

#### Determination of the rehydration ratio

Measurement of the rehydration ratio (RR) of dried leaves was carried out according to the method described by Doymaz *et al.* (2015) and Nozad *et al.* (2016). Based on this method, 5 g of the dehydrated samples were poured into a glass beaker (750 mL) containing 500 mL of distilled water (25°C) and kept for 24 h. Next, the leaves were removed from the beaker and their surface water was blotted up using a tissue paper. Finally, the weight of the resulted sample was measured precisely using a digital balance. In all cases, the tests were triplicated for each sample and the mean values of the three replications  $\pm$  standard deviation were reported. The RR calculation was carried out using Eq. 10 as employed by Doymaz *et al.* (2015), Nozad *et al.* (2016), and Salarikia *et al.* (2016):

$$RR = \frac{W_2 - W_1}{W_1} \quad (10)$$

Where RR denotes the rehydration ratio [kg water/kg dry matter (DM)],  $W_1$  is the weight of the dried leaves (kg), and  $W_2$  represents the weight of the rehydrated leaves (kg).

#### Color measurement

The color changes of leaves (fresh and hot-air dried) were quantified using image processing in MATLAB (Version 8.1.0.604 R2013a, The Math works, Inc., USA) (Nozad *et al.* 2016). The color test instrument was designed and constructed in the Department of Agricultural Machinery Engineering, University of Tabriz, Tabriz, Iran. It consists of a chamber with a trapezoidal cross section that was equipped by two  $D_{65}$  (daylight) lamps as the light source for illumination of sample. At first, a sample was put in the chamber. After zooming the lens and focusing, the images were taken by camera. A digital camera (Nikon, D3200, Japan) was used to capture images from leaf surfaces. The camera calibration was performed prior to each drying experiment.

In each experimental run, the color of the leaves (10 fresh and 10 dried samples) was measured as  $L^*$ ,  $a^*$  and  $b^*$  values, which known as Hunter parameters. It is well known that the  $L^*$  value represents the degree of lightness/darkness,  $a^*$  stands for the degree of redness/greenness, and  $b^*$  shows the degree of yellowness/blueness. Changes in the color of the leaves were calculated as follows (Nozad *et al.*, 2016; Salarikia *et al.*, 2016):

$$\Delta E = \sqrt{\Delta L^{*2} + \Delta a^{*2} + \Delta b^{*2}} \quad (11)$$

Where  $\Delta E$  denotes the total color change of leaves and  $\Delta L^*$ ,  $\Delta a^*$ , and  $\Delta b^*$  represents the difference between the color parameters of initial samples ( $L_0^*$ ,  $a_0^*$ , and  $b_0^*$ ) and final dried leaves ( $L^*$ ,  $a^*$ , and  $b^*$ ).

#### Determination of the essential oil content

The essential oil was extracted from the leaves using a flask connected to Clevenger hydro-distillation apparatus (Nozad *et al.*, 2016). Based on this method, a given amount of the samples (30 g) were put into a round-bottomed distillation flask filled with a given amount of distilled water (250 mL). Then, the

heating was performed for 3 h and the distilled essential oil collected in the side arm was separated. The data of essential oil (%) were expressed on the basis of dry matter weight.

#### Statistical analysis

The experimental data were analyzed statistically by the analysis of variance (ANOVA) using Minitab statistical software (Minitab Release 14, Minitab Inc., USA). The significant difference between the means was determined using Tukey's honestly significant difference (HSD) test at the significance level of 5% ( $p < 0.05$ ). The data were expressed as the mean  $\pm$  standard deviation and all experiments were carried out in triplicate.

#### Results and discussion

##### Drying kinetics

The effects of the drying temperature (45, 55, and 65°C) on the dimensionless moisture ratios of lemon verbena leaves are illustrated in Fig. 1.




Fig. 1. Moisture ratios (dimensionless) as a function of the drying time at different temperatures during hot-air drying of lemon verbena leaves

It can be seen that the moisture ratio declined quickly in the initial period of HAD (almost up to 75-90 min) and was subsequently followed by a gradual non-linear decrease (almost exponential) with an increase in the process time. This result was similar to the findings of other researchers (Doymaz

2012; Lemus-Mondaca *et al.*, 2015; Said *et al.*, 2015; Aral and Beşe 2016).

Fig. 1 also shows that all drying temperatures exhibited a relatively similar behavior for the dried samples. Moreover, by increasing the drying temperature, the moisture ratios and consequently the drying

kinetics were altered. Based on ANOVA results, the drying time was significantly ( $p < 0.05$ ) lower in the case of the samples dried at 65°C than those dehydrated at lower temperatures (45 or 55°C). The total drying time was 420, 345, and 285 min for the leaves dried at 45, 55, and 65°C, respectively. This result indicates that a temperature increase of 20°C (i.e. from 45 to 65°C) caused a reduction of approximately 135 min in the total drying time ( $p < 0.05$ ). This might be due to the increased vapor pressure in lemon verbena leaves at higher drying temperatures, which in turn results in a faster moisture loss from the samples and thus, a shorter drying time (Aral and Beşe 2016). Other studies have reported similar findings (Ertekin and Heybeli 2014; Lemus-Mondaca *et al.*, 2015; Said *et al.*, 2015; Aral and Beşe 2016).

### Modeling of the drying curves

In the literature, several empirical and semi-empirical models have been applied to describe the drying kinetics of food materials and plants under different drying conditions, which are based on the dimensionless moisture ratio as a function of drying time. In the present work, the experimental data of moisture loss in lemon verbena leaves during HAD at different drying temperatures (45, 55, and 65°C) were fitted to eighteen models summarized in Table 1 (Ertekin and Heybeli 2014), which allows of using and comparing different model correlations based on the fitting criteria ( $R^2$ , RMSE, and SSE) on a defined set of experimental data

**Table 2. Statistical analysis of different kinetic models for different drying temperatures**

Model number	T (°C)	R <sup>2</sup>	Adjusted R <sup>2</sup>	RMSE	SSE	Model number	T (°C)	R <sup>2</sup>	Adjusted R <sup>2</sup>	RMSE	SSE
1	45	0.9875	0.9875	0.0283	0.0225	10	45	0.9977	0.9976	0.0124	0.0041
	55	0.9870	0.9870	0.0294	0.0199		55	0.9981	0.9980	0.0116	0.0029
	65	0.9907	0.9907	0.0256	0.0125		65	0.9981	0.9980	0.0120	0.0026
2	45	0.9995	0.9995	0.0056	0.0008	11	45	0.9744	0.9735	0.0413	0.04611
	55	0.9996	0.9996	0.0050	0.0005		55	0.9723	0.9710	0.0439	0.04233
	65	0.9997	0.9997	0.0049	0.0004		65	0.9766	0.9753	0.0417	0.03132
3	45	0.9995	0.9995	0.0056	0.0008	12	45	0.9998	0.9998	0.0034	0.00028
	55	0.9996	0.9996	0.0050	0.0005		55	0.9998	0.9999	0.0032	0.00019
	65	0.9997	0.9997	0.0049	0.0004		65	0.9999	0.9998	0.0033	0.00017
4	45	0.9927	0.9924	0.0221	0.0131	13	45	0.9999	0.9999	0.0030	0.00021
	55	0.9916	0.9912	0.0241	0.0128		55	0.9997	0.9996	0.0051	0.00049
	65	0.9934	0.9931	0.0221	0.0088		65	0.9997	0.9996	0.0053	0.00042
5	45	0.9995	0.9995	0.0057	0.0008	14	45	0.9996	0.9996	0.0048	0.00063
	55	0.9996	0.9996	0.0051	0.0005		55	0.9995	0.9994	0.0061	0.00083
	65	0.9997	0.9996	0.0050	0.0004		65	0.9998	0.9998	0.0041	0.00030
6	45	0.9997	0.9997	0.0044	0.0005	15	45	0.9998	0.9998	0.0033	0.00029
	55	0.9997	0.9997	0.0046	0.0004		55	0.9999	0.9998	0.0032	0.00027
	65	0.9998	0.9997	0.0044	0.0003		65	0.9999	0.9998	0.0031	0.00026
7	45	0.9997	0.9997	0.0047	0.0005	16	45	0.9998	0.9998	0.0033	0.00030
	55	0.9997	0.9996	0.0048	0.0005		55	0.9999	0.9998	0.0030	0.00025
	65	0.9998	0.9997	0.0044	0.0003		65	0.9999	0.9999	0.0032	0.00017
8	45	0.9995	0.9995	0.0058	0.0008	17	<b>45</b>	<b>1</b>	<b>1</b>	<b>0.0007</b>	<b>0.000011</b>
	55	0.9996	0.9996	0.0052	0.0005		<b>55</b>	<b>1</b>	<b>0.9999</b>	<b>0.0013</b>	<b>0.000031</b>
	65	0.9997	0.9996	0.0052	0.0004		<b>65</b>	<b>0.9999</b>	<b>0.9999</b>	<b>0.0020</b>	<b>0.000056</b>
9	45	0.9995	0.9995	0.0058	0.0008	18	45	0.9998	0.9998	0.0035	0.00028
	55	0.9996	0.9996	0.0052	0.0005		55	0.9999	0.9998	0.0035	0.00022
	65	0.9997	0.9996	0.0052	0.0004		65	0.9997	0.9997	0.0049	0.00034

**Table 3. Estimated coefficients of different kinetics models at different drying temperatures\***

Model number	T (°C)	Coefficients		Model number	T (°C)	Coefficients	
1	45	a = 0.0102		10	45	a = 0.0124	b = 0.0014
	55	a = 0.0130			55	a = 0.0159	b = 0.0019
	65	a = 0.0177			65	a = 0.0212	b = 0.0022
2	45	a = 0.0230	n = 0.8315	11	45	a = 1.7520	k = 0.0128
	55	a = 0.0294	n = 0.8213		55	a = 1.7650	k = 0.0164
	65	a = 0.0360	n = 0.8346		65	a = 1.8030	k = 0.0229
3	45	a = 0.0107		12	45	a = 0.7404	g = 0.0350
	55	n = 0.8315			55	b = 0.2604	k = 0.0078
	65	a = 0.0137	n = 0.8213		65	a = 0.7025	g = 0.0402
4	45	a = 0.9339		13	45	b = 0.2977	k = 0.0096
	55	k = 0.0095			65	a = 0.7515	g = 0.0625
	65	a = 0.9370	k = 0.0121		65	b = 0.2490	k = 0.0138
5	45	a = 0.9500		14	45	a = 0.6834	k = 0.0051
	55	k = 0.0168			55	b = 0.3152	n = 1.0690
	65	a = 1.0020	n = 0.8298		65	g = 0.0268	
6	45	k = 0.0233		15	45	a = 1.0290	k = 0.0338
	55	a = 1.0020	n = 0.8195		55	b = -0.0292	n = 0.7970
	65	k = 0.0298			65	g = 3.1770	
7	45	a = 0.9993		16	45	a = 0.9833	k = 0.0334
	55	k = 0.0359	n = 0.8352		55	b = 0.0168	n = 0.8485
	65	a = 1.0060	k = 0.0259		65	g = 7.8360	
8	45	b = -3.089×10 <sup>-5</sup>	n = 0.8041	17	45	a = 0.2325	
	55	a = 1.0040	k = 0.0317		55	k = 0.0349	
	65	b = -2.328×10 <sup>-5</sup>	n = 0.8036		65	a = 0.2449	
9	45	a = 1.0010	k = 0.0380	18	45	k = 0.0417	
	55	b = -2.562×10 <sup>-5</sup>	n = 0.8199		55	a = 0.2307	
	65	a = 1.0210	k = 0.0104		65	k = 0.0612	
10	45	b = -0.0150	n = 0.8015	19	45	a = 0.2599	k = 0.0349
	55	a = 1.012	k = 0.0135		55	b = 0.0078	
	65	b = -0.0084	n = 0.8030		65	a = 0.2977	k = 0.0401
11	45	a = 1.0100	k = 0.0183	20	45	b = 0.0096	k = 0.0401
	55	b = -0.0087	n = 0.8154		55	a = 0.2487	k = 0.0624
	65	a = 0.7216	k = 0.0232		65	b = 0.0138	k = 0.0624
12	45	b = -0.2803	n = 0.8298	21	45	a = 0.2598	k = 0.0349
	55	a = 1.1360	k = 0.0298		55	b = 0.2246	k = 0.0349
	65	b = 0.1335	n = 0.8194		65	a = 0.2977	k = 0.0401
13	45	a = 0.5534	k = 0.0359	22	45	b = 0.2390	k = 0.0401
	55	b = -0.446	n = 0.8351		55	a = 0.2490	k = 0.0623
	65	a = -6.0690	k = 0.0108		65	b = 0.2214	k = 0.0623
14	45	b = -7.0710	n = 0.8298	23	45	a = 0.6936	g = 0.0114
	55	a = 1.3370	k = 0.0137		55	b = 0.2731	k = 0.0256
	65	b = 0.3346	n = 0.8195		65	c = -0.0333	n = 0.9001
15	45	a = 0.5256	k = 0.0186	24	45	a = 0.6157	g = 6.0080
	55	b = -0.4736	n = 0.8352		55	b = 1.9780	k = 0.0153
	65	a = 0.7216	k = 0.0232		65	c = 0.6156	n = -0.4296
16	45	b = -0.2803	n = 0.8298	25	45	a = 0.3155	g = 0.0001
	55	a = 1.1360	k = 0.0298		55	b = 0.6998	k = 0.0352
	65	b = 0.1335	n = 0.8194		65	c = 0.0144	n = 1.8710
17	45	a = 0.5534	k = 0.0359	26	45	a = 0.7466	g = 0.0373
	55	b = -0.446	n = 0.8351		55	b = 0.2643	h = 10.300
	65	a = -6.0690	k = 0.0108		65	c = -0.0109	k = 0.0079
18	45	b = -7.0710	n = 0.8298	27	45	a = -0.1141	g = 0.0495
	55	a = 1.3370	k = 0.0137		55	b = 0.4036	h = 0.0096
	65	b = 0.3346	n = 0.8195		65	c = 0.7106	k = 0.0802
19	45	a = 0.5256	k = 0.0186	28	45	a = 0.8052	g = 1.4230
	55	b = -0.4736	n = 0.8352		55	b = -1.424	h = 0.2176
	65	a = 0.7216	k = 0.0232		65	c = 1.6190	k = 0.0144

\* a, b, c, g, h, k, and n are model parameters (empirical constants).




Fig. 2. Comparison of the experimental (exp.) and predicted (pred.) moisture ratios for lemon verbena leaves dehydrated at different drying temperatures

Table 2 indicates that for the most of tested models,  $R^2$  and adjusted  $R^2$  values were higher than 0.999, and RMSE and SSE values were between 0.0007-0.0413 and 0.000011-0.04611, respectively.

Generally, the closer the experimental and predicted moisture ratios, the better they explain the adequacy of the regression model. As expected, the models with a larger number of coefficients (models 12, 13, 15-18 in Table 2) had higher  $R^2$  and lower RMSE and SSE values. The data in Table 2 suggested that the employed models were suitable to describe the drying behavior of lemon verbena leaves. However, for plotting the predicted moisture ratios against the drying time, only the model presented in this research (model-17 in Table 1) was fitted to the experimental data (Fig. 2) which had greater  $R^2$  (0.9999-1) and adjusted  $R^2$  (0.9999-1) values and lower RMSE (0.0007-0.0020) and SSE (0.000011-0.000056) values than the other 17 models (Table 2). The models coefficients (constants) obtained at different drying temperatures are represented in Table 3. By increasing drying temperature, different coefficients did not follow a similar trend. As can be obtained from Table 3, drying rate constant ( $k$ ) in the studied drying models increased with increasing drying temperatures. Thus, it may be assumed that this kinetic parameter would

be directly proportional to drying temperature (Lemus-Mondaca *et al.*, 2015). Similar results were reported by other investigators (Vega-Gálvez *et al.*, 2012; Lemus-Mondaca *et al.*, 2015). Furthermore, there was no clear trend on the effect of drying temperature on the other constants ( $a$ ,  $b$ ,  $c$ ,  $g$ ,  $h$ , and  $n$ ).

### Rehydration ratio (RR)

Comparison of the RR results for the samples dried at different drying temperatures are presented in Fig. 3.

It can be seen that there was no significant change ( $p > 0.05$ ) in the RR of the dried leaves when the drying temperature changed from 45 to 55°C. However, RR reduced significantly ( $p < 0.05$ ) with an increase in the air temperature from 45 to 65°C. This was attributed to the fact that HAD at higher temperatures resulted in a change in the structural features of the leaves (such as tissue collapse, development of a surface hard layer and volumetric shrinkage), which in turn can cause significant damages to the textural quality of the dried samples and therefore a decrease in RR (Doymaz 2012; Nozad *et al.*, 2016; Salarikia *et al.*, 2016). The highest RR value of the dried samples (78.24%) was observed for the leaves dried at 45°C, followed by those dehydrated at 55 (76.12%) and 65°C (73.36%). These results are in



agreement with the results reported by f Doymaz (2012), who stated that the higher the drying temperature (40, 50, and 60 °C) resulted in a lower RR of the grape leaves, concluding that higher drying temperatures led to greater changes in the structural attributes and thus, lower RR values. Nozad *et al.*, (2016) also observed that in HAD of

spearmint (*Mentha spicata* L.) leaves, an increase in the air temperature from 30 to 50°C had a considerable decreasing effect on RR. Similar findings were reported by other researchers (Jangam *et al.*, 2008; Vega-Gálvez *et al.*, 2012; Salarikia *et al.*, 2016).




Fig. 3. Rehydration ratio of lemon verbena leaves at different drying temperatures. Error bars show one standard deviation from the mean and means with different letters are significantly different ( $p < 0.05$ ).

#### Effective moisture diffusivity ( $D_{eff}$ )

In the present study, the value of  $D_{eff}$  at 45, 55, and 65°C was estimated by plotting  $\ln(MR)$  versus the drying time (Eq. 4) (Oberoi and Sogi 2015). The slope of the corresponding line represents the value of  $D_{eff}$ . The  $D_{eff}$  values varied from  $1.140 \times 10^{-10}$  to  $2.280 \times 10^{-9}$  m<sup>2</sup>/s (Table 4). Other researchers found similar values for the  $D_{eff}$  of dried foods (in particular leaf materials), which were in general in the range of  $10^{-12}$  to  $10^{-8}$  m<sup>2</sup>/s (Zogzas *et al.*, 1996).  $D_{eff}$  values have been determined for other herbs as follows: mint leaves  $3.067 \times 10^{-9}$  to  $1.941 \times 10^{-8}$  m<sup>2</sup>/s (Doymaz 2006), spinach leaves  $6.590 \times 10^{-10}$

to  $1.927 \times 10^{-9}$  m<sup>2</sup>/s (Doymaz 2009), nettle leaves  $1.744 \times 10^{-9}$  to  $4.992 \times 10^{-9}$  m<sup>2</sup>/s (Kaya and Aydin 2009), mint leaves  $0.965 \times 10^{-11}$  to  $1.190 \times 10^{-11}$  m<sup>2</sup>/s (Therdthai and Zhou 2009), olive leaves  $1.054 \times 10^{-9}$  to  $4.973 \times 10^{-9}$  m<sup>2</sup>/s (Erbay and Icier 2010), grape leaves  $4.13 \times 10^{-10}$  to  $1.83 \times 10^{-9}$  m<sup>2</sup>/s (Doymaz 2012), kaffir lime leaves  $2.61 \times 10^{-11}$  to  $9.24 \times 10^{-11}$  m<sup>2</sup>/s (Tasirin *et al.*, 2014), stevia leaves  $4.67 \times 10^{-10}$  to  $14.90 \times 10^{-10}$  m<sup>2</sup>/s (Lemus-Mondaca *et al.*, 2015), and wild edible plant (*Allium roseum*) leaves  $2.55 \times 10^{-12}$  to  $8.83 \times 10^{-12}$  m<sup>2</sup>/s (Said *et al.*, 2015).

Table 4. The values of effective moisture diffusivity ( $D_{eff}$ ) for lemon verbena leaves at different drying temperatures

Drying temperature (°C)	$D_{eff}$ (m <sup>2</sup> /s)
45	$1.140 \times 10^{-10}$ c
55	$1.710 \times 10^{-10}$ b
65	$2.280 \times 10^{-9}$ a

Different letters in the same column indicate significant differences ( $p < 0.05$ )

As can be realized from Table 4, significant

differences ( $p < 0.05$ ) of the  $D_{eff}$  values were

observed between the leaves dried at different drying temperatures. This indicates that hot-air temperature has a considerable effect on  $D_{\text{eff}}$  during HAD of plants and foods, as reported by numerous authors (Erbay and Icier 2010; Doymaz 2012; Lemus-Mondaca *et al.*, 2015; Said *et al.*, 2015; Aral and Beşe 2016). This may be related to the higher thermal energy transferring to the leaves at higher drying temperatures, which subsequently results in an increase in the kinetic energy of the water molecules (Aral and Beşe 2016). A higher  $D_{\text{eff}}$  value indicates the increasing rate of moisture loss with the rise of drying temperature (Fig. 1).

#### Activation energy ( $E_a$ )

The value of  $E_a$  in the drying of lemon verbena leaves was estimated from the slope of the linearized Arrhenius equation (Eq. 9) (Doymaz 2012) and was found to be 31.04 kJ/mol. The  $E_a$  value obtained in this research was in the range that reported for other aromatic plants and fruits. Experimentally-determined  $E_a$  values have been reported by several researchers, for example Ahmed *et al.* (2001) for coriander leaves (26.50 kJ/mol in the temperature range of 45-65°C), Doymaz (2006) for mint leaves (62.96 kJ/mol in the temperature range of 35-60°C), Doymaz (2009) for spinach leaves (34.35 kJ/mol in the temperature range of 50-80°C), Kaya and Aydın (2009) for nettle leaves (79.873-109.003 kJ/mol in the temperature range of 35-55°C and at airflow rates of 0.2-0.6 m/s), Erbay and Icier (2010) for olive leaves (60.97 kJ/mol in the temperature range of 50-70°C), Doymaz (2012) for grape leaves (64.56 kJ/mol in the temperature range of 40-60°C), Lemus-Mondaca *et al.* (2015) for stevia leaves (38.78 kJ/mol in the temperature range of 30-80°C), and Said *et al.* (2015) for wild edible plant (*Allium roseum*) leaves (46.80-52.68 kJ/mol in the temperature range of 30-80°C and 1 and 1.5 m/s airflow velocity).

It has been reported that the value of  $E_a$  is influenced by several factors, including the drying air temperature, the moisture content of food or herb, and variations in the  $D_{\text{eff}}$  value

with the drying temperature (Aghbashlo *et al.*, 2008), which makes it difficult to compare the  $E_a$  values for different foods and herbs dehydrated at different process conditions. However, from numerous conducted studies on the calculation of the  $E_a$  value, it can be concluded that long dehydration time, high initial moisture content, remarkable variation in the  $D_{\text{eff}}$  value with the drying temperature (at constant airflow rate) (Aghbashlo *et al.*, 2008) or with both temperature and airflow rate (Erbay and Icier 2010), low hot-air flow rate, low drying temperature, and textural changes in the sample due to the percentage of shrinkage and tissue collapse, are all the reasons for a considerable increase in the  $E_a$  value.

#### Color measurement

The value of color change ( $\Delta E$ ) represents the degree of total color change in dehydrated leaves compared to the color of fresh samples. The lower  $\Delta E$  the better the quality of dried leaves (Salarikia *et al.*, 2016). The color of aromatic plants and herbs are very sensitive to heat damage during HAD. Fig. 4 shows the value of  $\Delta E$  for dried samples. It can be seen that the value of  $\Delta E$  increase significantly ( $p < 0.05$ ) with increasing of hot-air temperature from 45 to 65°C. Chlorophyll a and chlorophyll b are responsible for natural green color of leaves. The increase in  $\Delta E$  is due to the increase in substitution of magnesium with hydrogen in chlorophyll with drying temperature. Under this condition, chlorophylls are converted to pheophytins (Therdthai and Zhou 2009). This finding confirmed the previous observations obtained by Therdthai and Zhou (2009) for mint leaves (*Mentha cordifolia* Opiz ex Fresen), Chenarbon *et al.* (2012) for St. John's wort (*Hypericum perforatum* L.) leaves, Akbudak and Akbudak (2013) for parsley, and Salarikia *et al.* (2016) for peppermint leaves.

#### Determination of the essential oil content

Essential oil content of lemon verbena leaves before (fresh sample) and after drying is shown in Fig. 5. The fresh samples had the

highest value of essential oil (1.10%) between the treatments. Furthermore, no significant difference ( $p>0.05$ ) of essential oil content was observed between the leaves dried at different temperatures, while significant difference ( $p<0.05$ ) of essential oil content was observed between fresh leaves and samples dehydrated at 55 and 65°C. The reduction in essential oil content with increasing drying temperature ( $p>0.05$ ) might be explained by the fact that the relatively high temperature of hot-air result

in an increase in rupture of oil glands and as a consequence, rapid evaporation or higher loss of volatile components (Argyropoulos and Müller 2014). This result is consistent with those obtained in previous studies, for example *Laurus nobilis* L. leaves (Sellami *et al.*, 2011), *Thymys daenensis* subsp. *daenensis*. Celak leaves (Rahimmalek and Goli, 2013), Lemon verbena (Shahhoseini *et al.*, 2013), and peppermint leaves (Salarikia *et al.*, 2016).




Fig. 4. Color change in lemon verbena leaves at different drying temperatures. Error bars show one standard deviation from the mean and means with different letters are significantly different ( $p<0.05$ ).




Fig. 5. Essential oil content of lemon verbena leaves at different drying temperatures. Error bars show one standard deviation from the mean and means with different letters are significantly different ( $p<0.05$ ).

## Conclusions

This study was focused on investigation of

some selected properties of lemon verbena leaves (moisture loss,  $D_{eff}$ , color change, essential oil content, and RR) during HAD at

different drying temperatures as well as the empirical modeling of the drying kinetics. A shorter drying time and a higher  $D_{\text{eff}}$  value were observed ( $p < 0.05$ ) with an increase in the drying temperature from 45 to 65°C. The values of  $D_{\text{eff}}$  ranged from  $1.140 \times 10^{-10}$  to  $2.280 \times 10^{-9}$  m<sup>2</sup>/s and the  $E_a$  value was found to be 31.04 kJ/mol, all of which were in agreement with the results reported by other investigators in the literature. Our results also showed that the percentage of RR was significantly ( $p < 0.05$ ) affected by the air temperature and its maximum (78.24%) and minimum (73.36%) values were attained at 45°C and 65°C, respectively. Essential oil content of the samples dried at different drying

temperatures was not significant ( $p > 0.05$ ) with respect to each other but was significantly ( $p < 0.05$ ) different from the fresh samples. Furthermore, the value of  $\Delta E$  increased significantly ( $p < 0.05$ ) with increasing of hot-air temperature from 45 to 65°C and ranged from 12.24 to 14.92, respectively. The results of modeling ( $R^2 > 0.99$  and low RMSE and SSE values for most of the tested models) indicated a good fit to the experimental data of moisture ratio. Among these, the model proposed in the present study had a better goodness of fit (with an adjusted  $R^2 \geq 0.999$  and the lowest RMSE and SSE) and was considered as the best model.

## References

- Aghbashlo, M., & Samimi-Akhijahani, H., 2008, Influence of drying conditions on the effective moisture diffusivity, energy of activation and energy consumption during the thin-layer drying of berberis fruit (*Berberidaceae*). *Energy Conversion and Management*, 49(10): 2865-2871.
- Ahmed, J., Shivhare, U., & Singh, G., 2001, Drying characteristics and product quality of coriander leaves. *Food and Bioprocess Technology*, 79(2): 103-106.
- Akbudak, N., & Akbudak, B., 2013, Effect of vacuum, microwave, and convective drying on selected parsley quality. *International Journal of Food Properties*, 16(1): 205-215.
- AOAC, 1984. Official Methods of Analysis, Association of the Official Analytical Chemists, Washington, DC.
- Aral, S., & Beşe, A. V., 2016, Convective drying of hawthorn fruit (*Crataegus* spp.): Effect of experimental parameters on drying kinetics, color, shrinkage, and rehydration capacity. *Food Chemistry*, 210: 577-584.
- Argyropoulos, D., & Müller, J., 2014, Changes of essential oil content and composition during convective drying of lemon balm (*Melissa officinalis* L.). *Industrial Crops and Products*, 52: 118-124.
- Chenarbon, H. A., Minaei, S., Bassiri, A. R., Almassi, M., Arabhosseini, A., & Motevali, A., 2012, Effect of drying on the color of St. John's wort (*Hypericum perforatum* L.) leaves. *International Journal of Food Engineering*, 8(4), doi: 10.1515/1556-3758.2545.
- Crank, J., 1975. The mathematics of diffusion, 2nd ed., Oxford University Press, London.
- Doymaz, İ., 2006, Thin-layer drying behaviour of mint leaves. *Journal of Food Engineering*, 74(3): 370-375.
- Doymaz, I., 2009, Thin-layer drying of spinach leaves in a convective dryer. *Journal of Food Process Engineering*, 32(1): 112-125.
- Doymaz, I., 2012, Air drying characteristics, effective moisture diffusivity and activation energy of grape leaves. *Journal of Food Processing and Preservation*, 36(2): 161-168.
- Doymaz, I., Kipcak, A. S., & Piskin, S., 2015, Characteristics of thin-layer infrared drying of green bean. *Czech Journal of Food Sciences*, 33(1): 83-90.
- Erbay, Z., & Icier, F., 2010, Thin-layer drying behaviors of olive leaves (*Olea europaea* L.). *Journal of Food Process Engineering*, 33(2): 287-308.
- Ertekin, C., & Heybeli, N., 2014, Thin-layer infrared drying of mint leaves. *Journal of Food Processing and Preservation*, 38(4): 1480-1490.
- Funes, L., Fernández-Arroyo, S., Laporta, O., Pons, A., Roche, E., Segura-Carretero, A., Fernández-Gutiérrez, A., & Micol, V., 2009, Correlation between plasma antioxidant capacity and verbascoside levels in rats after oral administration of lemon verbena extract. *Food Chemistry*, 117(4): 589-598.
- Jangam, S. V., Joshi, V. S., Mujumdar, A. S., & Thorat, B. N., 2008, Studies on dehydration of sapota (*Achras zapota*). *Drying Technology*, 26(3): 369-377.

- Kaya, A., & Aydın, O., 2009, An experimental study on drying kinetics of some herbal leaves. *Energy Conversion and Management*, 50(1): 118-124.
- Lemus-Mondaca, R., Vega-Gálvez, A., Moraga, N. O., & Astudillo, S., 2015, Dehydration of Stevia rebaudiana Bertoni leaves: kinetics, modeling and energy features. *Journal of Food Processing and Preservation*, 39(5): 508-520.
- Nozad, M., Khojastehpour, M., Tabasizadeh, M., Azizi, M., Ashtiani, S.-H. M., & Salarikia, A., 2016, Characterization of hot-air drying and infrared drying of spearmint (*Mentha spicata* L.) leaves. *Journal of Food Measurement and Characterization*, 10(3): 1-8.
- Oberoi, D. P. S., & Sogi, D. S., 2015, Drying kinetics, moisture diffusivity and lycopene retention of watermelon pomace in different dryers. *Journal of food science and technology*, 52(11): 7377-7384.
- Pereira, C. G., & Meireles, M. A. A., 2007, Evaluation of global yield, composition, antioxidant activity and cost of manufacturing of extracts from lemon verbena (*Aloysia triphylla* [L'Herit.] Britton) and mango (*Mangifera indica* L.) leaves. *Journal of Food Process Engineering*, 30(2): 150-173.
- Rahimmalek, M., & Goli, S. A. H., 2013, Evaluation of six drying treatments with respect to essential oil yield, composition and color characteristics of Thymys daenensis subsp. daenensis. *Celak leaves. Industrial Crops and Products*, 42: 613-619.
- Roshanak, S., Rahimmalek, M., & Goli, S. A. H., 2016, Evaluation of seven different drying treatments in respect to total flavonoid, phenolic, vitamin C content, chlorophyll, antioxidant activity and color of green tea (*Camellia sinensis* or *C. assamica*) leaves. *Journal of food science and technology*, 53(1): 721-729.
- Said, L. B. H., Najjaa, H., Farhat, A., Neffati, M., & Bellagha, S., 2015, Thin layer convective air drying of wild edible plant (*Allium roseum*) leaves: experimental kinetics, modeling and quality. *Journal of food science and technology*, 52(6): 3739-3749.
- Salarikia, A., Miraei Ashtiani, S. H., & Golzarian, M. R., 2016, Comparison of drying characteristics and quality of peppermint leaves using different drying methods. *Journal of Food Processing and Preservation*, doi: 10.1111/jfpp.12930 (in press).
- Sellami, I. H., Wannas, W. A., Bettaieb, I., Berrima, S., Chahed, T., Marzouk, B., & Limam, F., 2011, Qualitative and quantitative changes in the essential oil of *Laurus nobilis* L. leaves as affected by different drying methods. *Food Chemistry*, 126(2): 691-697.
- Shahhoseini, R., Ghorbani, H., Karimi, S. R., Estaji, A., & Moghaddam, M., 2013, Qualitative and quantitative changes in the essential oil of lemon verbena (*Lippia citriodora*) as affected by drying condition. *Drying Technology*, 31(9): 1020-1028.
- Tasirin, S. M., Puspasari, I., Lun, A., Chai, P., & Lee, W., 2014, Drying of kaffir lime leaves in a fluidized bed dryer with inert particles: Kinetics and quality determination. *Industrial Crops and Products*, 61: 193-201.
- Therdthai, N., & Zhou, W., 2009, Characterization of microwave vacuum drying and hot air drying of mint leaves (*Mentha cordifolia* Opiz ex Fresen). *Journal of Food Engineering*, 91(3): 482-489.
- Vega-Gálvez, A., Ah-Hen, K., Chacana, M., Vergara, J., Martínez-Monzó, J., García-Segovia, P., Lemus-Mondaca, R., & Di Scala, K., 2012, Effect of temperature and air velocity on drying kinetics, antioxidant capacity, total phenolic content, colour, texture and microstructure of apple (var. *Granny Smith*) slices. *Food Chemistry*, 132(1): 51-59.
- Zogzas, N., Maroulis, Z., & Marinos-Kouris, D., 1996, Moisture diffusivity data compilation in foodstuffs. *Drying Technology*, 14(10): 2225-2253.

## خشک کردن همرفتی لایه نازک برگ‌های به لیمو

عنایت‌الله نقوی<sup>1\*</sup> - صادق ریگی<sup>2</sup>

تاریخ دریافت: 1395/09/25

تاریخ پذیرش: 1396/02/02

## چکیده

برگ به‌لیمو یک افزودنی غذایی طعم‌زا و همچنین منبع خوبی از ترکیبات با ارزش مانند روغن‌های فرار، فلاوونوئیدها و اسیدهای فنلی است. با این‌حال، همانند بسیاری از گیاهان معطر دیگر، برگ به‌لیمو به دلیل داشتن محتوای رطوبت بالا فسادپذیر است. هدف از این کار پژوهشی مطالعه اثر دمای هوا (45، 55 و 65 درجه سانتی‌گراد) روی ویژگی‌های کیفی برگ به‌لیمو طی خشک‌کردن هوای داغ (HAD) بود. همچنین، کینتیک خشک‌کردن مدل‌سازی شد. نتایج نشان داد که دمای خشک‌کردن بالاتر منجر به کاهش معنی‌دار ( $p < 0/05$ ) نسبت جذب آب مجدد به علت تغییر ویژگی‌های ساختاری برگ‌های خشک شده گردید. محتوای روغن فرار برگ‌های خشک شده نیز به دلیل از دست رفتن مقادیر بالای اجزای فرار به طور معنی‌داری ( $p < 0/05$ ) در مقایسه با برگ‌های تازه متفاوت بود و در محدوده 0/42 تا 0/85 قرار داشت. علاوه بر این، با خشک‌کردن نمونه‌ها در 65 درجه سانتی‌گراد در مقایسه با 45 درجه سانتی‌گراد، افزایش معنی‌دار مقدار ضریب انتشار مؤثر رطوبت ( $D_{eff}$ ) و تغییر رنگ مشاهده شد. مقدار  $D_{eff}$  از  $1/140 \times 10^{-10}$  تا  $2/280 \times 10^{-9}$  m/s متغیر بود و مقدار انرژی فعال‌سازی  $31/04$  kJ/mol تعیین شد. بیشترین مقدار  $R^2$  ( $\geq 0/999$ ) و کمترین مقدار RMSE و SSE برای مدل نقوی و همکاران (پیشنهاد شده در این پژوهش) به دست آمد.

واژه‌های کلیدی: تغییر رنگ، خشک‌کردن همرفتی، ضریب انتشار مؤثر رطوبت، روغن فرار، مدل‌سازی، برگ به‌لیمو، نسبت جذب آب مجدد

1- باشگاه پژوهشگران جوان و نخبگان، واحد تبریز، دانشگاه آزاد اسلامی، تبریز، ایران

2 - دانش‌آموخته کارشناسی ارشد، گروه علوم و صنایع غذایی، دانشگاه آزاد اسلامی واحد سبزوار، سبزوار، ایران

\* - نویسنده مسئول : (Email: E\_naghavi@tabrizu.ac.ir)