



Isolation and identification of antioxidants components From Cumin seed (*Cuminum cyminum*)

S. Einafshar^{1*}, H. Poorazarang², R. Farhoosh², J. Asili³

Received: 2016.01.25

Accepted: 2016.08.16

Abstract

Natural materials are complicated compositions; therefore, a rapid screening of the active antioxidants is being challenged in the literature. Different polarity solvents were applied to isolate four fractions (F1, F2, F3 and F4) of methanolic extract of *Cuminum cyminum*. Their anti-oxidative properties were tested using radical scavenging and FRAP assays. F3 (with $IC_{50}=0.006\text{mg/mL}$ and $FRAP=521.95\text{ mmolFe}^{2+}/\text{L}$) was significantly the most active fraction. Guided isolation through bio-autography on TLC using 1, 1-diphenyl-2-picrylhydrazyl radical (DPPH) as a detection reagent led to the isolation of two antioxidant compounds from F3. F3 was injected to a preparative HPLC with the proper mobile phase (acetonitril: methanol/ water) and isolated two main compounds. These compounds were identified as Luteolin 7 glucoside and Apigenin 7 glucoside by means of ¹HNMR and ¹³CNMR and compare them with references.

Key words: Antioxidant, Apigenin, *Cuminum cyminum*, Luteolin.

Introduction

A growing amount of evidence has shown that free radicals such as superoxide radical ($O_2^{\cdot-}$), hydroxyl radical (OH^{\cdot}), peroxy radical (ROO^{\cdot}), and nitric oxide radical (NO^{\cdot}), attack biological molecules, such as lipids, proteins, enzymes, DNA and RNA. They play an important role in several pathological processes, such as cancer, atherosclerosis, and negative cellular changes associated with aging (Fisch *et al.*, 2003; Nakman *et al.*, 2003; shon, *et al.*, 2003; Valentao, *et al.*, 2002). Antioxidants are effective in protecting the body against these degenerative events. Synthetic antioxidants, such as butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT) that are commonly

used in lipid-containing food have safety and toxicity problems (Amarowicz *et al.*, 2000; Howell, 1968). There is an increasing interest to seek for antioxidants naturally present in vegetables, fruits and functional herbs. Natural materials are complicated compositions therefore a rapidly screening of the active antioxidants is being challenged. A crude natural product extract is generally an extremely complicated mixture of several compounds possessing varying chemical and physical properties. The fundamental strategy for separating these compounds is based on their physical and chemical properties that can be effectively separated them into various chemical groups (Amarowicz *et al.*, 2000).

Plant natural products are usually extracted with solvents of increasing polarity, for instance, at first extracted with petroleum ether, dichloromethane followed by more polar solvents, i.e., ethyl acetate. Methanolic extracts of plant materials contain a wide range of polar and moderately polar compounds (otsuka, 2006). By virtue of the co solubility, many compounds, which are insoluble individually in pure state in methanol can be extracted quiet easily with these solvents. Herein we use an activity guided

1. Department of Agricultural Engineering Institute, Khorasan Razavi Agricultural and Natural Resources Research and Education Center, AREEO, Mashhad, IRAN.

2. Ferdowsi University of Mashhad, Faculty of Agriculture, Department of Food Science and Technology, Mashhad, Iran.

3. Medical science of Mashhad, Faculty of medicine, Department of pharmacognosy.

Corresponding Author Email: Soodabeheyn@yahoo.com

DOI: 10.22067/ifstrj.v1395i0.53312

purification of natural antioxidants from methanolic extract of cumin seed (*Cuminum cyminum*) by partitioning between insoluble solvents (Satti *et al.*, 2009).

Cumin seed L. is widely used as a spice. Crushed cumin seeds are used as a condiment in a variety of dishes. The proximate composition of the seeds indicates that it contains fixed oil (10%), protein, cellulose, sugar and other mineral elements (Winton, 1939). Cumin seed contains essential oil (2%–5%) that imparts the characteristic aroma to the seeds and the physicochemical properties of it have been reported (Guenther, 1950).

Phenolic compounds present in spices with natural antioxidant properties have been studied for substitution of synthetic antioxidants. Compare to many traditional medicinal plants, little is known about the active antioxidant matter of Cumin seed. Active principles of cuminaldehyde (cumin) are reported to inhibit lipid peroxidation (Shobana and Akhilender, 2000) and it has been reported that the ether extract of cumin seed inhibits biosynthesis of eicosanoids (Satti *et al.*, 2009) which are a pain mediators. Modulating effects of cumin seed was explored and its active compound, immunomodulatory properties were evaluated in normal and immune-suppressed animals (Chauhan, *et al.*, 2009). A preliminary study of cumin seed chemical composition was made on its essential oil had shown the presence of flavonoid compounds (Gachkar *et al.*, 2007; Li *et al.*, 2009; Behera, *et al.*, 2004) which act as antioxidants in edible oils (Vekiari *et al.*, 1993). Einafshar (2012) showed that *Cuminum cyminum* has antioxidative effects in both bulk and emulsion systems.

In the present study an activity guided purification was conducted to isolate the free radical scavenging compounds from Cumin seed, the bioactive constituents from the seeds of *Cuminum cyminum* was isolated and quantified and their chemical compositions identified.

Materials and Methods

Materials

Cumin seeds were obtained from international company of Pegah Saffron and Zire Green Gold (Mashhad, Iran). Refined, bleached, and deodorized sunflower oil (SFO) with no added antioxidant was supplied by Shadgol (Nishbour, Iran). All chemicals and solvents used in this study were of analytical reagent grade and supplied by Merck and Sigma Chemical Companies.

Plant material

The sample seeds were dried in a cabin drier at 55°C for 2 days, ground into a fine powder in a mill (Mulinex Depose-Brevete S.G.C.G., France). The powders were extracted with methanol (1:20 w/v) by agitation in a dark place at ambient temperature for 48 h. The solvent was evaporated in vacuum at 40°C. The desolventized extract (DEX) stored at -18°C until use.

Extraction and isolation of antioxidant compounds

60 mg the methanolic extract of *Cumin seed* was dissolved in 100 ml methanol and extracted thrice using 100 ml petroleum ether under stir for 10 minutes. The upper phase of petroleum ether, contains nonpolar compounds such as lipids, chlorophylls and so on. This process is sometimes referred to as defatting. Although methanol and petroleum ether are not completely miscible, they are miscible to some extent. A small amount of water is added to methanol to obtain a 95% aqueous methanolic solution to get two distinct layers with similar volumes. Petroleum ether in the upper phase was removed by evaporation at 50°C under vacuum to give a residue named F1. The lower phase is partitioned in Dichloromethane and upper phase in ethylacetate successively and named F2 and F3 respectively. Less polar compounds are present in the dichloromethane soluble fraction and polar compounds, probably up to mono-glycosides, in the ethylacetate. The methanol fraction contains polar compounds mainly glycosides named F4.

The conventional spectrophotometric DPPH• scavenging capacity assay

The radical-scavenging activity was evaluated according to Chung *et al.* Two ml of the methanol solution of F1, F2, F3 and F4 were mixed with 1 ml of 0.5 mM DPPH• methanol solution and 2 ml of 0.1 M sodium acetate buffer (pH 5.5). After shaking, the mixture was incubated at room temperature in the dark for 30 min, and then the absorbance was measured at 517 nm using a Shimadzu UV-160A spectrometer, Kyoto (Japan). The IC₅₀ value was determined as the concentration of each sample required to give 50% DPPH• radical scavenging activity.

FRAP test

Acetate buffer (0.3 M, pH 3.6) was prepared by dissolving 3.1 g C₂H₃O₂Na.3H₂O and 16 ml of acetic acid in 1 l of distilled water. TPTZ (2, 4, 6 -tripiryridyl-S-triazine) solution was prepared by dissolving 23.4 mg of TPTZ in 7.5 ml of 40 mM HCl solution. Ferric solution (20 mM) was prepared using FeCl₃.6H₂O. The final working FRAP reagent was prepared freshly by mixing acetate buffer, TPTZ and ferric solutions at a ratio of 10:1:1.

In brief, 900 µl FRAP (ferric reducing-antioxidant power) working reagent was mixed with 90 µl distilled water and was warmed to 37 °C in a water bath. The reagent control reading was recorded at 595 nm, followed by adding 30 µl of sample solutions (100 mg in 10 ml of n-hexane). The absorbance was taken at 595 nm, against the control solution. A standard curve was prepared using different concentrations of FeSO₄.7H₂O (200–2000 µmol/l). All solutions were freshly prepared. The results were expressed in mmol Fe²⁺/l (Capannesi, *et al.*, 2000).

HPLC preparative analysis

F3 methanol solution (1.0 mg/mL) was filtered through cellulose acetate membrane filter (0.45 µm, Anpu Co., Shanghai, China) prior to HPLC analysis. 1 ml aliquot of the filtrate of concentration 3mg ml⁻¹ in methanol was injected into a Wellchrom KNAUER

HPLC, with reverse column (C18, 250 ×4.6 mm i.d., 5 µm), HPLC pump: KNAUER k-1001, UV detector: KNAUER k-2600, and eluted with a mobile phase containing solvent A (methanol) solvent B (acetonitril) and solvent C (water) (90:5:5) iso-critically pumped at a flow rate of 1.0 mL/min. In the chromatogram the peaks of interest were observed at t_R 22.5 min and 27 min, detection was at 250nm. Compounds were collected in a clean, weighed flask from 21.5 to 25 min at room temperature under a room temperature. Multiple preparative HPLC separation was done and fractions for two peaks combined.

Processing of the collected fractions

The solvent was removed from the samples at 20°C with N₂ gas and the residual aqueous phases were shell-frozen on dry ice prior to freeze-drying. The freeze-dried samples were stored at -20°C until NMR analyses were performed. Complete removal of solvent was done by freeze drier and resulted yellow and yellow fade compounds powder.

NMR identification of compounds

NMR spectroscopy experiments on the compounds were performed on a Bruker Avance 500 at 500 MHz for ¹H NMR and ¹³C NMR with CD₃OD and DMSO as solvent in 25 centigrade.

TLC bioautography analysis

An aliquot of AMS methanol solution (1 mg/mL, 3 mL) or individual pure isolate methanol solutions (1.0 mg/mL, 2 mL) was directly deposited (as spots or bands) onto the TLC plates. TLC plates were developed in a presaturated solvent chamber with ethyl acetate- acetonitril- formic acid- water- ethyl formiate (7:1:1:1.1:12) as developing reagents until the solvent front reached 1 cm from the top of plates. The developed TLC plates were then removed from Fig. 1. TLC plates stained with 2.54 mM DPPH• solution in methanol, and visualized (A) under visible light, (B) under UV 254 nm, and (C) under UV 366 nm. Three microlitres Silica gel 60 F254 TLC plates (Merck, Germany) were used for TLC

bioautography analysis.

Result and discussion

DPPH[•] scavenging activity and FRAP tests of isolates

The conventional spectrophotometric DPPH[•] scavenging capacity assay was first used to screen the potential antioxidant of F1, F2, F3 and F4. The DPPH[•] scavenging activities of all the fractions were estimated using the conventional spectrophotometric

DPPH[•] scavenging capacity assay. Table 1 shows the quantities related to the DPPH[•] radical-scavenging activities of the F1, F2, F3 and F4 compare to that of BHT (butylated hydroxy toluen) as an synthetic antioxidant. All fractions except F3 showed DPPH[•] radical-scavenging activities extremely lower than that of BHT (F3=0.006, BHT=0.007, F1=1.47, F2=0.119 and F4=0.24 mg mL⁻¹). F3 showed significant DPPH[•] scavenging activities which were similar to that of BHT.

Table 1. The DPPH radical-scavenging activity (IC₅₀) and ferric reducing-antioxidant power (FRAP) of the total extract and four fractions of *Cumin seed*. Diethylether fraction (F1), dichloromethane fraction (F2), Ethyl acetate fraction (F3) and methanolic fraction (F4).^τ

Antioxidant matter (100 ppm)	IC ₅₀ (mg/mL)	FRAP (mmol Fe ²⁺ /L)
Total extract	0.74 ± 0.10 b	459.46 ± 0.81e
F1	1.47 ± 0.03 a	229.63 ± 0.28 f
F2	0.119 ± 0.003 d	495.52 ± 0.37 d
F3	0.006 ± 0.001 f	521.95 ± 0.91 b
F4	0.24 ± 0.002 e	500.82 ± 0.69 c
BHT	0.007 ± 0.002 f	1955.72 ± 0.98 a

^τ Means ± SD (standard deviation) within a column with the same lowercase letters are not significantly different at P < 0.05

The antioxidant activity decreased as follows: F3 > F2 > F4 > F1

Many similarities were found to exist between the results of the FRAP test and those of the DPPH[•] radical-scavenging activity assay. Table 1 demonstrates that the reducing power of F3 (521.95 mmol/L) was significantly lower than that of BHT and higher than those of total extract (459.46 mmol/L) and other fractions. The chelating abilities of F3 on ferrous ion were good as shown by their low IC₅₀ value (0.006 mg/mL). This indicates that the chelating activity of F3 in metal ion may play an important role in their antioxidant activity.

In the present study the F3 was selected for further purification, since less is known about the antioxidants components in the seeds of *Cumin seed*.

Isolation of antioxidant compounds

F3 fraction was then monitored by a TLC bio-autography method to guide the separation procedure because this method gives quick access for detection and localization of the active compounds in a complicated plant extract (Tasdemiir, *et al.*, 2004). In this

method, the DPPH[•] scavenging activity was observed visually as white yellow spots on a purple background. Fig. 1-A shows the profile of the antioxidant components in the F3 fraction of the methanolic extract of *Cumin seed* under visible light. At least two spots (such as a and b) in the chromatograms were observed to have DPPH[•] scavenging activities. In addition, the same stained TLC plate was also inspected under UV 254 and 366 nm.

Note that the antioxidant spots shown in Fig. 2A were also observed in those of Fig. 2B and C. However, it needs to be pointed out that the different detection sensitivities observed among Fig. 2 A–C were due to the diverse nature of the anti-oxidative compounds. For example, spot (a) was easily better visualized under 254 and 366 nm than that under visible light.

The F3 fraction was directly subjected to a preparative HPLC C18 column and eluted isocratically with methanol, acetonitril and water (90:5:5). Finally, two compounds were separated and identified, the structures of which are Apigenin-7-*O* glucoside and Luteolin- 7- *O* glucoside presented in Fig 2

and 3.




Fig. 1. TLC plates stained with 2.54 mM DPPH_ solution in methanol, and visualized (A) under visible light, (B) under UV 254 nm, and (C) under UV 366 nm. Three microlitres of the 80% methanol extract (AMS, 1 mg/mL) of *Cuminum cyminum* was applied as dots on TLC layer. The spots marked with b and c indicate compounds with DPPH_ scavenging activities.




Fig 2. Apigenin-7-O glucoside




Fig 3. Luteolin- 7- O glucoside

Structure determinations of isolated compounds

The purity and identity of the compounds were established by use of spectroscopic techniques (^1H and ^{13}C NMR) and by comparison with the literature data. Compound 1 was also obtained as a yellow powder, and gave a positive reaction with natural products plus poly ethylen glycole reagent, probably indicating a flavonoid nature. Its UV spectrum was consistent with that of a flavonoid with maxima at 252, 265, and 347 nm (González and Molina, 2006) A direct comparison of ^1H , ^{13}C NMR data (Table 2) with the reported data

(Shen et al., 2004; Li *et al.*, 2007) led to identification of 2 as Luteolin-7-Glucosyde. This is the first report on the isolation of luteolin-7 -O glucoside from Cumin seed.

Compound 2 was obtained as a pale yellow powder, and showed characteristic flavonoid reaction with natural products plus poly ethylen glycole reagent. Its detailed UV (MeOH, λ_{max} 265, 335 nm), ^1H and ^{13}C NMR data (Table 3), negative ESI MS data (m/z 269 [M^- H^-]), agreed well with the previous reported data (Shen et al., 2004; Li *et al.*, 2007). Therefore, this compound was

identified as Apigenin-7-*O* Glucoside, and isolated from Cumin seed for the first time.

Table 2. NMR data of compound 1

Position	¹ H-NMR data		¹³ C-NMR data	
	Compound 1	Ref [23,24]	Compound 1	Ref [23,24]
2	-	-	165.8	165.8
3	6.63,s	6.63,s	103.7	103.7
4	-	-	183.1	183.1
5	-	-	161.8	161.8
6	6.44, d,1.8H ₂	6.44, d,1.8H ₂	100.5	100.5
7	-	-	163.9	163.9
8	6.83,d,J=1.8H ₂	6.83,d,J=1.8H ₂	95.8	95.8
9	-	-	158	158
10	-	-	106.3	106.3
Glucose	-	-	122.6	122.6
1'	7.43,bs	7.43,bs	113.8	113.8
2'	-	-	146.3	146.3
3'	-	-	150.4	150.4
4'	6.95,d,J=8H ₂	6.95,d,J=8H ₂	116.5	116.5
5'	7.41,d,J=8H ₂	7.41,d,J=8H ₂	120.4	120.4
6'	-	-	-	-
1''	5.1,d,J=7.8	5.1,d,J=7.8	100.7	100.7
2''	3.49,t,J=9H ₂	3.49,t,J=9H ₂	73.8	73.8
3''	3.56,t,J=9H ₂	3.56,t,J=9H ₂	76.8	76.8
4''	3.42,t,J=9H ₂	3.42,t,J=9H ₂	70.3	70.3
5''	360,m,	360,m,	77.4	77.4
6''	3.68,dd,J=12.2,5.6 H _a 3.85,dd,J=12.2, 1.8H _b	3.68,dd,J=12.2,5.6 H _a 3.85,dd,J=12.2, 1.8H _b	61.3	61.3

Table 3. NMR data of compound 2

Position	¹ H-NMR data		¹³ C-NMR data	
	Compound 2	Ref[23,24]	Compound 2	Ref[23,24]
Aglycon	-	-	-	-
2	-	-	166.77	164.7
3	6.8, S	6.8,S	104.14	103.4
4	-	-	184.08	182.4
5	-	-	162.93	161.9
6	6.5, S	6.5, d, J=2 Hz	96.07	95
7	-	-	164.81	163.1
8	6.6, S	6.6, d, J=2 Hz	101.19	104.7
9	-	-	158.96	157.3
10	-	-	107.1	105.7
1'	-	-	123.08	121.2
2'	7.9, d, J=8.0 Hz	7.9, d, J=8.8 Hz	129.64	128.9
3'	6.9, d, J=8.5 Hz	6.9, d, J=8.8 Hz	117.05	116.4
4'	-	-	162.93	161.4
5'	6.9, d, J=8.5 Hz	6.9, d, J=8.8 Hz	117.05	116.4
6'	7.9, d, J=8.0 Hz	7.9, d, J=8.8 Hz	129.64	128.9
Glucose	-	-	-	-
1''	5.2, d, J=7H ₂	5.2, d, J=7 Hz	101.64	99.9
2''	3.4 -4	3.4 -4	73.9	73.9
3''	-	-	75.2	75.2
4''	-	-	71	71
5''	-	-	76.1	76.1
6''	-	-	66.7	66.7

References

Amarowicz, R., Naczki, M., Shahidi, F. 2000. Antioxidant activity of various fractions of non-tannin phenolics of Canola hulls, *Journal of Agricultural and Food Chemistry*,48: 2755–2759.

- Behera, S., Nagarajan, S., Jagan, S., M Rao, L. J. 2004. Microwave heating and conventional roasting of cumin seeds (*Cuminum cyminum* L.) and effect on chemical composition of volatiles, *Journal of Food Chemistry*, 87: 25–29.
- Capannesi, C., Palchetti, I., Mascini, M., Parenti, A. 2000. Electrochemical sensor and biosensor for polyphenols detection in olive oils, *Food Chemistry*, 71: 553–562.
- Chauhan, P. S., Satti, N. K., Suri, K. A., Amina, M., Bani, S. 2009. Stimulatory effect of *Cuminum cyminum* and flavonoid glycoside on Cyclosporine- A and restraint stress induced immune- Suppression in Swiss albino mice, *Chemico- Biological Interactions* 2009; 185: 66-72.
- Gachkar, L., Yadegari, D., Rezaei, M. B., Taghizadeh, M. 2011. Chemical and biological characteristics of *Cuminum cyminum* and *Rosmarinus officinalis* essential oils, *Food Chemistry*, 102: 898–904.
- Chung, Y., Chien, C., Teng, K., Chou, S. 2006. Antioxidative and mutagenic properties of *Zanthoxylum ailanthoides* Sieb & zucc, *Food Chemistry*, 97: 418–425.
- Einafshar, S., Purazarang, H., Farhoosh, R., Seyedi, S. M. 2012. Antioxidant activity of the essential oil and methanolic extract of cumin seed (*Cuminum cyminum*), *European Journal of Food Lipid Sciences and Technology*, 114: 168-174.
- Fisch KM, Bo'hm V, Wright AD, Ko' nig GM. 2003. Antioxidative meroterpenoids from the brown alga *Cystoseira crinita*, *Journal of Natural Products*, 66: 968–975.
- Gonza' lez-Lavaut, J. A., Molina-Torres, J. 2006. Flavonoid glycosides from Cuban *Erythroxyllum* species, *Biochemical Systematics and Ecology*, 34: 539–542.
- Guenther, E. 1950. The essential oils. Vol. 4, D. Von Nostrand Company Inc. New York, 615–619.
- Howell, J. C. 1986. Food antioxidants: international perspectives welcome and introductory remarks, *Food and Chemical Toxicology*, 24: 997-1003.
- Ito, N., Hirose, M., Fukushima, S., Tsuda, H. 1986. Studies on antioxidants: their carcinogenic and modifying effects on chemical carcinogenesis, *Food and Chemical Toxicology*, 24: 1071–1082.
- Li, H. J., Luo, Y. G., He, Z.H., Zhang, G. L. 2007. Chemical constituents from *Lonicera saccata*, *Chinese Journal of Applied and Environmental Biology*, 13:188–191.
- Li, X. M., Tian, S. L., Pang, Z. C., Shi, J. Y. 2009. Extraction of *Cuminum cyminum* essential oil by combination technology of organic solvent with low boiling point and steam distillation, *Food Chemistry*; 115: 1114–1119.
- Nakamura Y, Watanabe S, Miyake N, Kohno H, Osawa T. 2003. Dihydrochalcones: evaluation as novel radical scavenging antioxidants, *Journal of Agricultural and Food Chemistry*, 51: 3309–3312.
- Otsuka, H. 2006. Crystallization in Final stages of purification, In: Methods in Biotechnology 20" Natural products isolation" second Ed. Satyait D. Sarker Zahid Latif, Alexander I.Gray (eds.). Human press, New Jersi, 114-126.
- Satti, N. K., Amina, M., Dutti, P., Sharma, B. 2009. A simple and reliable preparative high performance liquid chromatographic technique for isolation of a bioactive flavones diglycoside from and extract of *Cuminum cyminum* seeds, *Acta chromatographica*, 21, 499-512.
- Shen, J., Liang, J., Peng, S. L., Ding, L. S. 2004. Chemical constituents from *Saussurea stella*, *Natural Product Research and Development*. 16: 391–394.
- Shobana, S., Akhilender, K. N. 2000. Antioxidant activity of selected Indian spices, *Prostaglandins Leukotrienes and Essential Fatty Acids*, 62: 107–110.
- Shon, M. Y., Kim, T. H., Sung, N. J. 2003. Antioxidants and free radical scavenging activity of *Phellinus baumii* (*Phellinus* of *Hymenochaetaceae*) extracts *Food Chemistry*, 82: 593–597.
- Tasdemir, D., Donmez, A. A., Calis, I., Ruedi, P. 2004. Evaluation of biological activity of Turkish plants. Rapid screening for the antimicrobial, antioxidant, and acetylcholinesterase inhibiting potential by TLC bioautographic method, *Pharmaceutical Biology*, 42: 374–383.
- Valentao, P., Fernandes, E., Carvalho, F., Andrade, P. B. 2002. Antioxidative properties of Cardoon (*Cynara cardunculus* L.) infusion against superoxide radical, hydroxyl radical, and hypochlorous acid, *Journal of Agricultural and Food Chemistry*, 50: 4989–4993.
- Vekiari, S. A., Oreopoulou, V., Tzia, C., 1993. Thomopoulos CD. Oregano Flavonoids as lipid Antioxidants, *Journal American Oil Chemistry Society*, 70: 483–489.
- Winton, A. L., Winton, K. B. 1939. The structure and composition of foods. Vol. 4, Wiley, New York, 420–423.

جداسازی و شناسایی ترکیبات دارای خواص آنتی‌اکسیدانی
زیره سبز (*Cuminum cyminum*)سودابه عین‌افشار^{1*} - هاشم پورآذرننگ²، رضا فرهوش²، جواد اصیلی³

تاریخ دریافت: 1394/11/05

تاریخ پذیرش: 1395/05/26

چکیده

مواد طبیعی ترکیبات پیچیده‌ای هستند که استخراج سریع مواد فعال آنتی‌اکسیدانی از اهمیت خاصی برخوردار است. برای جداسازی چهار جزء (F1، F2، F3 و F4) موجود در عصاره متانولی زیره سبز از حلال‌هایی با قطبیت متفاوت استفاده شد. خواص آنتی‌اکسیدانی اجزاء فوق، با استفاده از آزمون تعیین قدرت‌گیرندگی رادیکال و آزمون FRAP تعیین شد. جزء F3 با (میلی-گرم بر میلی لیتر) IC500/006 = و (میلی مول آهن 2+ در لیتر) 521/95 = FRAP فعال‌ترین جزء بود. با استفاده از کروماتوگرافی لایه نازک (TLC) و ارزیابی قدرت‌گیرندگی رادیکال آزاد (DPPH) دو ترکیب دارای خاصیت آنتی‌اکسیدانی از جزء F3 استخراج شد. به‌منظور تعیین ترکیب شیمیایی این دو جزء با استفاده از کروماتوگرافی تهیه‌ای (HPLC preparative) به همراه فاز متحرک مناسب (استونیتریل: متانول / آب) مقدار لازم از این دو ترکیب استخراج شد و با استفاده از HNMR و CNMR و مقایسه با منابع موجود، دو ترکیب لوتئولین 7 گلوکوزید و آپیزنین 7 گلوکوزید شناسایی گردیدند.

واژه‌های کلیدی: آنتی‌اکسیدان، آپیزنین، زیره سبز، لوتئولین.

1- استادیار، بخش تحقیقات فنی و مهندسی کشاورزی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی خراسان رضوی، سازمان تحقیقات، آموزش و ترویج کشاورزی، مشهد، ایران.

2- استاد، گروه علوم و صنایع غذایی، دانشکده کشاورزی، دانشگاه فردوسی مشهد.

3- عضو هیئت علمی گروه فارماکوتوزی، دانشکده داروسازی دانشگاه علوم پزشکی مشهد.

* - نویسنده مسئول : (Email: Soodabeheyn@yahoo.com)