



## Water relations, pigment stabilization, photosynthetic abilities and growth improvement in salt stressed rice plants treated with exogenous potassium nitrate application

Suriyan Cha-um<sup>a,\*</sup>, Kongake Siringam<sup>b</sup>, Niran Juntawong<sup>b</sup>,  
Chalermopol Kirdmanee<sup>a</sup>

<sup>a</sup>National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), 113 Thailand Science Park, Paholyothin Road, Klong 1, Klong Luang, Pathumthani 12120, Thailand.

<sup>b</sup>Department of Botany, Faculty of Science, Kasetsart University, Bangkok 10900, Thailand.

\*Corresponding author. E-mail: suriyanc@biotec.or.th

Received 27 June 2009; Accepted after revision 9 May 2010; Published online 14 June 2010

---

### Abstract

Potassium is a major nutrient which may play an important role in many processes such as ion homeostasis in plant cells and osmotic adjustment of guard cells during stomatal opening and closing. Pathumthani 1 (PT1) rice has been reported as being a salt sensitive cultivar and has been selected as a model plant in this study to investigate the possibility of improving the osmotic potential, pigment stabilization, photosynthetic efficiency and growth characteristics of this plant under salinity stress by potassium nitrate ( $\text{KNO}_3$ ) application. Sodium ion accumulation in root and leaves of  $\text{KNO}_3$  (11.8 mM) treated plants declined as compared to the control plants. In control plants, however, due to decline in potassium ion content, the  $\text{Na}^+ / \text{K}^+$  ratio increased significantly. A positive relation between  $\text{Na}^+$  accumulation and osmotic potential was found. Osmotic potential ( $\Psi_s$ ) in the root and leaf tissues of PT1 rice treated with 11.8 mM  $\text{KNO}_3$  was maintained at low levels, which was accompanied with the stabilization of photosynthetic pigments, high photosynthetic performance and better growth characters under 200 mM NaCl. The photosynthetic ability in plants with  $\text{KNO}_3$  application was positively related to plant dry weight. Exogenous  $\text{KNO}_3$  application to rice crops may play a vital role as a short-gun technique for the improvement of salt tolerance.

**Keywords:** Chlorophyll fluorescence; Net photosynthetic rate; *Oryza sativa*; Osmotic potential; Potassium; Quantum yield; Salinity.

---

### Introduction

Soil salinity is a serious abiotic stress, directly and indirectly affecting plant growth and development, leading to loss of productivity especially in glycophyte species (Hasegawa et al., 2000; Qadir et al., 2008). Sodium ion ( $\text{Na}^+$ ) is a major contaminant of salt affected soil

and has been reported as being toxic to plants which is quickly absorbed and taken up by root cells (Tester and Davenport, 2003; Malagoli et al., 2008).

One of the most important salt defense mechanisms in higher plants, especially halophyte species is  $\text{Na}^+/\text{K}^+$  homeostasis, which is defined as a primary defense response when plants are exposed to salt stress (Cuin et al., 2003; Munns and Tester, 2008). There are many documents reporting on this issue in terms of  $\text{Na}^+ / \text{K}^+$  ratio,  $\text{Na}^+$  and  $\text{K}^+$  interactions, potassium transporter proteins and the potassium transport gene family (Schachtman and Liu, 1999; Rodríguez-Navarro, 2006; Amrutha et al., 2007; Chen et al., 2007; Gierth and Mäser, 2007; Britto and Kronzucker, 2008; Cuin et al., 2008; Alemán et al., 2009; Szczerba et al., 2009). The role of potassium in salt tolerance mechanisms has been investigated, including ion homeostasis, osmoregulation and antioxidant systems (Cakmak, 2005; Chen et al., 2007; Szczerba et al., 2009). There are several factors influencing  $\text{Na}^+/\text{K}^+$  homeostasis, such as different genetic resources (Golldack et al., 2003; Kader et al., 2006; Huang et al., 2008), mutant lines (Wu et al., 1996; Zhu et al., 1998) and over-expression of potassium-related gene (s) (Rubio et al., 1999; Obata et al., 2007; Takahashi et al., 2007; Mangano et al., 2008). In addition,  $\text{K}^+$  is known to function in osmotic adjustment in the guard cell controlling the stomatal movements and thus  $\text{CO}_2$  assimilation in photosynthesis (Chartzoulakis et al., 2006; Degl'Innocenti et al., 2009).

Potassium application has been reported as being effective in salt tolerance mechanism of plants through  $\text{Na}^+/\text{K}^+$  homeostasis (Chen et al., 2007; Cuin et al., 2008; Kader and Lindberg, 2008; Alemán et al., 2009), osmoregulation (Szczerba et al., 2009) and antioxidant systems (Cakmak, 2005).  $\text{K}^+$  in plant tissues evidently decreases when plants are exposed to salt stress, especially rice genotypes (Basu et al., 2002; Castillo et al., 2007; Ahmad et al., 2007). Exogenous potassium in the culture medium is directly absorbed and taken up by the root tissues and translocated into the whole plant body (Akram et al., 2007; Akram et al., 2009; Kaya et al., 2007; Shaibur et al., 2008; Zheng et al., 2008). An alternative way to improve salt tolerance in plants is to increase the level of endogenous potassium by the application of potassium salts (Ahmad and Jabeen, 2005; Kaya et al., 2007; Zheng et al., 2008; Akram et al., 2009). Thus growth parameters including plant height and dry weight in salt stressed wheat pretreated with 16 mM  $\text{KNO}_3$  were improved in both salt tolerant (DK961) and salt sensitive (JN17) cultivars (Zheng et al., 2008). Improved growth characters, in terms of shoot dry weight, root dry weight and fruit weight, were also found in salt stressed melons treated with 5 mM  $\text{KNO}_3$  (Kaya et al., 2007). Leaf area and fruit yield of salt stressed *Lagenaria siceraria* were improved by 2.47 mM  $\text{KNO}_3$  (Ahmad and Jabeen, 2005).

Rice (*Oryza sativa* L. spp. *indica*) is the main cereal crop in Asian countries. It plays a major role as a staple food, representing 50-80% of people's daily calorie intake (Khush, 2005). In previous reports, rice was found to be highly sensitive to salt stress in both vegetative and reproductive stages (Zeng et al., 2001; Moradi and Ismail, 2007), leading to crop yield losses of more than 50% when exposed to saline media with electrical conductivity of  $6.65 \text{ dS m}^{-1}$  (Zeng and Shannon, 2000). In Thailand, the Pathumthani 1 (PT1) cultivar is a long grain, aromatic rice with high cooking quality and a soft texture (Laohakunjit and Kerdchoechuen, 2007). It is widely cultivated in irrigated paddy fields and has been reported as being salt susceptible (Cha-um et al., 2007). The objective of this study was to investigate the ameliorative effects of potassium nitrate application on some

physiological responses such as water balance, ion homeostasis and photosynthetic parameters of salt-stressed rice plants.

## Materials and Methods

### *Plant materials and treatments*

Seeds of the salt-sensitive rice cultivar PT1 (*Oryza sativa* L. spp. *indica* cv. Pathumthani 1), provided by Pathumthani Rice Research Center, (Rice Research Institute, Department of Agriculture, Ministry of Agriculture and Cooperative, Thailand) were manually de-husked, surface-sterilized first in 5% Clorox<sup>®</sup> for 60 min, followed by 30% Clorox<sup>®</sup> for 30 min, and then rinsed three times with sterile distilled water. Surface-sterilized seeds were germinated on 0.25% Phytigel<sup>®</sup>-solidified MS media with 3% sucrose (photomixotrophic condition) in a 250 mL glass vessel. All media were adjusted to pH 5.7 and autoclaved. Rice seedlings were cultured *in vitro* under conditions of 25±2 °C ambient temperature, 60±5% relative humidity (RH) and 60±5 μ mol m<sup>-2</sup> s<sup>-1</sup> photosynthetic photon flux density (PPFD) provided by fluorescent lamps with 16 h d<sup>-1</sup> photoperiod. Seven-day-old seedlings were aseptically transferred to MS-liquid sugar-free media (photoautotrophic conditions) with 0, 9.4 (control), 11.8 and 24.1 mM KNO<sub>3</sub> using vermiculite as supporting material. The uncovered vessels containing photoautotrophic seedlings were transferred aseptically to a culture box chamber (Carry Box Model P-850, size 26×36×19 cm, Japan) with RH maintained at 65±5% by 1.5 L saturated NaCl solution (360 gL<sup>-1</sup>). The number of air exchanges in the culture box chambers was increased to 5.1±0.3 h<sup>-1</sup> by perforating the sides of the plastic chamber with 32 holes and placing gas-permeable microporous polypropylene film (0.22 μ m pore size) over each hole (Cha-um et al., 2004). The chamber containing the rice seedlings was acclimated for 14 days in a Plant Growth Incubator under a temperature shift of 28±2 °C/25±2 °C (light/dark), 500±100 μ mol mol<sup>-1</sup> CO<sub>2</sub> concentration, 60±5% RH, 120±5 μ mol m<sup>-2</sup> s<sup>-1</sup> PPFD provided by fluorescent lamps with 12 h d<sup>-1</sup> photoperiod. The culture media were adjusted to 200 mM NaCl (salt stress) for 4 days. Osmotic potential, photosynthetic pigments, photosynthetic performance and growth characters were analyzed.

### *Analytical methods and data collection*

One hundred milligrams of whole plant materials were ground in liquid nitrogen. Sodium and potassium ions in plant materials were extracted by acidic methods (HNO<sub>3</sub> and HClO<sub>4</sub>) and assayed according to Dionisio-Sese and Tobita (1998) using an Atomic Absorption Spectrophotometer (AA, Model M6, Thermo Elemental, MA, USA).

Osmolarity in the roots and leaves of rice seedlings was measured, according to Lanfermeijer et al. (1991). A hundred milligrams each of fresh root and leaf tissue were cut into small pieces, transferred to 1.5 mL micro tube then stirred with a glass rod. Twenty micro liters of extracted solution was dropped directly onto a disc-shaped filter paper in an osmometer chamber (Wescor, USA). The osmolarity was then measured.

Chlorophyll a (Chl<sub>a</sub>), chlorophyll b (Chl<sub>b</sub>) and total chlorophyll (TC) concentrations were determined following the method of Shabala et al. (1998) and the total carotenoids (C<sub>x+c</sub>) concentration was determined following the method of Lichtenthaler (1987). One hundred milligrams of leaf material was collected from the second and third nodes of the shoot tip. The leaf samples were placed in a 25 mL glass vial, along with 10 mL 95.5% acetone, and blended using a homogenizer. The glass vials were sealed with parafilm to prevent evaporation and then stored at 4 °C for 48 h. The Chl<sub>a</sub> and Chl<sub>b</sub> concentrations were measured using a UV-visible spectrophotometer at 644 nm and 662 nm. The C<sub>x+c</sub> concentration was also measured by spectrophotometer at 470 nm. A solution of 95.5% acetone was used as a blank.

The chlorophyll fluorescence emission from the adaxial surface of the third leaf from the shoot tip was monitored using a fluorescence monitoring system in the pulse amplitude modulation mode, as previously described by Loggini et al. (1999). A leaf, adapted to dark conditions for 30 min using leaf-clips, was initially exposed to the modulated measuring beam of far-red light (LED source with typical peak at wavelength 735 nm). Original (F<sub>0</sub>) and maximum (F<sub>m</sub>) fluorescence yields were measured under weak modulated red light (<0.5 μ mol m<sup>-2</sup> s<sup>-1</sup>) with 1.6 s pulses of saturating light (>6.8 μ mol m<sup>-2</sup> s<sup>-1</sup> PAR) and autocalculated using FMS software for Windows<sup>®</sup>. The variable fluorescence yield (F<sub>v</sub>) was calculated by the equation of F<sub>m</sub>-F<sub>0</sub>. The ratio of variable to maximum fluorescence (F<sub>v</sub>/F<sub>m</sub>) was calculated as maximum quantum yield of PSII photochemistry. The photon yield of PSII (Φ<sub>PSII</sub>) in the light was calculated by Φ<sub>PSII</sub> = (F<sub>m</sub>'-F)/F<sub>m</sub>' after 45 s of illumination, when steady state was achieved.

Net photosynthetic rate (P<sub>n</sub>) was calculated by comparing the CO<sub>2</sub> concentration inside the glass vessel containing the rice seedlings with that of outside. CO<sub>2</sub> concentrations inside and outside the glass vessel (C<sub>in</sub> and C<sub>out</sub>) at steady state were measured by Gas Chromatography (GC; Model GC-17A, Shimadzu Co. Ltd., Japan). The GC capillary column and detector were a GS-Q (J&W Scientific<sup>®</sup>, Germany) and a thermal conductivity detector (TCD), respectively. The detector and injector were set to a temperature of 250 °C. The temperature of the column was set at 30 °C for 1 min at initial state, then increased by 20 °C per min to 100 °C, and held for 1 min. The net photosynthetic rate of *in vitro* cultivated plantlets was calculated according to the method of Fujiwara et al. (1987), as follows:

$$[P_n] = K \times E \times V (C_{out} - C_{in}) / \text{Leaf area}$$

where K is a conversion factor converting the amount of CO<sub>2</sub> from volume to mole (40.5 mol m<sup>-3</sup> at 28 °C), E is the number of air exchanges per hour (2.32 h<sup>-1</sup>) and V is the volume of air in the vessel (0.0025 m<sup>3</sup>).

Shoot fresh weight, root fresh weight, shoot dry weight, root dry weight and leaf area of rice seedlings were measured as described by Cha-um et al. (2006). Rice seedlings were dried in a hot-air oven (Mettler, Model 500, Germany) for 2 days, and then incubated in a desiccator before measurement of dry weight. The leaf area of rice seedlings was measured using a Leaf Area Meter DT-scan (Delta-Scan Version 2.03, Delta-T Devices, Ltd., UK).

### Experimental design

The experiment was arranged as a Completely Randomized Design (CRD) with six replicates ( $n=6$ ). The mean values obtained were compared by Least Significant Difference (LSD) and analyzed using SPSS software. The correlations between physiological and biochemical parameters were evaluated using Pearson's correlation coefficients.

### Results and Discussion

Sodium ions ( $\text{Na}^+$ ) in the salt stressed root and leaf tissues of rice seedlings without  $\text{KNO}_3$  treatment were accumulated to 18.26 and 33.66  $\text{mg g}^{-1}\text{FW}$ , whereas those in the 11.8  $\text{KNO}_3$  pretreatment were lowest at 3.64 and 15.70  $\text{mg g}^{-1}\text{FW}$ , respectively (Table 1). In contrast, potassium ions in the root and leaf tissues were dropped to 2.96 and 4.80  $\text{mg g}^{-1}\text{FW}$  when exposed to 200 mM NaCl, leading to enhanced  $\text{Na}^+ / \text{K}^+$  ratio in the plant tissues (Table 1). Potassium content in root and leaf tissues of PT1 salt sensitive rice was unaltered, whereas it was enriched in HJ salt tolerant rice when exposed to salt stress (Siringam et al., 2009). It is possible that  $\text{K}^+$  accumulation may play a key role in salt tolerance mechanisms in rice. The  $\text{K}^+$  was enriched in root and leaf tissues of salt stressed seedlings pretreated with 11.8 mM  $\text{KNO}_3$ , controlling the low level of  $\text{Na}^+$  in salt-stressed seedlings. In olive,  $\text{Na}^+$  content in the leaf tissues of the exogenously applied plants with potassium salts is lower than those without K treatment when exposed to 100 mM NaCl, whereas  $\text{K}^+$  was accumulated (Chartzoulakis et al., 2006). Similarly, the  $\text{Na}^+$  in wild barley (*Hordeum maritimum* L.) treated with 3 mM  $\text{K}^+$  subsequently exposed to 100 mM NaCl is lowest, while  $\text{K}^+$  is highest (Degl'Innocenti et al., 2009). The  $\text{Na}^+$  reduction and  $\text{K}^+$  accumulation in the salt stressed plants pretreated with potassium is directly reduced the  $\text{Na}^+ / \text{K}^+$  ratios, which are reported in wild barley (Degl'Innocenti et al., 2009), olive (Chartzoulakis et al., 2006), *Arabidopsis* (Kaddour et al., 2009), melon (Kaya et al., 2007), sunflower (Akram et al., 2009) and winter wheat (Zheng et al., 2008). In the present study, exogenous application of  $\text{KNO}_3$  was adopted to improve salt tolerance of PT1 rice plants exposed to 200 mM NaCl through the increase of the internal potassium levels.

Table 1. Sodium ( $\text{Na}^+$ ), potassium ( $\text{K}^+$ ) ions and Na:K ratios in the root and leaf tissues of PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $\text{KNO}_3$  subsequently grown under 200 mM NaCl salt stress for 4 days.

$\text{KNO}_3$ (mM)	Root			Leaf		
	$\text{Na}^+$ ( $\text{mg g}^{-1}\text{FW}$ )	$\text{K}^+$ ( $\text{mg g}^{-1}\text{FW}$ )	Na:K	$\text{Na}^+$ ( $\text{mg g}^{-1}\text{FW}$ )	$\text{K}^+$ ( $\text{mg g}^{-1}\text{FW}$ )	Na:K
0	18.26 <sup>a</sup>	2.96 <sup>d</sup>	6.16 <sup>a</sup>	33.66 <sup>a</sup>	4.80 <sup>d</sup>	7.08 <sup>a</sup>
9.4	12.92 <sup>b</sup>	4.17 <sup>c</sup>	3.10 <sup>b</sup>	25.11 <sup>b</sup>	7.43 <sup>c</sup>	3.38 <sup>b</sup>
11.8	3.64 <sup>d</sup>	5.62 <sup>a</sup>	0.65 <sup>d</sup>	15.70 <sup>c</sup>	12.93 <sup>a</sup>	1.22 <sup>c</sup>
24.1	5.94 <sup>c</sup>	4.96 <sup>b</sup>	1.21 <sup>c</sup>	17.34 <sup>c</sup>	10.96 <sup>b</sup>	1.58 <sup>c</sup>
ANOVA	**	**	**	**	**	**

Different letters in each column show significant difference at  $P \leq 0.01$  (\*\*) by Least Significant Difference (LSD).

$\text{Na}^+$  accumulation in the salt-stressed root (Figure 1 A) and leaf tissues (Figure 1 B) was positively correlated to osmotic potential ( $\Psi_s$ ), with  $r^2 = 0.99$  for both tissues. Also, the  $\Psi_s$  in the leaf tissues was higher than in the root tissues when plants were exposed to 200 mM

NaCl (Figure 1). A major role of K in plant cells is osmoregulation, which is marked by the status of osmotic potential ( $\Psi_s$ ). The  $\Psi_s$  in both root and leaf tissues of salt stressed PT1 seedlings under the  $\text{KNO}_3$  treatments was better than those without  $\text{KNO}_3$  treatment. Improvement of the plant water relations including water content, water use efficiency, relative water content and water potential in K pretreated plants exposed to salt stress has been reported (Kaya et al., 2007; Zheng et al., 2008; Akram et al., 2009; Kaddour et al., 2009). The reduction of  $\Psi_s$  in the salt stressed leaves was positively correlated to total chlorophyll content ( $r^2 = 0.96$ ) (Figure 2). Chlorophyll a ( $\text{Chl}_a$ ), chlorophyll b ( $\text{Chl}_b$ ), total chlorophyll (TC) and total carotenoids ( $C_{x+c}$ ) concentrations in salt stressed leaves of PT1 rice seedlings were at 167.02, 53.72, 220.74 and 82.20  $\mu\text{g g}^{-1}$  FW in 11.8 mM  $\text{KNO}_3$  treated seedlings which were greater than those without  $\text{KNO}_3$  by 2.49, 2.49, 2.49 and 2.54 times, respectively (Table 2). The chlorophyll content in salt stressed wheat pretreated with 16 mM  $\text{KNO}_3$  is maintained (Zheng et al., 2008). Also, the photosynthetic pigments,  $\text{Chl}_a$ ,  $\text{Chl}_b$  and TC in salt stressed melon plants pretreated with 5 mM  $\text{KNO}_3$  are stabilized better than those without  $\text{KNO}_3$  (Kaya et al., 2007). The  $\text{Chl}_a$  content in salt stressed leaves of PT1 rice was positively correlated with maximum quantum efficiency of PSII ( $F_v/F_m$ ) with  $r^2 = 0.89$  (Figure 3). The  $F_v/F_m$ , photon yield of PSII ( $\Phi_{\text{PSII}}$ ) and net photosynthetic rate ( $P_n$ ) of rice seedlings exposed to NaCl stress (200 mM) were greatest under 11.8 mM  $\text{KNO}_3$  pretreatment. The levels in plants pretreated with 9.4, 11.8 and 24.1 mM  $\text{KNO}_3$  were higher than those in plants without  $\text{KNO}_3$  by 1.77, 2.21 and 10.36 times, respectively (Table 3). Maximum quantum yield of PSII ( $F_v/F_m$ ) in salt stressed sunflowers without K diminished to a greater degree than in plants treated with K, leading to high  $P_n$  (Akram et al., 2009). In the dark reaction of photosynthesis,  $\text{K}^+$  in the guard cells affects  $\text{CO}_2$  assimilation through controlling stomatal movements. Proper stomatal function in salt stressed leaves treated with exogenous potassium may be maintained by enriched  $\text{K}^+$  in the cells, leading to high  $P_n$  (Chartzoulakis et al., 2006; Akram et al., 2009; Degl'Innocenti et al., 2009). This is in congruence with the present results that show high  $P_n$  in salt stressed PT1 pretreated with 11.8 mM  $\text{KNO}_3$ , as well as improved growth when compared to plants without pretreatment of  $\text{KNO}_3$ . A positive relationship between  $\Phi_{\text{PSII}}$  and  $P_n$  was found ( $r^2 = 0.99$ ) (Figure 4). In addition, physiological parameters, including  $\text{Chl}_a$ ,  $\text{Chl}_b$ , TC,  $C_{x+c}$ ,  $F_v/F_m$ ,  $\Phi_{\text{PSII}}$  and  $P_n$ , were positively correlated to each other to a highly significant ( $P \leq 0.01$ ) level in statistical analysis (Table 4). Photosynthetic activity in rice pretreated with  $\text{KNO}_3$  and subsequently exposed to salt stress was enhanced, leading to high growth rates ( $r^2 = 0.73$ ) (Figure 5). Growth parameters, shoot fresh weight (SFW), root fresh weight (RFW), shoot dry weight (SDW), root dry weight (RDW) and leaf area (LA), were similarly affected in  $\text{KNO}_3$  pretreated plants under salt stress. Growth parameters of salt stressed seedlings under 11.8 mM  $\text{KNO}_3$  application were highest and significantly better than those without  $\text{KNO}_3$  treatment (Table 5).

In conclusion, the exogenous application of 11.8 mM  $\text{KNO}_3$  to PT1 rice seedlings produced optimal effects under 200 mM NaCl stress and led to the better ionic balance, pigment stabilization, reduced chlorophyll fluorescence emission and greater net photosynthetic rate, leading to improved growth performance. A short-gun technique, the exogenous application of  $\text{KNO}_3$  to rice crops is an alternative procedure to be adopted in improving salt tolerance in rice crop grown on salt-affected land.




Figure 1. Relationship between sodium ions ( $\text{Na}^+$ ) and osmotic potential ( $\Psi_s$ ) in root (A) and leaf tissues (B) of PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $\text{KNO}_3$  subsequently exposed to 200 mM  $\text{NaCl}$  salt stress for 4 days. Error bars represent by  $\pm$ SE.




Figure 2. Relationship between osmotic potential ( $\Psi_s$ ) in the leaf tissues and total chlorophyll contents in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $\text{KNO}_3$  subsequently exposed to 200 mM  $\text{NaCl}$  salt stress for 4 days. Error bars represent by  $\pm$ SE.

Table 2. Chlorophyll a ( $\text{Chl}_a$ ), chlorophyll b ( $\text{Chl}_b$ ), total chlorophyll (TC) and total carotenoids ( $\text{C}_{x+c}$ ) concentrations in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $\text{KNO}_3$  subsequently grown under 200 mM  $\text{NaCl}$  salt stress for 4 days.

$\text{KNO}_3$ (mM)	$\text{Chl}_a$ ( $\mu\text{g g}^{-1}$ FW)	$\text{Chl}_b$ ( $\mu\text{g g}^{-1}$ FW)	TC ( $\mu\text{g g}^{-1}$ FW)	$\text{C}_{x+c}$ ( $\mu\text{g g}^{-1}$ FW)
0	67.07 <sup>c</sup>	21.62 <sup>d</sup>	88.69 <sup>b</sup>	32.34 <sup>c</sup>
9.4	90.69 <sup>b</sup>	30.92 <sup>c</sup>	121.61 <sup>b</sup>	43.87 <sup>b</sup>
11.8	167.02 <sup>a</sup>	53.72 <sup>a</sup>	220.74 <sup>a</sup>	82.20 <sup>a</sup>
24.1	155.98 <sup>a</sup>	45.42 <sup>b</sup>	201.40 <sup>a</sup>	73.59 <sup>a</sup>
ANOVA	**	**	**	**

Different letters in each column show significant difference at  $P \leq 0.01$  (\*\*) by Least Significant Difference (LSD).




Figure 3. Relationship between chlorophyll a content and maximum quantum yield of PSII ( $F_v/F_m$ ) in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $KNO_3$  subsequently exposed to 200 mM NaCl salt stress for 4 days. Error bars represent by  $\pm$ SE.

Table 3. Maximum quantum yield of PSII ( $F_v/F_m$ ), photon yield of PSII ( $\Phi_{PSII}$ ) and net photosynthetic rate ( $P_n$ ) in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $mgL^{-1}$   $KNO_3$  subsequently grown under 200 mM NaCl salt stress for 4 days.

$KNO_3$ (mM)	$F_v/F_m$	$\Phi_{PSII}$	$P_n$ ( $\mu mol CO_2 m^{-2} s^{-1}$ )
0	0.501 <sup>d</sup>	0.288 <sup>d</sup>	0.22 <sup>c</sup>
9.4	0.676 <sup>c</sup>	0.389 <sup>c</sup>	0.89 <sup>b</sup>
11.8	0.887 <sup>a</sup>	0.637 <sup>a</sup>	2.28 <sup>a</sup>
24.1	0.770 <sup>b</sup>	0.443 <sup>b</sup>	1.13 <sup>b</sup>
ANOVA	**	**	**

Different letters in each column show significant difference at  $P \leq 0.01$  (\*\*\*) by Least Significant Difference (LSD).




Figure 4. Relationship between photon yield of PSII ( $\Phi_{PSII}$ ) and net photosynthetic rate ( $P_n$ ) in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM  $KNO_3$  subsequently exposed to 200 mM NaCl salt stress for 4 days. Error bars represent by  $\pm$ SE.



Table 4. Relationship between physiological and biochemical parameters of rice seedlings treated with 0, 9.4, 11.8 and 24.1 mM KNO<sub>3</sub> subsequently exposed to 200 mM NaCl for 4 days.

Parameters	Chl <sub>a</sub>	Chl <sub>b</sub>	TC	C <sub>x+c</sub>	F <sub>v</sub> /F <sub>m</sub>	Φ <sub>PSII</sub>	P <sub>n</sub>
Chl <sub>a</sub>	1	-	-	-	-	-	-
Chl <sub>b</sub>	0.976**	1	-	-	-	-	-
TC	0.999**	0.986**	1	-	-	-	-
C <sub>x+c</sub>	0.981**	0.960**	0.980**	1	-	-	-
F <sub>v</sub> /F <sub>m</sub>	0.879**	0.911**	0.890**	0.886**	1	-	-
Φ <sub>PSII</sub>	0.843**	0.886**	0.857**	0.850**	0.928**	1	-
P <sub>n</sub>	0.811**	0.851**	0.824**	0.820**	0.898**	0.958**	1

Significant levels at P≤0.01 is represented by \*\* using Pearson's correlation coefficients.




Figure 5. Relationship between net photosynthetic rate (P<sub>n</sub>) and plant dry weight in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM KNO<sub>3</sub> subsequently exposed to 200 mM NaCl salt stress for 4 days. Error bars represent by ±SE.

Table 5. Shoot fresh weight (SFW), root fresh weight (RFW), shoot dry weight (SDW), root dry weight (RDW) and leaf area (LA) in PT1 rice treated with 0, 9.4, 11.8 and 24.1 mM KNO<sub>3</sub> subsequently grown under 200 mM NaCl salt stress for 4 days.

KNO <sub>3</sub> (mM)	SFW (mg)	RFW (mg)	SDW (mg)	RDW (mg)	LA (cm <sup>2</sup> )
0	145.1 <sup>b</sup>	31.3 <sup>b</sup>	31.7 <sup>b</sup>	3.0 <sup>b</sup>	5.50 <sup>d</sup>
9.4	158.9 <sup>b</sup>	34.4 <sup>ab</sup>	32.6 <sup>b</sup>	3.4 <sup>ab</sup>	7.48 <sup>c</sup>
11.8	181.3 <sup>a</sup>	36.9 <sup>a</sup>	37.1 <sup>a</sup>	3.8 <sup>a</sup>	12.24 <sup>a</sup>
24.1	159.9 <sup>b</sup>	36.2 <sup>a</sup>	32.6 <sup>b</sup>	3.7 <sup>a</sup>	10.55 <sup>b</sup>
ANOVA	*	*	*	*	**

Different letters in each column show significant difference at P≤0.05 (\*) or P≤0.01 (\*\*) by Least Significant Difference (LSD).

### Acknowledgements

The authors wish to thank the National Center for Genetic Engineering and Biotechnology (BIOTEC) as a funding source (Grant number; BT-B-02-RG-BC-4905) and Dr. Teeraporn Busaya-angoon at Pathumthani Rice Research Center, for providing of PT1 rice seeds.

## References

- Ahmad, M.S.A., Javed, F., Ashraf, M., 2007. Iso-osmotic effect of NaCl and PEG on growth, cations and free proline accumulation in callus tissue of two indica rice (*Oryza sativa* L.) genotypes. *Plant Growth Regul.* 53: 53-63.
- Ahmad, R., Jabeen, R., 2005. Foliar spray of mineral elements antagonistic to sodium-a technique to induce salt tolerance in plants growing under saline conditions. *Pak. J. Bot.* 37: 913-920.
- Akram, M.S., Ashraf, M., Akram, N.A., 2009. Effectiveness of potassium sulfate in mitigating salt-induced adverse effects on different physio-biochemical attributes in sunflower (*Helianthus annuus* L.). *Flora*, 204: 471-483.
- Akram, M.S., Athar, H., Ashraf, M., 2007. Improving growth and yield of sunflower (*Helianthus annuus* L.) by foliar application of potassium hydroxide (KOH) under salt stress. *Pak. J. Bot.* 39: 769-776.
- Alemán, F., Nieves-Cordones, M., Martínez, V., Rubio, F., 2009. Potassium/sodium steady-state homeostasis in *Thellungiella halophila* and *Arabidopsis thaliana* under long-term salinity conditions. *Plant Sci.* 176: 768-774.
- Amrutha, R.N., Sekhar, P.N., Varshney, R.K., Kavi Kishor, P.B., 2007. Genome-wide analysis and identification of genes related to potassium transporter families in rice (*Oryza sativa* L.). *Plant Sci.* 172: 708-721.
- Basu, S., Gangopadhyay, G., Mukherjee, B.B., 2002. Salt tolerance in rice *in vitro*: Implication of accumulation of Na<sup>+</sup>, K<sup>+</sup> and proline. *Plant Cell Tiss. Org. Cult.* 69: 55-64.
- Britto, D.T., Kronzucker, H.J., 2008. Cellular mechanisms of potassium transport in plants. *Physiol. Plant.* 113: 637-650.
- Cakmak, I., 2005. The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *J. Plant Nutr. Soil Sci.* 168: 521-530.
- Castillo, E.G., Tuong, T.P., Ismail, A.M., Inubushi, K., 2007. Response to salinity in rice: Comparative effects of osmotic and ionic stresses. *Plant Prod. Sci.* 10: 159-170.
- Chartzoulakis, K., Psarras, G., Vemmos, S., Loupassaki, M., Bertaki, M., 2006. Response of two olive cultivars to salt stress and potassium supplement. *J. Plant Nutri.* 29: 2063-2078.
- Cha-um, S., Supaibulwatana, K., Kirdmanee, C., 2004. Biochemical and physiological responses of Thai jasmine rice (*Oryza sativa* L. ssp. *indica* cv. KDML105) to salt stress. *Sci. Asia*, 30: 247-253.
- Cha-um, S., Supaibulwatana, K., Kirdmanee, C., 2006. Water relation, photosynthetic ability and growth of Thai jasmine rice (*Oryza sativa* L. ssp. *indica* cv. KDML105) to salt stress by application of exogenous glycinebetaine and choline. *J. Agron. Crop Sci.* 192: 25-36.
- Cha-um, S., Vejchasarn, P., Kirdmanee, C., 2007. An effective defensive response in Thai aromatic rice varieties (*Oryza sativa* L. ssp. *indica*) to salinity. *J. Crop Sci. Biotechnol.* 10: 257-264.
- Chen, Z., Pottosin, I.I., Cuin, T.A., Fuglsang, A.T., Tester, M., Jha, D., Zepeda-Jazo, I., Zhou, M., Palmgren, M.G., Newman, I.A., Shabala, S., 2007. Root plasma membrane transporters controlling K<sup>+</sup>/Na<sup>+</sup> homeostasis in salt-stressed barley. *Plant Physiol.* 145: 1714-1725.
- Cuin, T.A., Miller, A.J., Laurie, S.A., Leigh, R.A., 2003. Potassium activities in cell compartments of salt-grown barley leaves. *J. Exp. Bot.* 54: 657-661.
- Cuin, T.A., Betts, S.A., Chalmandrier, R., Shabala, S., 2008. A root's ability to retain K<sup>+</sup> correlates with salt tolerance in wheat. *J. Exp. Bot.* 59: 2697-2760.
- Degl'Innocenti, E., Hafsi, C., Guidi, L., Navari-Izzo, F., 2009. The effect of salinity on photosynthetic activity in potassium-deficient barley species. *J. Plant Physiol.* 166: 1968-1981.
- Dionisio-Sese, M.L., Tobita, S., 1998. Antioxidant responses of rice seedlings to salinity stress. *Plant Sci.* 135: 1-9.
- Fujiwara, K., Kozai, T., Watanabe, I., 1987. Fundamental studies on environment in plant tissue culture vessels. (3) Measurements of carbon dioxide gas concentration in closed vessels containing tissue cultured plantlets and estimates of net-photosynthetic rates of the plantlets. *J. Agric. Method*, 4: 21-30.
- Gierth, M., Mäser, P., 2007. Potassium transporters in plants-Involvement in K<sup>+</sup> acquisition, redistribution and homeostasis. *FEBS Letts.* 581: 2348-2356.
- Gollmack, D., Quigley, F., Michalowski, C.B., Kamasani, U.R., Bohnert, H.J., 2003. Salinity stress-tolerant and-sensitive rice (*Oryza sativa* L.) regulate AKT1-type potassium channel transcripts differently. *Plant Mol. Biol.* 51: 71-81.
- Hasegawa, P.M., Bressan, R.A., Zhu, J.K., Bohnert, H.J., 2000. Plant cellular and molecular responses to high salinity. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 51: 463-499.
- Huang, S., Spielmeier, W., Lagudah, E.S., Munns, R., 2008. Comparative mapping of *HKT* genes in wheat, barley, and rice, key determinants of Na<sup>+</sup> transport, and salt tolerance. *J. Exp. Bot.* 59: 927-937.
- Kaddour, R., Nasri, N., M'rah, S., Berthomieu, P., Lachaal, M., 2009. Comparative effect of potassium on K and Na uptake and transport in two accessions of *Arabidopsis thaliana* during salinity stress. *C.R. Biol.* 332: 784-794.
- Kader, M.A., Lindberg, S., 2008. Cellular traits for sodium tolerance in rice (*Oryza sativa* L.). *Plant Biotechnol.* 25: 247-255.

- Kader, M.D., Seidel, T., Gollmack, D., Lindberg, S., 2006. Expressions of *OsHKT1*, *OsHKT2*, and *OsVHA* are differentially regulated under NaCl stress in salt-sensitive and salt-tolerant rice (*Oryza sativa* L.) cultivars. *J. Exp. Bot.* 57: 4257-4268.
- Kaya, C., Tuna, A.L., Ashraf, M., Altunlu, H., 2007. Improved salt tolerance of melon (*Cucumis melo* L.) by the addition of proline and potassium nitrate. *Environ. Exp. Bot.* 60: 397-403.
- Khush, G.S., 2005. What it will take to feed 5.0 billion rice consumers in 2030. *Plant Mol. Biol.* 59: 1-6.
- Lanfermeijer, F.C., Koerselman-Kooij, J.W., Borstlap, A.C., 1991. Osmosensitivity of sucrose uptake by immature pea cotyledons disappears during development. *Plant Physiol.* 95: 832-838.
- Laohakunjit, N., Kerchoechuen, O., 2007. Aroma enrichment and the change during storage of non-aromatic milled rice coated with extracted natural flavor. *Food Chem.* 101: 339-344.
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods Enzymol.* 148: 350-380.
- Loggini, B., Scartazza, A., Brugnoli, E., Navari-Izzo, F., 1999. Antioxidant defense system, pigment composition, and photosynthetic efficiency in two wheat cultivars subjected to drought. *Plant Physiol.* 119: 1091-1099.
- Malagoli, P., Britto, D.T., Schulze, L.M., Kronzucker, H.J., 2008. Futile Na<sup>+</sup> cycling at the root plasma membrane in rice (*Oryza sativa* L.): kinetics, energetics, and relationship to salinity tolerance. *J. Exp. Bot.* 59: 4109-4117.
- Manganano, S., Silberstein, S., Santa-Maria, M.E., 2008. Point mutations in the barley HvHAK1 potassium transporter lead to improved K<sup>+</sup>-nutrition and enhanced resistance to salt stress. *FEBS Letts.* 582: 3922-3928.
- Moradi, F., Ismail, A.M., 2007. Responses of photosynthesis, chlorophyll fluorescence and ROS-Scavenging systems to salt stress during seedling and reproductive stages in rice. *Ann. Bot.* 99: 1161-1173.
- Munns, R., Tester, M., 2008. Mechanisms of salinity tolerance. *Ann. Rev. Plant Biol.* 59: 651-681.
- Obata, T., Kitamoto, H.K., Nakamura, A., Fukuda, M., Tanaka, Y., 2007. Rice shaker potassium channel OsKAT1 confers tolerance to salinity stress on yeast and rice cells. *Plant Physiol.* 144: 1978-1985.
- Qadir, M., Tabei, A., Akhtar, J., Larbi, A., Minhas, P.S., Khan, M.A., 2008. Productivity enhancement of salt-affected environments through crop diversification. *Land Degrad. Develop.* 19: 429-453.
- Rodríguez-Navarro, A., Rubio, F., 2006. High-affinity potassium and sodium transport systems in plants. *J. Exp. Bot.* 57: 1149-1160.
- Rubio, F., Schwarz, M., Gassmann, W., Schroeder, J.I., 1999. Genetic selection of mutations in the high affinity K<sup>+</sup> transporter HKT1 that define functions of a loop site for reduced Na<sup>+</sup> permeability and increased Na<sup>+</sup> tolerance. *J. Biol. Chem.* 274: 6839-6847.
- Schachtman, D., Liu, W., 1999. Molecular pieces to the puzzle of the interaction between potassium and sodium uptake in plants. *Trend. Plant Sci.* 4: 281-287.
- Shabala, S.N., Shabala, S.I., Martynenko, A.I., Babourina, O., Newman, I.A., 1998. Salinity effect on bioelectric activity, growth, Na<sup>+</sup> accumulation and chlorophyll fluorescence of maize leaves: a comparative survey and prospects for screening. *Aust. J. Plant Physiol.* 25: 609-616.
- Shaibur, M.R., Shamim, A.H.M., Kawai, S., 2008. Growth response of hydroponic rice seedlings at elevated concentrations of potassium chloride. *J. Agric. Rural Develop.* 6: 43-53.
- Siringam, K., Juntawong, N., Cha-um, S., Kirdmanee, C., 2009. Relationships between sodium ion accumulation and physiological characteristics in rice (*Oryza sativa* L. spp. *indica*) seedlings grown under iso-osmotic salinity stress. *Pak. J. Bot.* 41: 1837-1850.
- Szczerba, M.W., Britto, D.T., Kronzucker, H.J., 2009. K<sup>+</sup> transport in plants: Physiology and molecular biology. *J. Plant Physiol.* 166: 447-466.
- Takahashi, R., Liu, S., Takano T., 2007. Cloning and functional comparison of a high-affinity K<sup>+</sup> transporter gene *PhaHKT1* of salt-tolerant and salt-sensitive reed plants. *J. Exp. Bot.* 58: 4387-4395.
- Tester, M., Davenport, R., 2003. Na<sup>+</sup> tolerance and Na<sup>+</sup> transport in higher plants. *Ann. Bot.* 91: 503-527.
- Wu, S.J., Ding, L., Zhu, J.K., 1996. *SOS1*, a genetic locus essential for salt tolerance and potassium acquisition. *Plant Cell*, 8: 617-627.
- Zeng, L., Shannon, M.C., 2000. Salinity effects on seedling growth and yield components of rice. *Crop Sci.* 40: 996-1003.
- Zeng, L., Shannon, M.C., Lesch, S.M., 2001. Timing of salinity stress affects rice growth and yield components. *Agric. Water Manage.* 48: 191-206.
- Zheng, Y., Jia, A., Ning, T., Xu, J., Li, Z., Jiang, G., 2008. Potassium nitrate application alleviates sodium chloride stress in winter wheat cultivars differing in salt tolerance. *J. Plant Physiol.* 165: 1455-1465.
- Zhu, J.K., Liu, J., Xiong, L., 1998. Genetic analysis of salt tolerance in *Arabidopsis*: Evidence for a critical role of potassium nutrition. *Plant Cell*, 10: 1181-1191.

Article reference # ijpp09-207; Editorial responsibility: H.R. Sadeghipour